

- Biológia, Bratislava, 48: 45-47.  
 Wagnerová Z. (1992): Rostlinné spoločenstvo s dominanciou *Petasites albus* v subalpínskom stupni Krkonoš, *Philonoti seriatae-Petasitetum albi as.nova.* - Opera Corcontica, Praha, 29: 73-83.

## NOVÁ LICHENOLOGICKÁ LITERATURA II.

sestavil J.Liška

- Adler M.T. (1990): An artificial key to the genera of the Parmeliaceae (Lichenes, Ascomycotina). - *Mycotaxon* 38: 331-347.
- Aguirre-Hudson B. (1991): A taxonomic study of the species referred to the ascomycete genus *Leptorhaphis*. - *Bull. Brit. Mus. Nat. Hist., Bot.*, 21/2: 85-192.
- Ahmadjian V. (1991): Molecular biology of lichens: a look to the future. - *Symbiosis* 11: 249-254.
- Ahti T., ed. (1990): William Nylander's collected lichenological papers. Vols. I-III. - 2144 p., J.Cramer, Berlin & Stuttgart.
- Aptroot A. (1991): A conspectus of Normandina (Verrucariaceae, lichenized Ascomycetes). - *Willdenowia* 21: 263-267.
- Aptroot A. (1991): A monograph of the Pyrenulaceae (excluding Anthracothecium and Pyrenula) and the Requinellaceae, with notes on the Pleomassariaceae, the Trypetheliaceae and Mycomicrothelia (lichenized and non-lichenized Ascomycetes). - *Biblioth. Lichenol.* 44: 1-178.
- Aptroot A. (1991): Lichenologische literature 1987-1990. - *Buxbaumiella* 26: 45-49.
- Armaleo D. & Clerc P. (1991): Lichen chimeras: DNA analysis suggests that one fungus forms two morphotypes. - *Exp. Mycol.* 15: 1-10.
- Armstrong R.A. (1992): Soroidal dispersal from individual soralia in the lichen *Hypogymnia physodes* (L.) Nyl. - *Environm. Experiment. Bot.* 32: 55-63.
- Baral H.O. (1992): Vital versus herbarium taxonomy: morphological differences between living and dead cells of ascomycetes, and their taxonomic implications. - *Mycotaxon* 44: 333-390.
- Bates J.W. & Farmer A.M., eds. (1992): Bryophytes and lichens in a changing environment. - 404 p., Oxford Univ. Press, Oxford.
- Boom P.P.G. van den (1992): The saxicolous species of the lichen genus *Lecania* in The Netherlands, Belgium and Luxembourg. - *Nova Hedwigia* 54: 229-254.
- Brasch J. & Jacobsen P. (1991): Flechten und ihre Allergene. - *Allergologie* 14: 99-103.
- Bruss O. (1990): Studien über die Flechtengattung *Catapyrenium* (Verrucariaceae). I. Die Gattung *Catapyrenium* in Europa - *Ergänzungen.* - *Linz. Biol. Beitr.* 22: 327-334.
- Brown D.H. (1991): Lichen mineral studies - currently clarified or confused? - *Symbiosis* 11: 207-223.
- Brown D.H. (1992): Impact of agriculture on bryophytes and lichens in a changing environment. - In: Bates J.W. & Farmer A.M., eds., *Bryophytes and lichens in a changing environment*, p. 259-283, Oxford Univ. Press, Oxford.
- Brown D.H. & Brown R.M. (1991): Mineral cycling and lichens: the physiological basis. - *Lichenologist* 23: 293-307.
- Cieslinski S. & Czerwowska K. (1991): Threatened lichens in Poland and their conservation. - *Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich* 106: 133-149.
- Cieslinski S., Czerwowska K. & Fabiszewski J. (1992): Czerwona lista porostów zagrożonych w Polsce. - In: Zarzycki K., Wojewoda W. & Heinrich Z., eds., *Lista*

- roślin zagrożonych w Polsce (ed. 2), p. 57-74, Inst. Bot. W.Szafera PAN, Kraków.
- Cielinski S., Czyżewska K. & Glanc K. (1992): Resources of lichens at the Białowieża National Park (N.E. Poland) and their changes. - Veröff. Geobot. Inst. ETH, Stiftung Rübel, Zürich 107: 392-401.
- Crittenden P.D. & Porter N. (1991): Lichen-forming fungi: potential sources of novel metabolites. - Trends in Biotechnology 9: 409-414.
- Eriksson O.E. & Hawksworth D.L. (1991): Systema Ascomycetum Vol. 9, Parts 1-2. - 271 p., Kew & Umea.
- Eriksson O.E. & Hawksworth D.L. (1991): Systema Ascomycetum Vol. 10, Parts 1-2. - 167 p., Kew & Umea.
- Esslinger T.L. (1989): Systematics of Oropogon (Alectoriaceae) in the New World. - Syst. Bot. Monogr. 28: 1-111.
- Esslinger T.L. (1992): The brown Parmelia type specimens of A.N.Oxner. - Lichenologist 24: 13-20.
- Faltnowicz W. (1992): The lichens of Western Pomerania (NW Poland). An ecogeographical study. - Polish Bot. Stud. 4: 1-182.
- Gams W. (1991): What are names in current use? - Mycotaxon 40: 319-322.
- Garty J. (1992): The postfire recovery of rock-inhabiting algae, microfungi, and lichens. - Can. J. Bot. 70: 301-312.
- Geppert H. & Stordeur R. (1991): Ein Beitrag zur Kenntnis der Flechtenflora des Erzgebirges. 1. Mitteilung. - Wiss. Zeitschr. Univ. Halle 40/6: 69-92.
- Hafellner J. (1991): Die Gattung *Aspicilia*, ihre Ableitungen nebst Bemerkungen über cryptolecanorine Ascocarpororganisation bei anderen Genera der Lecanorales (*Ascomycetes lichensisati*). - Acta Bot. Malacit. 16: 133-140.
- Hafellner J. (1992): A new checklist of lichenized and lichenicolous fungi of the Madeira Archipelago. - 29 p., Inst. Bot. Karl-Franzens-Univ., Graz.
- Hallingbäck T. (1989): Occurrence and ecology of the lichen *Lobaria scrobiculata* in southern Sweden. - Lichenologist 21: 331-341.
- Hauck M. (1992): Rote Liste der gefährdeten Flechten in Niedersachsen und Bremen. - Informationsdienst Naturschutz Niedersachsen 12/1: 1-44.
- Hawksworth D.L. (1991): Lists of names in current use: a new initiative to address a continuing problem. - Mycotaxon 40: 445-468.
- Herzig R. & Urech M. (1991): Flechten als Bioindikatoren. Integriertes biologisches Messsystem der Luftverschmutzung für das Schweizer Mittelland. - Biblioth. Lichenol. 43: 1-283.
- Hestmark G. (1990): Thalloconidia in the genus *Umbilicaria*. - Nordic J. Bot. 9: 547-574.
- Hestmark G. (1991): Teleomorph-anamorph relationships in *Umbilicaria* I. Making the connections. - Lichenologists 23: 343-359.
- Hestmark G. (1991): Teleomorph-anamorph relationships in *Umbilicaria* II. Patterns in propagative morph production. - Lichenologist 23: 361-380.
- Hestmark G. (1991): To sex, or not to sex... Structures and strategies of reproduction in the family Umbilicariaceae (Lecanorales, Ascomycetes). - Sommerfeltia, Suppl. 3: 1-47.
- Hill D.J. (1992): Lobe growth in lichen thalli. - Symbiosis 12: 43-55.
- Honegger R. (1991): Functional aspects of the lichen symbiosis. - Ann. Rev. Plant Physiol. Plant Molecul. Biol. 42: 553-578.
- Honegger R. (1991): Fungal evolution: symbiosis and morphogenesis. - In: Margulis L. & Fester R., eds., Symbiosis as a source of evolutionary innovation, p. 319-340, MIT Press, Cambridge (Massachusetts).
- Huneck S. (1991): New results in the chemistry of lichens. - Symbiosis 11: 225-248.
- Hyyvärinen M., Halonen P. & Kauppi M. (1992): Influence of stand age and structure on the epiphytic vegetation in the middle-boreal forests of Finland. - Lichenologist 24: 165-180.

- Jacobsen P. (1992): Flechten in Schleswig-Holstein: Bestand, Gefährdung und Bedeutung als Bioindikatoren. - Mittel. Arbeitsgemeinschaft Geobot. Schleswig-Holstein u. Hamburg 42: 1-234.
- Jørgensen P.M. (1991): Difficulties in lichen nomenclature. - Mycotaxon 40: 497-501.
- Kappen L. & Breuer M. (1991): Ecological and physiological investigations in continental Antarctic cryptogams. II. Moisture relations and photosynthesis of lichens near Casey Station, Wilkes Land. - Antarctic Science 3: 273-278.
- Kiszka J. & Piorecki J. (1991): Porosty (Lichenes) Pogórza Przemyskiego. - 137 p., Uniwa, Warszawa.
- Korf R.P. (1991): Shall we abandon the principle of priority (and other nomenclatural caveats) for the sake of expediency? - Mycotaxon 40:459-468.
- Kümmerling H., Leuckert Ch. & Wirth V. (1991): Chemische Flechtenanalysen VI. *Lepraria incana* (L.) Ach. - Nova Hedwigia 53: 507-517.
- Lawrey J.D. (1991): The species-area curve as an index of disturbance in saxicolous communities. - Bryologist 94: 377-382.
- Legaz M.E. & Vicente C. (1991): An experimental approach to molecular biology of lichens: regulation of catabolite-sensitive promoters in some Usneaceae. - J. Hattori Bot. Lab. 70: 167-180.
- Leuckert Ch. & Knoph J.-G. (1992): European taxa of saxicolous *Lecidella* containing chloroxanthones: identification of patterns using thin layer chromatography. - Lichenologist 24: 383-397.
- Leuckert Ch., Knoph J.-G. & Hertel H. (1992): Chemotaxonomische Studien in der Gattung *Lecidella* (Lecanorales, Lecanoraceae) II. - Herzogia 9: 1-17.
- Leuckert Ch. & Rux K.-D. (1991): Die Flechtenflora von Berlin (West) mit besonderer Berücksichtigung epiphytischer und epigäischer Sippen (Rote Liste). - Landschaftsentwicklung und Umweltforschung 6: 119-124.
- Literski B. (1992): Verbreitung einiger Flechtenarten in Europa. - Herzogia 9: 149-166.
- Makryj T.V. (1990): Lišajniki Bajkalskogo chrebtta. - 199 p., Nauka (Sibirskoje otd.), Novosibirsk.
- Malyševa N.V. (1992): Lišajniki Staro-Petergofskogo parka Biologičeskogo Instituta Sankt-Peterburgskogo universiteta izmenenie lichenoflory za 65 let. - Vestn. Sankt-Peterburg. Univ., ser. 3, 3/17: 33-38.
- McCarthy P.M. (1992): Bibliography of Australian lichens: 1807-1991. - 62 p., National Herbarium of Victoria, Melbourne.
- Mikulín A.G. (1990): Opredelitel' lišajnikov poluostrova Kamčatka. - 137 p., Vladivostok.
- Orange A. (1992): A key to the *Cladonia chlorophaea* group in Europe, using microcrystal tests. - Brit. Lichen Soc. Bull. 70: 36-42.
- Nagarkar M.B. (1990): Lichenology in the Indian subcontinent - a review of recent work. - Acta Bryolichenol. 2: 65-78.
- Poelt J. (1990): Parasitische Arten der Flechtengattung *Rhizocarpon*: eine weitere Übersicht. - Mitteil. Bot. Staatssamml. München 29: 515-538.
- Poelt J. (1991): Homologies and analogies in the evolution of lichens. - In: Hawksworth D.L., ed., *Frontiers in mycology*, p. 85-97, CAB International, Wallingford.
- Poelt J. & Obermayer W. (1991): Beiträge zur Kenntnis der Flechtenflora des Himalaya II. Die Gattung *Bryonora* (Lichenes, Lecanoraceae) zugleich eine Revision aller Arten. - Nova Hedwigia 53: 1-26.
- Poelt J. & Petutschnig W. (1992): *Xanthoria candelaria* und ähnliche Arten in Europa. - Herzogia 9: 103-114.
- Polonyi S. & Türk R. (1991): Wachstum und Lobendifferenzierung der Blattflechte *Parmelia sulcata*. - Phytol., Horn, 31: 81-95.



- Purvis O.W. et al., eds. (1992): The lichen flora of Great Britain and Ireland. - 710 p., Brit. Lichen Soc. & Natural Hist. Mus. Publ., London.
- Rambold G., Schuhwerk F. & Triebel D. (1991): Die grosssystematischen Einheiten der Ordnung Lecanorales (Ascomycetes) und ihre ökologischen Präferenzen. - *Mitteil. Bot. Staatssamml. München* 30: 385-400.
- Rambold G. & Schwab A.J. (1990): Rusty coloured species of the lichen genus *Miriquidica* (Lecanoraceae). - *Nordic J. Bot.* 10: 117-121.
- Rambold G. & Triebel D. (1992): The inter-lecanoralean Associations. - *Biblioth. Lichenol.* 48: 1-201 p.
- Richardson D.H.S. (1991): Lichens and man. - In: Hawksworth D.L., ed., *Frontiers in mycology*, p. 187-210, CAB International, Wallingford.
- Scholz P. (1992): Rote Liste der Flechten des Landes Sachsen-Anhalt. - *Ber. Landesamt. Umweltschutz Sachsen-Anhalt* 1992/1: 38-43.
- Schöller H. (1991): Flechtenverbreitung und Klima. Vegetationsökologische Untersuchungen zur Rolle der Flechten in naturnahen Traubeneichenwäldern der Taunus. - *Biblioth. Lichenol.* 42: 1-250.
- Schreiner E. & Hafellner J. (1992): Sorediöse, corticole Krustenflechten im Ostalpenraum I. - *Biblioth. Lichenol.* 45: 1-291.
- Seppälä M., Gray J. & Ricard J. (1991): Development of low-centered ice-wedge polygons in the northernmost Ungava Peninsula, Québec, Canada. - *Boreas* 20: 259-
- Siebel H.N. et al. (1992): Rode lijst van in Nederland verwenen en bedreigde mossen en korstmossen. - *Gorteria* 18: 1-20.
- Tibell L. (1991): Revision of some taxa of Caliciales described by W. Nylander. - *Ann. Bot. Fenn.* 28: 117-121.
- Tindal E. (1991): A monograph of the genus *Toninia* (Lecideaceae, Ascomycetes). - *Opera Bot.* 110: 1-137.
- Tønnsberg T. (1992): The sorediate and isidiate, corticolous, crustose lichens in Norway. - *Sommerfeltia* 14: 1-331.
- Wedin M. (1990): Ascocarp and spore ontogeny in two species of *Sphaerophorus*. - *Nordic J. Bot.* 10: 539-545.
- Wei J.-C. (1991): An enumeration of lichens in China. - 278 p., *Int. Acad. Publ.*, Beijing.
- Wolseley P.A. & Aguirre-Hudson B. (1991): Lichens as indicators of environmental change in the tropical forests of Thailand. - *Global Ecol. & Biogeogr. Lett.* 1: 170-175.