

- Pišút I. (1995): Interessante Flechtenfunde aus Mittel-, Süd- und Südosteuropa 2. - Bibl. Lichenol. 58: 281-287.
- Pišút I. (1996): Changes in the epiphytic lichen flora of Slovakia. - In: IAL 3 - Abstracts, p. 240, Salzburg.
- Pišút I. (1996): Zaujímavejšie nálezy lišajníkov zo Slovenska 3. [Interessantere Flechtenfunde aus der Slowakei 3.] - Bull Slov. Bot. Spoloč. 18: 27-29.
- Pišút I. & Lackovičová A. (1995): Lišajníky biosférickej rezervácie Východné Karpaty. [Lichens of the Biosphere Reserve Východné Karpaty (Eastern Slovakia).] - Bryonora, Praha, 16: 25-26.
- Pišút I., Lackovičová A. & Lisická E. (1996): Lichens in Slovakia. - In: IAL 3 - Abstracts, p. 177, Salzburg.
- Tretiach M., Giralt M. & Vězda A. (1996): Gyalideopsis mexicana, a new lichen species from Chihuahua, Mexico. - Bryologist 99: 236-239.
- Vězda A. (1996): Notes on the exsiccatum "Vězda, Lichenes rariores" with an Index to fascicles 1-20 (Nos. 1-200). - 8 p., Brno.
- Vězda A. (1996): Lichenes rariores exsiccati. Fasciculus unus et vicesimus - sextus et vicesimus (numeris 201-260). - 24 p., Brno.
- Vězda A. (1996): Zemřel čestný člen ČSB a emeritní profesor dr. Josef Poelt (1924-1995). - Preslia 67(1995): 319-320.
- Wagner B. (1995): Spilonema paradoxum a Thermutis velutina - dvě překvapení v herbáři PRC. [Spilonema paradoxum and Thermutis velutina - two interesting lichens from PRC herbarium.] - Bryonora, Praha, 15: 8-9.

### NOVÁ BRYOLOGICKÁ LITERATURA V.

cestavil Z. Soldán

- Ahrens M., Schröder W. & Meinunger L. (1996): Tortula brevissima Schiffn. - über Neufunde in Deutschland. - Bryol. Mitt. 1: 31-38.
- Akiyama H. (1995): Rheophytic mosses: their morphological, physiological, and ecological adaptations. - Acta Phytotax. Geobot. 46: 77-98.
- Aleffi M. & Schumacker R. (1995): Check-list and red-list of the liverworts (Marchantiophyta) and hornworts (Anthocerotophyta) of Italy. - Fl. Medit. 5: 73-161.
- Anderson D.C., Davis R.B. & Janssens J.A. (1995): Relationships of the bryophytes and lichens to environmental gradients in Maine peatlands. - Vegetatio 120: 147-159.
- Andriessen L., Sotiaux A., Nagels C. & Sotiaux O. (1995): Aneura maxima (Schiffn.) Steph. in Belgium, new for the European liverwort flora. - J. Bryol. 18: 803-806.
- Bakken S. (1995): Effect of nitrogen supply and irradiance on growth and nitrogen status in the moss *Dicranum majus* from differently polluted areas. - J. Bryol. 18: 707-721.
- Bates J.W. (1995): Numerical analysis of bryophyte-environment relationships in a lowland English flora. - Fragn. Flor. Geobot. 40: 471-490.
- Bednarek-Ochyra H. (1995): The genus *Racomitrium* (Muscini, Grimmiaceae) in Poland: taxonomy, ecology and phytogeography. - Fragn. Flor. Geobot. 2: 3-307.
- Benkert D., Erzberger P., Klawitter J., Linder W., Linke C., Schaepe A., Steinland M. & Wiehle W. (1995): Liste der Moose von Brandenburg und Berlin mit Gefährdungsgraden. - Verhandl. Bot. Ver. Berlin Brandenburg 128: 1-68.
- Bisang I. (1995): On the phenology of *Anthoceros agrestis* (Anthocerotae, Anthocerotaceae), with special reference to central Europe. - Fragn. Flor. Geobot. 40: 513-518.
- Bisang I. (1995): The diaspore bank of hornworts (Anthocerotae, Bryophyta) and its role in the maintenance of populations in cultivated fields. - Crypt. Helvetica 18: 107-118.

- Bischler-Causse H. (1995): La flora hepaticológica del Mediterráneo. - Stud. Bot. 13: 275-278.
- Blom H.H. (1996): A revision of the *Schistidium apocarpum* complex in Norway and Sweden. - Bryophytorum Biblioth. 49: 1-333.
- Boisselier-Dubayle M.C., Jubier M.F. & Bischler H. (1995): Genetic variability in the three subspecies of *Marchantia polymorpha* (Hepaticae): isozymes, RFLP and RADP markers. - Taxon 44: 363-376.
- Carey P.D., Preston C.D., Hill M.O., Usher M.B. & Wright S.M. (1995): An environmentally defined biogeographical zonation od Scotland designed to reflect species distributions. - J. Ecol. 83: 833-846.
- Casas C. (1995): Modificaciones a New 'Checklist of Spanish Mosses' II. - Bol. Soc. Española Briol. 6: 5-6.
- Churchill S.P., Griffin D. III. & Lewis M. (1995): Moss diversity of the tropical Andes. - In: Churchill S.P. & al., eds., Biodiversity and conservation of neotropical montane forests, p. 335-346, The New York Botanical Garden, New York.
- Cronberg N. (1995): Isozyme electrophoresis of *Sphagnum*: an analysis of methodology. - Lindbergia 20: 40-48.
- Delgadillo M.C. (1995): Neotropical moss floras: species common to North and South America. - Trop. Bryol. 10: 1-6.
- De Luna E. (1995): The circumscription of phytogenetic relationships of the Hedwigiaceae (Muscii). - Syst. Botany 20: 347-373.
- Duckett J.G. y Read D.J. (1995): Ericoid mycorrhizas and rhizoid-ascomycete associations in liverworts share the same mycobiont: Isolation of the partners and resynthesis of the associations in vitro. - New Phytologist 129: 439-447.
- During H. (1995): Population regulation in tuber-bearing mosses: a simulation model. - Lindbergia 20: 26-34.
- Edwards E., Duckett J.G. & Richardson J.B. (1995): Hepatic characters in the earliest land plants. - Nature 374: 635-636.
- Fife A.J. (1995): Checklist of the mosses of New Zealand. - Bryologist 98: 313-337.
- Frego G.A. & Carleton T.J. (1995): Microsite conditions and spatial pattern in a boreal bryophyte community. - Canad. J. Bot. 73: 544-551.
- Geissler P. & Maier E. (1995): Lectotypification of central European *Grimmia* species (Muscii, Grimmiaceae). - Candollea 50: 495-514.
- Gos L. & Ochyra R. (1994): New or otherwise interesting distribution data for species of *Seligeria* (Muscii, Seligeriaceae) for Eurasia. - Fragm. Flor. Geobot. 39: 383-390.
- Gradstein S.R. (1995): Diversity of Hepaticae and Anthocerotae in montane forests of the tropical Andes. - In: Churchill S.P. & al., eds., Biodiversity and conservation of neotropical montane forests, p. 321-334, The New York Botanical Garden, New York.
- Gradstein S.R. & van Melick H.M.H. (1996): De Nederlandse Levermossen en Hauvmossen. Flora en verspreidingsatlas van de Nederlandse Hepaticae en Anthocerotae. - 366 p., Naturhistorische Bibliotheek Nr. 64, Koninklijke Nederlandse Natuurhistorische Vereniging, Utrecht.
- Greven H. & Sotiaux A. (1995): *Grimmia limprichtii*, a bryophyte with a disjunct distribution in the Alps and Himalayas. - Bryologist 98: 239-241.
- Hallingbäck T. (1995): The practices of bryophyte conservation. - Crypt. Helvetica 18: 119-130.
- Hedenäs L. (1995): Higher taxonomic level relationships among diplolepidous pleurocarpous mosses - a cladistic overview. - J. Bryol. 18: 723-782.
- Hedenäs L., Godow S. & Hylander K. (1996): 1. *Dichelyma capillaceum* (Dicks.) Myr. (Bryopsida: Fontinalaceae). - J. Bryol. 19: 157-179.
- Hederson T.A. & Longton R.E. (1995): Pattern of life history variation in the Funariales, Polytrichales and Pottiales. - J. Bryol. 18: 639-676.

- Hodgetts N.G. (1995): *Plagiochila brittanica* Paton (Hepaticae) new to Switzerland and continental Europe. - *Cryptogamie Bryol. Lichénol.* 16: 305-307.
- Huber H. (1996): Bestimmungsschlüssel für die einheimischen Arten der Gattung *Cinclidotus* nach Blatt-Merkmalen. - *Meylania* 11: 8-11.
- Karttunen K. & Toivonen H. (1995): Ecology of aquatic bryophyte assemblages in 54 small Finnish lakes and their changes in 30 years. - *Ann. Bot. Fennici* 32: 75-90.
- Kooijman A.M. & Westhoff V. (1995): Variation in habitat factors and species composition of *Scorpidium scorpioides* communities in NW-Europe. - *Vegetatio* 117: 133-150.
- Koperski M. (1996): Bemerkungen zu *Scleropodium cespitans* (Müll. Hall.) L.F.Koch. - *Bryol. Mitt.* 1: 28-30.
- Li Y. & Vitt D.H. (1995): The dynamics of moss establishment: temporal responses to a moisture gradient. - *J. Bryol.* 18: 677-688.
- Monje-Najera J. & Blanco M.A. (1995): The influence of leaf characteristics on epiphytic cover: a test of hypotheses with artificial leaves. - *Trop. Bryology* 11: 5-10.
- Mosler H.-J. (1995): Psychological foundation of intervention to protect bryophytes. - *Crypt. Helvetica* 18: 159-168.
- Muotka T. & Virtanen R. (1995): The stream as a habitat template for bryophytes: species distribution along gradients in disturbance and substratum heterogeneity. - *Freshwater Biol.* 33: 141-160.
- Mühlenhoff D. & Büdel (1995): Vergleich der Wassermoos- und Wasserflechtenvegetation zweier Bachtypen (Waldbäche/Wiesenbäche) im Spessart. - *Nova Hedwigia* 61: 525-546.
- Müller F. (1995): *Rhytidadelphus subpinnatus* - Verbreitung und Ökologie in Deutschland. - *Herzogia* 11: 101-110.
- Müller F. & al. (1995): Artenliste der Moose Sachsen. - Mater. Naturschutz Landschaftspflege 10: 1-67.
- Müller N. (1996): Pilotprojekt: Verbreitungskarten für die Moose der Schweiz. - *Meylania* 11: 23-31.
- Neinhuis C. & Jetter R. (1985): Ultrastructure and chemistry of epicuticular wax crystals in Polytrichales sporophytes. - *J. Bryol.* 18: 399-406.
- Odrzykoski I.J. (1995): Sibling species of *Conocephalum conicum* complex (Hepaticae, Conocephalaceae) in Poland: distribution maps and description of an identification method based on isozyme markers. - *Fragm. Flor. Geobot.* 40: 393-404.
- Otnyjukova T.N. (1995): Sporophyte abnormalities as a cause for decline and disappearance of mosses in polluted areas. - *Crypt. Helvetica* 18: 67-76.
- Økland R.H. (1995): Population biology of the clonal moss *Hylocomium splendens* in Norwegian boreal forests. 1. Demography. - *J. Ecol.* 83: 697-712.
- Pakarinen P. (1995): Classification of boreal mires in Finland and Scandinavia: a review. - *Vegetatio* 118: 29-38.
- Papp B. & Rajczy M. (1995): Changes of bryophyte vegetation and habitat conditions along a section of the river Danube in Hungary. - *Crypt. Helvetica* 18: 95-106.
- Potter L., Foot J.P., Caporn S.J.M. & Lee J.A. (1996): Responses of four *Sphagnum* species to acute ozone fumigation. - *J. Bryol.* 19: 19-32.
- Raeymaekers G. (1995): Policy aspect of bryophyte conservation in the European Union. - *Crypt. Helvetica* 18: 181-188.
- Schubert R. (1994): Moose - Bryophyta. - In: Rothmaler W., ed., *Excursionsflora von Deutschland*, Band 1. Niedere Pflanzen, 3. ed., p. 627-790, Gustav Fischer Verlag, Jena & Stuttgart.
- Sérgio C., Ochyra R. & Séneca A. (1995): *Dicranum crassifolium* (Musci, Dicranaceae), a new species from southern Europe. - *Fragm. Flor. Geobot.* 40: 203-214.