

- Řezáč M. (2000): Arachnofauna (Araneae, Opiliones) Klánovického lesa a jeho okolí (Praha) s důrazem na zdejší rašeliniště. – Muz. a Současnost, Roztoky, ser. natur., 14: 3-26.
- Sedláková I. et Chytrý M. (1999): Sekundární sukcese vřesovišť v Národním parku Podyjí po vypálení a pokosení: využití pro management. – Příroda, Praha, 14: 51-72.
- Sedláková I. et Chytrý M. (1999): Mohla být disturbance příčinou změny jihomoravského suchého trávníku ve vřesovišti? – Zprávy Čes. Bot. Společ., Praha, 34, Mater. 17: 25-36.
- Sofron J. (1998): Notizen zu den ausgesuchten Pflanzengesellschaften des zentralen Brdywaldes. – Folia Mus. Rer. Natur. Bohemiae-Bot., Plzeň, 41: 1-40.
- Stančík D. (1999): Změny vegetace Břehyňského rybníku v průběhu posledních 30. let. – Zprávy Čes. Bot. Společ., Praha, 34, Mater. 17: 107-122.
- Vacek S., Podhrázský V. et Souček J. (1999): Dynamika poškození smrkových a bukových porostů CHKO Orlické hory VIII. Analýza změn v PR Pod Vrchmezím a PR Sedloňovský vrch. – Příroda, Praha, 14: 145-167.
- Valachovič M. et Janovicová K. (1999): Altitudinal differentiation of oligotrophic water-spring vegetation in Slovakia. – Thaissia - J. Bot., Košice, 9: 49-62.
- Wagnerová Z. (1997): Synantropní květina u Vrbatovy boudy v Krkonoších. – Opera Corcontica, Vrchlabí, 34: 133-141.
- Zelený V. (1999): Rostliny Bílinska. – 135 p., Grada Publishing, Praha.

NOVÁ LICHENOLOGICKÁ LITERATURA IX.

cestavil Z. Palice

- Aptroot A., Brand M. et Spier L. (1998): Fellhanera viridisorediata, a new sorediate species from sheltered trees and shrubs in western Europe. – Lichenologist 30: 21-26.
- Aptroot A. et van den Boom P.P.G. (1998): Pyrenocollema chlorococcum, a new species with a chlorococcoid photobiont from zinc-contaminated soils and wood. – Cryptogamie, Bryol.-Lichénol. 19: 193-196.
- Aptroot A. et al. (1999): Checklist van de Nederlandse lichenen en lichenicole fungi. – Buxbaumiella 50: 4-64.
- Arhoun M. et al. (1998): An automatic system for stepwise treatment of solid samples and application to pollution evaluation by measuring ion lixiviation rates in lichens. – Lab Robotics and Automation 11: 121-126.
- Armaeo D. et Miao V. (1999): Symbiosis and DNA methylation in the Cladonia lichen fungus. – Symbiosis 26: 143-163.
- Arup U. et Grube M. (1999): Where does Lecanora demissa (Ascomycota, Lecanorales) belong? – Lichenologist 31: 419-430.
- Bartók K. (1998): Mapping of Peltigera species in Romania. – Sauteria 9: 13-24.
- Bartók K. et al. (1998): Studies on ¹³⁷Cs content of lichens in mountain regions of Romania. – Sauteria 9: 249-256.
- Beck A. (1999): Photobiont inventory of a lichen community growing on heavy-metal-rich rock. – Lichenologist 31: 501-510.
- Beck A., Friedl T. et Rambold G. (1998): Selectivity of photobiont choice in a defined lichen community: inferences from cultural and molecular studies. – New Phytol. 139: 709-720.
- Bennett J.P. et Wetmore C.M. (1999): Covariance of lichen and vascular plant floras. – Rhodora 101: 277-297.

- Berger F. (1999): Kompilierte Liste weiterer Flechtenfunde aus dem unteren Rannatal (Mühlviertel, Oberösterreich, Österreich) und Aspekte zur dessen Unterschutzstellung. – Beitr. Naturk. Oberösterreichs 181-203.
- Berger F., Priemetzhofer F. et Türk R. (1998): Neue und seltene Flechten und lichenicole Pilze aus Oberösterreich, Österreich IV. – Beitr. Naturk. Oberösterreichs 6: 397-416.
- Bielczyk U. (1998): Lista gatunków porostów prawnie chronionych w Polsce. - Fragn. Flor. Geobot. Ser. Polon. 5: 251-258.
- Bielczyk U. (1999): Materiały do geograficznego rozmieszczenia porostów (Lichenes) w Polsce. 1. Porosty Tatr. – Fragn. Flor. Geobot. Ser. Polon. 6: 245-253.
- Breuss O. (1998): Eine neue Verrucaria-Art mit Goniocystenthallus. – Linzer Biol. Beitr. 30: 277-279.
- Breuss O. (1998): Drei neue holz- und borkenbewohnende Verrucaria-Arten mit einem Schlüssel der bisher bekannten Taxa. - Linzer Biol. Beitr. 30: 831-836.
- Büdel B., Karsten U. et Garcia-Pichel F. (1997): Ultraviolet-absorbing scytonemin and mycosporine-like amino acid derivatives in exposed, rock-inhabiting cyanobacterial lichens. – Oecologia 112: 165-172.
- Buschbom J. et Kappen L. (1998): The role of microclimate for the lichen vegetation pattern on rock surfaces in the subarctic. – Sauteria 9: 79-94.
- Cubero O.F. et al. (1999): DNA extraction and PCR amplification method suitable for fresh, herbarium-stored, lichenized, and other fungi. – Plant Syst. Evol. 216: 243-249.
- Dettki H. (1998): Dispersal of fragments of two pendulous lichen species. – Sauteria 9: 123-131.
- Dettki H. et Esseen P.-A. (1998): Epiphytic macrolichens in managed and natural forest landscapes: a comparison at two spatial scales. – Ecography 21: 613-624.
- Edwards H.G.M., Russell N.C. et Seaward M.R.D. (1997): Calcium oxalate in lichen biodeterioration studied using FT-Raman spectroscopy. – Spectrochimica Acta, Part A 53: 99-105.
- Ekman S. (1999): PCR optimization and troubleshooting, with special reference to the amplification of ribosomal DNA in lichenized fungi. – Lichenologist 31: 517-531.
- Elix J. (1999): Detection and identification of secondary lichen substances: Contributions by the Uppsala School of Lichen Chemistry. – Symb. Bot. Ups. 32: 103-121.
- Esseen P.-A. et Renhorn K.E. (1998): Edge effects on an epiphytic lichen in fragmented forests. – Conservation Biology 12: 1307-1317.
- Figueira R. et al. (1998): In situ studies on sea-salt uptake by epiphytic lichens. – Sauteria 9: 143-150.
- Figueira R. et al. (1999): Natural levels of saline elements in lichens: determination of cellular fractions and their importance as saline tracers. – Lichenologist 31: 183-196.
- Friedl T. et Rokitta C. (1997): Species relationships in the lichen alga Trebouxia (Chlorophyta, Trebouxiophyceae): molecular phylogenetic analyses of nuclear-encoded large subunit rRNA gene sequences. – Symbiosis 23: 125-148.
- Fryday A. (1999): The genus Rimularia Nylander in the British Isles. – British Lichen Soc. Bull. 84: 18-21.
- Gastberger M. et al. (1998): Long term study of ^{137}Cs in lichens at different locations in Austria. – Sauteria 9: 273-280.
- Gauslaa Y., Ohlson M. et Rolstad J. (1998): Fine-scale distribution of the epiphytic lichen Usnea longissima on two even-aged neighbouring Picea abies trees. – Journ. Veget. Sci. 9: 95-102.
- Giralt M., van den Boom P.P.G. et Matzer M. (1997): The lichen genus Rinodina in Belgium, Luxembourg and The Netherlands. – Mycotaxon 61: 103-151.
- Goffinet B. et Goward T. (1998): Is Nephroma silvae-veteris the cyanomorph of Lobaria oregana? Insights from molecular, chemical and morphological characters. – In: Glenn M.G.

- et al. (eds.), *Lichenographia Thomsoniana: North American Lichenology in Honor of John W. Thomson*. Mycotaxon Ltd., Ithaca, New York, p. 41-52.
- Goward T., Brodo I.M. et Clayden S.R. (1998): Rare Lichens of Canada. A Review and Provisional Listing. – 74 p., Committee on the Status of Endangered Wildlife in Canada, Ottawa.
- Grube M. et al. (1999): An exceptional group-I intron-like insertion in the SSU rDNA of lichen mycobionts. – *Current Genetics* 35: 536-541.
- Gruber R. et Türk R. (1998): Die Flechtenflora und -vegetation im Gemeindegebiet von Kollerschlag (Mühlviertel, Oberösterreich) – eine Erhebung unter Naturschutzaspekten. – *Beitr. Naturk. Oberösterreichs* 6: 65-106.
- Halonen P. et al. (1999): The lichen genus *Usnea* in East Fennoscandia. III. The shrubby species. – *Ann. Bot. Fennici* 36: 235-256.
- Hauck M. et Runge M. (1999): Occurrence of pollution-sensitive epiphytic lichens in woodlands affected by forest decline: a new hypothesis. – *Flora* 194: 159-168.
- Heibel E., Lumbsch H.T. et Schmitt I. (1999): Genetic variation of *Usnea filipendula* (Parmeliaceae) populations in western Germany investigated by RAPDs suggests reinvasion from various sources. – *Amer. J. Bot.* 86: 753-757.
- Heinken T. (1999): Dispersal patterns of terricolous lichens by thallus fragments. – *Lichenologist* 31: 603-612.
- Hermansson J., Pystina T.N. et Kudrjavceva D.I. (1998): *Predvaritelnyj spisok lišajnikov Respubliki Komi*. – 133 p., Syktyvkar.
- Hestmark G. (1997): Gap-dynamics, recruitment and individual growth in populations of *Lasallia pustulata*. – *Mycological Research* 101: 1273-1280.
- Ihlen P.G. (1998): The lichenicolous fungi on species of the genera *Baeomyces*, *Dibaeis*, and *Icmadophila* in Norway. – *Lichenologist* 30: 27-57.
- Ivanova N.V. et al. (1999): Phylogenetic analysis of the lichen family Umbilicariaceae based on nuclear ITS1 and ITS2 rDNA sequences. – *Lichenologist* 31: 477-489.
- Jüriado I. (1998): A revision of the *Lecanora subfuscata* group in Estonia. – *Folia Cryptogamica Estonica* 32: 15-20.
- Kappen L. (1999): Analogien zwischen Verhaltensweisen von Symbiosen, speziell der Flechten, und pflanzlichen Organismen. In: Beyschlag W. et Steinlein T. (eds.), *Ökologie pflanzlicher Interaktionen*. Bielefelder Ökologische Beiträge, p. 121-133.
- Kärnefelt I., Emanuelsson K. et Thell A. (1998): Anatomy and systematics of usneoid genera in the Parmeliaceae. – *Nova Hedwigia* 67: 71-92.
- Lange O.L., Green T.G.A. et Reichenberger H. (1999): The response of lichen photosynthesis to external CO₂ concentration. – *Journal of Plant Physiology* 154: 157-166.
- Lumbsch H.T. (1998): Taxonomic use of metabolic data in lichen-forming fungi. – In: Frisvad J.C., Bridge P.D. et Arora D.K. (eds.), *Chemical Fungal Taxonomy*. Marcel Dekker, Inc., New York, p. 345-387.
- Martínez I. et Burgaz A.R. (1998): Revision of the genus *Solorina* (lichenes) in Europe based on spore size variation. – *Ann. Bot. Fennici* 35: 137-142.
- Mattsson J.-E. et Wedin M. (1999): A re-assessment of the family Alectoriaceae. – *Lichenologist* 31: 431-440.
- McCarthy D.P. (1999): A biological basis for lichenometry?. – *Journ. Biogeogr.* 26: 379-386.
- McCune B. et al. (1997): Repeatability of community data: species richness versus gradient scores in large-scale lichen studies. – *Bryologist* 100: 40-46.
- Miller J.E. et Brown D.H. (1999): Studies of ammonia uptake and loss by lichens. – *Lichenologist* 31: 85-93.
- Molitor F. et Diederich P. (1997): Les pyrénotriches aquatiques du Luxembourg et leurs champignons lichénicoles. – *Bull. Soc. Nat. Luxemb.* 98: 69-92.
- Murtagh G.J. et al. (1999): Use of randomly amplified polymorphic DNA markers as a tool to study variation in lichen-forming fungi. – *Lichenologist* 31: 257-267.

- Niewiadomska E., Jarowiecka D. et Czarnota P. (1998): Effect of different levels of air pollution on photosynthetic activity of some lichens. – *Acta Soc. Bot. Polon.* 67: 259-262.
- Orange A. (1998): *Dermatocarpon leptophyllum* and related species in the British Isles. – *Lichenologist* 30: 1-20.
- Palmqvist K. et al. (1998): Photosynthetic capacity in relation to nitrogen content and its partitioning in lichens with different photobionts. – *Plant, Cell and Environment* 21: 361-372.
- Priemetzhofer F. (1999): Die Flechtenflora der Stadtmauern von Freistadt (Oberösterreich) – Beitr. Naturk. Oberösterreichs 7: 127-141.
- Printzen C. et al. (1999): A study on the genetic variability of *Biatora helvola* using RAPD markers. – *Lichenologist* 31: 491-499.
- Rambold G. et Triebel D. (1999): Generic concepts in lichenized and lichenicolous ascomycetes since 1950 - a historical approach. – *Symb. Bot. Ups.* 32: 123-164.
- Randlane T. et Saag A. (1999): Second checklist of lichenized, lichenicolous and allied fungi of Estonia. – *Folia Cryptogamica Estonica* 35: 1-132.
- Rydgren K., Hestmark G. et Okland R.H. (1998): Revegetation following experimental disturbance in a boreal old-growth *Picea abies* forest. – *Journ. Veget. Sci.* 9: 763-776.
- Sarrión F., Aragón G. et Burgaz A.R. (1999): Studies on mazaediate lichens and calicioid fungi of the Iberian Peninsula. – *Mycotaxon* 71: 169-198.
- Seaward M.R.D. (1997): Major impacts made by lichens in biodeterioration processes. – *International Biodeterioration and Biodegradation* 40: 269-273.
- Seaward M.R.D. (1998): Time-space analyses of the British lichen flora, with particular reference to air quality surveys. – *Folia Cryptogamica Estonica* 32: 85-96.
- Selosse M.-A. et Le Tacon F. (1998): The land flora: a phototroph-fungus partnership? – *Trends Ecol. Evol.* 13: 15-20.
- Selva S.B. et Tibell L. (1999): Lichenized and non-lichenized calicioid fungi from North America. – *Bryologist* 102: 377-397.
- Sérusiaux E. et al. (1999): New or interesting lichens and lichenicolous fungi from Belgium and Luxembourg. VIII. – *Lejeunia*, N. S., 162: 1-95.
- Sipman H.J.M. (1999): Lichens. – In: Luteyn J.L. (ed.), *Páramos: a Checklist of Plant Diversity, Geographical Distribution, and Botanical Literature. Memoirs of The New York Botanical Garden*, New York Botanical Garden, Bronx, NY, p. 41-53.
- Søchting U. (1999): Lichens of Bhutan – Biodiversity and Use. – 30 p., Botanical Institute, Department of Mycology, University of Copenhagen, Copenhagen.
- Stenroos S. et DePriest P.T. (1998): Small insertions at a shared position in the SSU rDNA of Lecanorales (lichen-forming Ascomycetes). – *Current Genetics* 33: 124-130.
- Stenroos S. et DePriest P.T. (1998): SSU rDNA phylogeny of cladoniiform lichens. – *Amer. J. Bot.* 85: 1548-1559.
- Suija A. (1998): Lecideoid lichens from Estonia - an annotated checklist. – *Folia Cryptogamica Estonica* 32: 107-112.
- Temina M. (1998): Growth of lichens on limestone outcrops in northern Estonia. – *Sauteria* 9: 173-179.
- Trass H. (1998): List of physcioid macrolichens of Russian Far East and Siberia. – *Folia Cryptogamica Estonica* 33: 147-151.
- Türk R., Breuss O. et Üblagger J. (1998): Die Flechten im Bundesland Niederösterreich. – *Wiss. Mitt. Niederösterr. Landesmuseum* 11: 1-315.
- Türk R. et Hafellner J. (1999): Flechten. Rote Liste gefährdeter Flechten (Lichenes) Österreichs. 2. Fassung. – In: Niklfeld H. (ed.), *Rote Listen gefährdeter Pflanzen Österreichs*, Graz, p. 187-228.
- Valladares F., Sancho L.G. et Ascaso C. (1998): Water storage in the lichen family Umbilicariaceae. – *Botanica Acta* 111: 99-107.
- van den Boom P.P.G. et Giralt M. (1996): Contribution to the flora of Portugal, lichens and lichenicolous fungi I. – *Nova Hedwigia* 63: 145-172.

- van den Boom P.P.G. et Giralt M. (1999): Contribution to the flora of Portugal, lichens and lichenicolous fungi II. – *Nova Hedwigia* 68: 183-195.
- van den Boom P.P.G. et al. (1998): A lichenological excursion in May 1997 near Han-sur-Lesse and Saint-Hubert, with notes on rare or critical taxa of the flora of Belgium and Luxembourg. – *Lejeunia*, N. S., 158: 1-54.
- Wedin M. et Döring H. (1999): The phylogenetic relationship of the Sphaerophoraceae, Austropeltum and Neophyllis (lichenized Ascomycota) inferred by SSU rDNA sequences. – *Mycological Research* 103: 1131-1137.
- Wedin M., Döring H. et Mattsson J.-E. (1999): A multi-gene study of the phylogenetic relationships of the Parmeliaceae. – *Mycological Research* 103: 1185-1192.
- Winka K., Ahlberg C. et Eriksson O.E. (1998): Are there lichenized Ostropales? – *Lichenologist* 30: 455-462.
- Wirth V., Cezanne R. et Eichler M. (1999): Beitrag zur Kenntnis der Dynamik epiphytischer Flechtenbestände. – *Stuttg. Beitr. Naturk.*, Ser. A 595: 1-17.
- Zamfir M., Margaritescu S. et Moberg R (1998): Contribution to the lichen flora of Romania. 1. The lichen flora of Bucegi Mts. (Muntenia). – *Herzogia* 13: 189-197.
- Zoller S., Scheidegger C. et Sperisen C. (1999): PCR primers for the amplification of mitochondrial small subunit ribosomal DNA of lichen-forming ascomycetes. – *Lichenologist* 31: 511-516.

UZÁVĚRKA PŘÍSTÍHO ČÍSLA: 15. října 2000

Uvítáme příspěvky psané na počítači (NLQ) nebo psacím strojem s novou páskou a bez rukopisních oprav a dalších poznámek (z důvodu možnosti převodu textu na PC scannerem bez ručního přepisování). Nejvíce vitény jsou příspěvky na disketě (Word for Windows, popř. ASCII file). Při zasílání příspěvků je též možno využít elektronickou poštu s následujícími adresami: sold@natur.cuni.cz nebo liska@ibot.cas.cz. Hlavní články prosíme s anglickým abstraktem. Redaktoři si vyhrazují právo na úpravu rukopisů, popř. krácení.