

NOVÁ LICHENOLOGICKÁ LITERATURA XI.

sestavil Z. Palice

- Aptroot A., Sipman H.J.M. et van Herk C.M. (2001): Cladonia monomorpha, a neglected cup lichen from Europe. – *Lichenologist* 33: 271-283.
- Aptroot A. et al. (2001): Origin and distribution of recently described lichens from the Netherlands. – *Akt. Lichenol. Mitteil. N.F.* 5: 13-25.
- Arup U. et al. (2001): Contributions to the lichen flora of Slovenia VIII. Lichenized and lichenicolous fungi from Veliki Snežnik. – *Bibl. Lichenol.* 78: 17-25.
- BeGora M.D. et Fahselt D. (2000): An alternative method for the quantification of lichen secondary products. – *Bryologist* 103: 563-567.
- BeGora M.D. et Fahselt D. (2001): Usnic acid and atranorin concentrations in lichens in relation to bands of UV irradiance. – *Bryologist* 104: 134-140.
- Berger F. (2000): Die Flechtenflora der Schlägner Schlinge im oberösterreichischen Donautal. – *Beitr. Naturk. Oberösterr.* 9: 369-451.
- Bjelland T. (2001): Comparative studies of the distribution and ecology of some oceanic species in the genus *Leptogium* (Lecanorales, Ascomycotina) in Norway. – *Nova Hedw.* 72: 1-44.
- Boqueras M. (2000): Líquens epífits i fongs liquenícoles del sud de Catalunya: flora i comunitats. – 556 p., Institut d'Estudis Catalans, Barcelona.
- Brodо I.M., Sharnoff D.S et Sharnoff S. (2001): Lichens of North America. – 795 p., Yale University Press, New Haven.
- Casselman K.D. (2001): Lichen dyes: the new source book. – 82 p., Dover Publications, Mineola, New York.
- Coppins B.J. et Tønsberg T. (2001): A new xanthone-containing Micarea from northwest Europe and the Pacific Northwest of North America. – *Lichenologist* 33: 93-96.
- Crespo A., Blanco O. et Hawksworth D.L. (2001): The potential of mitochondrial DNA for establishing phylogeny and stabilising generic concepts in the parmelioid lichens. – *Taxon* 50: 807-819.
- Dobson F.S. (2000): Lichens. An illustrated guide to the British and Irish species. – 431p., Richmond Publishing Co. Ltd, Slough.
- Döring H. et Wedin M. (2000): Homology assessment of the boundary tissue in fruiting bodies of the lichen family Sphaerophoraceae (Lecanorales, Ascomycota). – *Plant Biology* 2: 361-367.
- Frahm J.-P., Specht A. et Reisenrath K. (2000): Allelopathic effect of crustaceous lichens on epiphytic bryophytes and vascular plants. – *Nova Hedw.* 70: 245-254.
- Friedl T. et al. (2000): The distribution of group I introns in lichen algae suggests that lichenization facilitates intron lateral transfer. – *Mol. Phylog. Evol.* 14: 342-352.
- Fryday A.M. (2001): Phytosociology of terricolous lichen vegetation in the Cairngorm Mountains, Scotland. – *Lichenologist* 33: 331-351.
- Fryday A.M. (2001): The lichen vegetation associated with areas of late snow-lie in the Scottish Highlands. – *Lichenologist* 33: 121-150.
- Galun M. et Peleg-Zuriel L. (2000): Growth enhancement of lichen-forming fungi. – *Bibl. Lichenol.* 75: 105-111.
- Gauslaa Y. et Solhaug K.A. (2001): Fungal melanins as a sun screen for symbiotic green algae in the lichen *Lobaria pulmonaria*. – *Oecologia* 126: 462-471.
- Geebelen W. et Hoffmann M. (2001): Evaluation of bio-indication methods using epiphytes by correlating with SO₂-pollution parameters. – *Lichenologist* 33: 249-260.
- Gilbert O. (2000): Lichens. – 288p., New Naturalist Library, Harper Collins Publishers, London.
- Giralt M. (2001): The Lichen genera *Rinodina* and *Rinodinella* (lichenized Ascomycetes, Physciaceae) in the Iberian Peninsula. – *Bibl. Lichenol.* 79: 1-160.
- Grube M. et Arup U. (2001): Molecular and morphological evolution in the Physciaceae (Lecanorales, lichenized Ascomycotina), with special emphasis on the genus *Rinodina*. – *Lichenologist* 33: 63-72.
- Günzl B., Fischer P. et Schmidt M. (2000): Lichenologische Untersuchungen zur Primärsukzession an Felsblöcken eines Gletschervorfeldes in den Hohen Tauern. – *Jahrb. Ver. Schutz Bergwelt* 65: 167-172.

- Hafellner J. (2000): Zur Biodiversität lichenisierter und lichenicoler Pilze in den Eisenerzer Alpen (Steiermark). – Mitt. Naturwiss. Ver. Steiermark 130: 71-106.
- Hafellner J. (2001): Bemerkenswerte Flechtenfunde in Österreich. – Fritschiana 28: 1-30.
- Hafellner J. et Türk R. (2001): Die lichenisierten Pilze Österreichs – eine Checkliste der bisher nachgewiesenen Arten mit verbreitungsangaben. – Staphia 76: 1-167.
- Harris R.C., Brodo I.M. et Tønsberg T. (2000): Lecanora thysanophora, a common leprose lichen in North America. – Bryologist 103: 790-793.
- Hauck M., Jung R. et Runge M. (2000): Does water-holding capacity of bark have an influence on lichen performance in dieback-affected spruce forests? – Lichenologist 32: 407-409.
- Hauck M. et al. (2001): Long-distance transported sulphur as a limiting factor for the abundance of *Lecanora conizaeoides* in montane spruce forests. – Lichenologist 33: 267-269.
- Hedenås H. et Ericson L. (2000): Epiphytic macrolichens as conservation indicators: successional sequence in *Populus tremula* stands. – Biol. Conserv. 93: 43-53.
- Helms G. et al. (2001): Identification of photobionts from the lichen family Physciaceae using algal-specific ITS rDNA sequencing. – Lichenologist 33: 73-86.
- Ihlen P.G., Gjerde I. et Sætersdal M. (2001): Structural indicators of richness and rarity of epiphytic lichens on *Corylus avellana* in two different forest types within a nature reserve in south-western Norway. – Lichenologist 33: 215-229.
- Jensen M. (2000): Chlorophyll fluorescence properties of the lichen *Lobaria pulmonaria* at the CO₂ compensation point. – Bibl. Lichenol. 75: 183-188.
- Jensen M., Chakir S. et Feige G.B. (1999): Osmotic and atmospheric dehydration effects in the lichens *Hypogymnia physodes*, *Lobaria pulmonaria*, and *Peltigera aphthosa*: an in vivo study of the chlorophyll fluorescence induction. – Photosynthetica 37: 393-404.
- Kantvilas G. et Jarman S.J. (1999): Lichens of Rainforest in Tasmania and South-eastern Australia. – 212 p., Australian Biological Resources, Canberra.
- Kasalicky T. et al. (2000): A comparison of ITS and LSU nrDNA phylogenies of *Fulgensia* (Teloschistaceae, Lecanorales), a genus of lichenised ascomycetes. – Can. J. Bot. 78: 1580-1589.
- Keller C. (2000): Die Wasserflechten der Teigitsch zwischen der Langmannsperrre und dem Kraftwerk Arnstein (Steiermark, Österreich). – Herzogia 14: 49-58.
- Kermitt T. et Gaulaa Y. (2001): The vertical gradient of bark pH of twigs and macrolichens in a *Picea abies* canopy not affected by acid rain. – Lichenologist 33: 353-359.
- Knop J.-G. et Leuckert C. (2000): Chemotaxonomische Studien in der Gattung *Lecidella* (Lecanorales, Lecanoraceae) III. Die gesteinbewohnenden Arten mit farblosem Hypothecium unter besonderer Berücksichtigung von europäischem Material. – Herzogia 14: 1-26.
- Kranner I. et Lutzoni F. (1999): Evolutionary consequences of transition to a lichen symbiotic state and physiological adaptation to oxidative damage associated with poikilohydry. – In: Lerner H.R., ed., Plant responses to environmental stresses: from phytohormones to genome reorganization, p. 591-628, Marcel Dekker, New York et Basel.
- Kroken S. et Taylor J.W. (2000): Phylogenetic species, reproductive mode, and specificity of the green alga *Trebouxia* forming lichens with the fungal genus *Letharia*. – Bryologist 103: 645-660.
- Kroken S. et Taylor J.W. (2001): A gene genealogical approach to recognize phylogenetic species boundaries in the lichenized fungus *Letharia*. – Mycologia 93: 38-53.
- Kumar K.C.S. et Müller K. (1999): Lichen metabolites. 2. Antiproliferative and cytotoxic activity of gyrophoric, usnic, and diffractaic acid on human keratinocyte growth. – J. Natur. Products 62: 821-823.
- Kumpula J. (2001): Winter grazing of reindeer in woodland lichen pasture: Effect of lichen availability on the condition of reindeer. – Small Ruminant Res. 39: 121-130.
- LaGreca S. et Lumbsch H.T. (2001): The phylogenetic position of the Candelariaceae (Lecanorales) inferred from anatomical and molecular data. – Bibl. Lichenol. 78: 211-222.
- Lohtander K. et al. (2000): The family Physciaceae in Fennoscandia: phylogeny inferred from ITS sequences. – Mycologia 92: 728-735.
- Lücking R. et Kalb K. (2000): Follikole Flechten aus Brasilien (vornehmlich Amazonien), inklusive einer checkliste und Bemerkungen zu *Coenogonium* und *Dimerella* (Gyalectaceae). – Bot. Jahrb. Syst. Pflanzengesch. Pflanzengeogr. 122: 1-61.
- Lumbsch H.T. et Schmitt I. (2001): Molecular data suggest that the lichen genus *Pertusaria* is not monophyletic. – Lichenologist 33: 161-170.

- Lumbsch H.T. et al. (2001): Molecular systematics supports the recognition of an additional order of Ascomycota: the Agyriales. – *Mycol. Res.* 105: 16-23.
- Lumbsch H.T. et al. (2001): ITS sequence data suggest variability of ascus types and support ontogenetic characters as phylogenetic discriminators in the Agyriales (Ascomycota). – *Mycol. Res.* 105: 265-274.
- Lutzoni F., Pagel M. et Reeb V. (2001): Major fungal lineages are derived from lichen symbiotic ancestors. – *Nature* 411: 937-940.
- Matwejuk A. (2000): Water content in terricolous lichens. – *Acta Soc. Bot. Polon.* 69: 55-63.
- McCarthy P.M. (2000): Key to the saxicolous taxa of Porina. – *Lichenologist* 32: 1-13.
- Miądlikowska J. et Lutzoni F. (2000): Phylogenetic revision of the genus *Peltigera* (lichen-forming Ascomycota) based on morphological, chemical, and large subunit nuclear ribosomal DNA data. – *Int. J. Plant Sci.* 161: 925-958.
- Moe B. et Botnen A. (2000): Epiphytic vegetation on pollarded trunks of *Fraxinus excelsior* in four different habitats at Grinde, Leikanger, western Norway. – *Plant Ecol.* 151: 143-159.
- Molina M.C. et Crespo A. (2000): Comparison of development of axenic cultures of five species of lichen-forming fungi. – *Mycol. Res.* 104: 595-602.
- Motiejūnaitė J. (2002): Lapiškosios ir krumiškosios kerpes (Ascomyctes lichenisati – Species foliosae et fruticosae). – Valstiečiu Laikraštis, Vilnius.
- Mottershed D. et Lucas G. (2000): The role of lichens in inhibiting erosion of a soluble rock. – *Lichenologist* 32: 601-609.
- Myllys L., Lohntander K. et Tehler A. (2001): β -tubulin, ITS and group I intron sequences challenge the species pair concept in *Physcia aipolia* and *P. caesia*. – *Mycologia* 93: 335-343.
- Nimis P.L. (1999): Biomonitoraggio di metalli in traccia tramite licheni in aree a rischio del Friuli-Venezia Giulia. – *Stud. Geobot.* 18: 3-49.
- Nordin A. et Mattsson J.E. (2001): Phylogenetic reconstruction of character development in Physciaceae. – *Lichenologist* 33: 3-23.
- Orange A. (2000): *Verrucaria elaeina*, a misunderstood European lichen. – *Lichenologist* 32: 411-422.
- Ott S. et Lumbsch H.T. (2001): Morphology and phylogeny of ascomycete lichens. – In: Hock B. et Esser K., ed., *The mycota. IX. fungal associations*, p. 189-210, Springer Verlag, Berlin et Heidelberg.
- Øvstedal D.O. et Lewis Smith R.I. (2001): Lichens of Antarctica and South Georgia: a guide to their identification and ecology. – 411p., *Studies in Polar Research*, Cambridge University Press, Cambridge, England.
- Owe-Larsson B. et Rambold G. (2001): The sorediate species of the lichen genus *Miriquidica* (Lecanorales, Lecanoraceae). – *Bibl. Lichenol.* 78: 335-364.
- Palmqvist K. (2000): Carbon economy of lichens. – *New Phytol.* 148: 11-36.
- Ramstad S. et Hestmark G. (2001): Population structure and size-dependent reproductive effort in *Umbilicaria sordidochroa*. – *Mycologia* 93: 453-458.
- Roitto M. et al. (2001): Does the lichen mat alleviate the effects of wet deposited nickel on soil microorganisms and Scots pine (*Pinus sylvestris* L.) seedlings? – *Plant and Soil* 230: 267-277.
- Rosso A.L., McCune B. et Rambo T.R. (2000): Ecology and conservation of a rare, old-growth-associated canopy lichen in silvicultural landscape. – *Bryologist* 103: 117-127.
- Sanders W.B. (2001): Preliminary light microscope observations of fungal and algal colonization and lichen thallus initiation on glass slides placed near foliicolous lichen communities within a lowland tropical forest. – *Symbiosis* 31: 85-94.
- Scheidegger C. et al. (2001): Evolutionary trends in the Physciaceae. – *Lichenologist* 33: 25-45.
- Schultz M., Arendholz W.-R. et Büdel B. (2001): Origin and evolution of the lichenized Ascomycete order Lichinales: monophly and systematic relationships inferred from ascus, fruiting body and SSU rDNA evolution. – *Plant Biol.* 3: 116-123.
- Sérusiaux E. et al. (2001): *Fellhanera gyrophorica*, a new European species with conspicuous pycnidia. – *Lichenologist* 33: 285-289.
- Sillett S.C. et al. (2000): Dispersal limitations of epiphytic lichens result in species dependent on old-growth forests. – *Ecol. Appl.* 10: 789-799.
- Sparrus L.A. et Aptroot A. (2000): *Fellhanera ochracea*, a new corticolous lichen species from sheltered habitats in western Europe. – *Lichenologist* 32: 515-520.
- Spier L. et Aptroot A. (2000): *Fellhaneretum myrtillicolae* ass. nov., the lichen association on *Vaccinium myrtillus*. – *Herzogia* 14: 43-47.

- Sundberg B., Näsholm T. et Palmqvist K. (2001): The effect of nitrogen on growth and key thallus components in the two tripartite lichens, *Nephroma arcticum* and *Peltigera aphthosa*. – Plant, Cell and Environm. 24: 517-527.
- Suppan U., Prügger J. et Mayrhofer H. (2000): Catalogue of the lichenized and lichenicolous fungi of Slovenia. – Bibl. Lichenol. 76: 1-215.
- Tehler A. et al. (2000): Phylogenetic analyses of the fungi based on large rDNA data sets. – Mycologia 92: 459-474.
- Thell A. et Miao V. (1999): Phylogenetic analysis of ITS and group I intron sequences from European and North American samples of cetrarioid lichens. – Ann. Bot. Fenn. 35: 275-286.
- Thell A., Stenroos S. et Myllys L. (2000): A DNA study of the *Cetraria aculeata* and *C. islandica* groups. – Folia Cryptog. Eston. 36: 95-106.
- Tibell L. (2001): Photobiont association and molecular phylogeny of the lichen genus *Chaenotheca*. – Bryologist 104: 191-198.
- Tønsberg T., Türk R. et Hofmann P. (2001): Notes on the lichen flora of Tyrol (Austria). – Nova Hedw. 72: 487-497.
- Uliczka H. et Angelstam P. (1999): Occurrence of epiphytic macrolichens in relation to tree species and age in managed boreal forest. – Ecography 22: 396-405.
- van den Boom P.P.G. et Coppins B.J. (2001): *Micarea viridileprosa* sp. nov., an overlooked lichen species from Western Europe. – Lichenologist 33: 87-91.
- van Herk K. et Aptroot A. (2000): The sorediate *Punctelia* species with lecanoric acid in Europe. – Lichenologist 32: 233-246.
- Weber B. et Büdel B. (2001): Mapping and analysis of distribution patterns of lichens on rural medieval churches in north-eastern Germany. – Lichenologist 33: 231-248.
- Wedin M., Döring H. et Ekman S. (2000): Molecular phylogeny of the lichen families Cladoniaceae, Sphaerophoraceae, and Stereocaulaceae (Lecanorales, Ascomycotina). – Lichenologist 32: 171-187.
- Wetmore C.M. (2001): The *Caloplaca citrina* group in North and Central America. – Bryologist 104: 1-11.
- Zavada M.S. et Simoes P. (2001): The possible demi-lichenization of the basidiocarps of *Trametes versicolor* (L.:Fries) Pilát (Polyporaceae). – Northeast. Natur. 8: 101-112.
- Zedda L. (2000): *Lecanora leuckertiana* sp. nov. (lichenized Ascomycetes, Lecanorales) from Italy, Greece, Morocco and Spain. – Nova Hedw. 71: 107-112.