

## NOVÁ BRYOLOGICKÁ LITERATURA XIV.

### New bryological literature, XIV

Jan Kučera

Jihočeská Univerzita, biologická fakulta, Branišovská 31, CZ-370 05 České Budějovice, e-mail: kucera@bf.jcu.cz

- Adamczak M., Buczkowska K., Baczkiewicz A. & Wachowiak W. (2005): Comparison of allozyme variability in Polish populations of two species of *Ptilidium* Nees (*Hepaticae*) with contrasting degrees of sexual reproduction. – Cryptogamie: Bryologie 26: 151–165.
- Afonina O. M. (2004): Vidy *Hypnum* sekcií *Hamulosa* (*Musci*, *Hypnaceae*) v Rossii [The *Hypnum* sect. *Hamulosa* (*Musci*, *Hypnaceae*) in Russia]. – Arctoa 13: 9–28.
- Agnello G., Manneville O. & Asta J. (2004): Mosses and lichens as bioindicators (s. l.) of wetlands state: example of four protected areas in the Isère Department (France). – Revue D'Ecologie – La Terre et la Vie 59: 147.
- Aguirre J. & Rangel J. O. (2005): Species diversity and richness of the mosses of the Colombian Choco region. – Journal of the Hattori Botanical Laboratory 97: 97–116.
- Ahrens M. (2003): Untersuchungen zum Lebenszyklus von *Acaulon triquetrum* (*Bryopsida*, *Pottiaceae*). – Herzogia 16: 239–274.
- Aicardi O. (2004): Contribution à l'inventaire de la bryoflore française (année 2003). – Bulletin de la Société Botanique du Centre-Ouest, Nouvelle Série 35: 337–342.
- Akhani H. & Kürschner H. (2004): An annotated and updated checklist of the Iranian bryoflora. – Cryptogamie: Bryologie 25: 315–347.
- Albertos B., Lara F., Garilletti R. & Mazimpaka V. (2005): A survey of the epiphytic bryophyte flora in the northwest of the Iberian Peninsula. – Cryptogamie: Bryologie 26: 263–289.
- Aleffi M., Cortini Pedrotti C. & Gafta D. (2003): Bryogeographic patterns in the small islands surrounding the Italian peninsula, Sicily and Sardinia. – Bocconea 16: 93–103.
- Aleffi M., Sabovljević M. & Tacchi R. (2004): *Gymnostomum lanceolatum* M. J. Cano, Ros & J. Guerra (*Pottiaceae*, *Musci*), new to Italy. – Cryptogamie: Bryologie 25: 175–177.
- Al-Masri M. S., Mamish S., Al-Haleem M. A. & Al-Shamali K. (2005): *Lycopodium cernuum* and *Funaria hygrometrica* as deposition indicators for radionuclides and trace metals. – Journal of Radioanalytical and Nuclear Chemistry 266: 49–55.
- Amblard-Gross G., Maul A., Férid J.-F., Carrot F. & Ayraut S. (2004): Spatial variability of sampling: Grid size impact on atmospheric metals and trace elements deposition mapping with mosses. – Journal of Atmospheric Chemistry 49: 39–52.
- Amon J. P., Jacobson C. S. & Shelley M. L. (2005): Construction of fens with and without hydric soils. – Ecological Engineering 24: 341–357.
- Anand M., Laurence S. & Rayfield B. (2005): Diversity relationships among taxonomic groups in recovering and restored forests. – Conservation Biology 19: 955–962.
- Andersson M. S. & Gradstein S. R. (2005): Impact of management intensity on non-vascular epiphyte diversity in cacao plantations in western Ecuador. – Biodiversity and Conservation 14: 1101–1120.
- Araújo M. B. (2004): Matching species with reserves – uncertainties from using data at different resolutions. – Biological Conservation 118: 533–538.
- Arazi T., Talmor-Neiman M., Stav R., Riese M., Huijser P. & Baulcombe D. C. (2005): Cloning and characterization of micro-RNAs from moss. – Plant Journal 43: 837–848.
- Ashkenas L. R., Johnson S. L., Gregory S. V., Tank J. L. & Wollheim W. M. (2004): A stable isotope tracer study of nitrogen uptake and transformation in an old-growth forest stream. – Ecology 85: 1725–1739.
- Aude E. & Ejrnæs R. (2005): Bryophyte colonisation in experimental microcosms: the role of nutrients, defoliation and vascular vegetation. – Oikos 109: 323–330.
- Augusto L., Dupouey J.-L. & Ranger J. (2003): Effects of tree species on understory vegetation and environmental conditions in temperate forests. – Annals of Forest Science 60: 823–831.
- Austreim G., Hassel K. & Mysterud A. (2005): The role of life history traits for bryophyte community patterns in two contrasting alpine regions. – Bryologist 108: 259–271.

- Axtell M. J. & Bartel D. P. (2005): Antiquity of microRNAs and their targets in land plants. – *Plant Cell* 17: 1658–1673.
- Bakalin V. A. (2003): A preliminary check-list of the hepatics of Kamchatka Peninsula (Russian Far East). – *Arctoa* 12: 83–90.
- Bakalin V. A. (2004): *Macrodiplophyllum imbricatum* (*Scapaniaceae, Hepaticae*) – novyj vid dlya flory Rosii [*Macrodiplophyllum imbricatum* (*Scapaniaceae, Hepaticae*), a new species to the flora of Russia]. – *Botanicheskiy Zhurnal* 89: 1016–1020.
- Bakalin V. A. (2004): Notes on *Lophozia* V. Comments on Sect. *Sudeticae*, *Longidentatae* and *Savicziae*. – *Arctoa* 13: 229–240.
- Bakalin V. A. (2005): *Schofieldia* i *Cryptocoleopsis* – novye rody pechenochnikov (*Hepaticae*) dlya flory Rossii. – *Botanicheskiy Zhurnal* 90: 594–603.
- Bakalin V. A. & Taran G. S. (2004): Rod *Riccia* (*Hepaticae*) v Sibiri i Vostochnom Kazakhstane [The genus *Riccia* (*Hepaticae*) in Siberia and the East Kazakhstan]. – *Botanicheskiy Zhurnal* 89: 1283–1293.
- Baldwin L. K. & Bradfield G. E. (2005): Bryophyte community differences between edge and interior environments in temperate rain-forest fragments of coastal British Columbia. – *Canadian Journal of Forest Research – Revue Canadienne de Recherche Forestière* 35: 580–592.
- Bardunov L. V. & Vasil'ev A. N. (2005): Puti formirovania ekologicheskikh grupp mkhov vo flore tajgi. – *Botanicheskiy Zhurnal* 90: 90: 527–535.
- Bargagli R., Agnorelli C., Borghini F. & Monaci F. (2005): Enhanced deposition and bioaccumulation of mercury in Antarctic terrestrial ecosystems facing a coastal polynya. – *Environmental Science & Technology* 39: 8150–8155.
- Bargagli R., Skotnicki M. L., Marri L., Pepi M., Mackenzie A. & Agnorelli C. (2004): New record of moss and thermophilic bacteria species and physico-chemical properties of geothermal soils on the northwest slope of Mt. Melbourne (Antarctica). – *Polar Biology* 27: 423–431.
- Barker D. H., Stark L. R., Zimpfer J. F., McLetchie N. D. & Smith S. D. (2005): Evidence of drought-induced stress on biotic crust moss in the Mojave Desert. – *Plant Cell and Environment* 28: 939–947.
- Basiliko N., Knowles R. & Moore T. R. (2004): Roles of moss species and habitat in methane consumption potential in a northern peatland. – *Wetlands* 24: 178–185.
- Bates J. W., Thompson K. & Grime J. P. (2005): Effects of simulated long-term climatic change on the bryophytes of a limestone grassland community. – *Global Change Biology* 11: 757–769.
- Beckett R. P., Marschall M. & Laufer Z. (2005): Hardening enhances photoprotection in the moss *Atrichum androgynum* during rehydration by increasing fast- rather than slow-relaxing quenching. – *Journal of Bryology* 27: 7–12.
- Beckett R. P., Minibayeva F. V., Lüthje S. & Böttger M. (2004): Reactive oxygen species metabolism in desiccation-stressed thalli of the liverwort *Dumontiera hirsuta*. – *Physiologia Plantarum* 122: 3–10.
- Bednarek-Ochyra H. (2004): *Codriophorus corrugatus* (*Bryopsida, Grimmiaceae*), a new species from east Asia and southern Alaska. – *Bryologist* 107: 377–384.
- Bednarek-Ochyra H. (2004): Does *Racomitrium aciculare* occur in China and South Africa? – *Bryologist* 107: 197–201.
- Bednarek-Ochyra H. (2004): On the occurrence of *Racomitrium aquaticum* (Schrad.) Brid. (*Grimmiaceae, Musci*) in Asiatic Russia. – *Cryptogamie: Bryologie* 25: 83–89.
- Bednarek-Ochyra H. & Ochyra R. (2004): Lectotypification of *Codriophorus aquaticus* (*Bryopsida, Grimmiaceae*). – *Bryologist* 107: 480–488.
- Behre K.-E., Hölzer A. & Lemdahl G. (2005): Botanical macro-remains and insects from the Eemian and Weichselian site of Oerel (northwest Germany) and their evidence for the history of climate. – *Vegetation History and Archaeobotany* 14: 31–53.
- Belkina O. A. & Likhachev A. Yu. (2004): Flora listostebel'nykh mkhov gornykh massivov Chil'tal'd i Ionn-Nyugoayv (Murmanskaya oblast') [Moss flora of Chiltald and Ionn-Nyugoayv mountains (Murmansk Province, Russia)]. – *Arctoa* 13: 211–222.
- Belland R. J. (2005): A multivariate study of moss distributions in relation to environment in the Gulf of St. Lawrence region, Canada. – *Canadian Journal of Botany – Revue Canadienne de Botanique* 83: 243–263.
- Benscoter B. W., Kelman-Wieder R. & Vitt D. H. (2005): Linking microtopography with post-fire succession in bogs. – *Journal of Vegetation Science* 16: 453–460.
- Berg C. & Dengler J. (2005): Moose und Flechten als diagnostische Arten von Pflanzengesellschaften – eine Übersicht aus Mecklenburg-Vorpommern. – *Herzogia* 18: 145–161.

- Bisang I. & Hedenäs L. (2005): Sex ratio patterns in dioicous bryophytes re-visited. – *Journal of Bryology* 27: 207–220.
- Bleuel C., Wesenberg D., Sutter K., Miersch J., Braha B., Barlocher F. & Krauss G.-J. (2005): The use of the aquatic moss *Fontinalis antipyretica* L. ex Hedw. as a bioindicator for heavy metals – 3. Cd<sup>2+</sup> accumulation capacities and biochemical stress response of two *Fontinalis* species. – *Science of the Total Environment* 345: 13–21.
- Blockeel T. L., Vanderpoorten A., Sotiaux A. & Goffinet B. (2005): The status of the mid-western European endemic moss, *Brachythecium appleyardiae*. – *Journal of Bryology* 27: 137–141.
- Bond-Lamberty B., Wang C. K. & Gower S. T. (2004): Net primary production and net ecosystem production of a boreal black spruce wildfire chronosequence. – *Global Change Biology* 10: 473–487.
- Bonini I., Casini F., Chiarucci A. & De Dominicis V. (2005): The bryophyte flora of the geothermal field of Sasso Pisano (Pisa, Italy). – *Cryptogamie: Bryologie* 26: 291–299.
- Borghini F., Grimalt J. O., Sánchez-Hernández J. C. & Bargagli R. (2005): Organochlorine pollutants in soils and mosses from Victoria Land (Antarctica). – *Chemosphere* 58: 271–278.
- Bowker M. A., Belnap J., Rosentreter R. & Graham B. (2004): Wildfire-resistant biological soil crusts and fire-induced loss of soil stability in Palouse prairies, USA. – *Applied Soil Ecology* 26: 41–52.
- Brackel W. & Howein H. (2004): *Dicranum viride* in Ober- und Mittelfranken – Standortansprüche und Vergesellschaftung. – *Berichte der Bayerischen Botanischen Gesellschaft zur Erforschung der Heimischen Flora* 73/74: 129–134.
- Bragazza L., Limpens J., Gerdol R., Grosvernier P., Hájek M., Hájek T., Hájková P., Hansen I., Iacumin P., Kutnar L., Rydin H. & Tahvanainen T. (2005): Nitrogen concentration and δ<sup>15</sup>N signature of ombrotrophic *Sphagnum* mosses at different N deposition levels in Europe. – *Global Change Biology* 11: 106–114.
- Brown R. C. & Lemmon B. E. (2004): γ-Tubulin, microtubule arrays, and quadripolarity during sporogenesis in the hepatic *Aneura pinguis* (*Metzgeriales*). – *Journal of Plant Research* 117: 371–376.
- Brullo S., Privitera M. & Puglisi M. (2004): Bryophyte vegetation of the fumaroles from some Mediterranean and Macaronesian territories. – *Nova Hedwigia* 78: 367–387.
- Buck W. R., Cox C. J., Shaw A. J. & Goffinet B. (2005): Ordinal relationships of pleurocarpous mosses, with special emphasis on the *Hookeriales*. – *Systematics and Biodiversity* 2: 121–145.
- Budyakova A. A., Ignatov M. S., Yatsentyuk S. P. & Troitsky A. V. (2003): Systematic position of *Habrodon* (*Habrodontaceae, Musci*) as inferred from nuclear ITS1 and ITS2 and chloroplast *trnL* intron and *trnL-trnF* spacer sequence data. – *Arctoa* 12: 137–150.
- Buryová B. & Shaw A. J. (2005): Phenotypic plasticity in *Philonotis fontana* (*Bryopsida: Bartramiaceae*). – *Journal of Bryology* 27: 13–22.
- Callaghan T. V., Björn L. O., Chernov Y., Chapin T., Christensen T. R., Huntley B., Ims R. A., Johansson M., Jolly D., Jonasson S., Matveyeva N., Panikov N., Oechel W., Shaver G., Elster J., Jónsdóttir I. S., Laine K., Taulavuori K., Taulavuori E. & Zöckler C. (2004): Responses to projected changes in climate and UV-B at the species level. – *Ambio* 33: 418–435.
- Campisi P. & Provenzano F. (2004): New record in Sicily of *Gigaspermum mouretii* (*Gigaspermaceae, Musci*), rare species in Europe. – *Flora Mediterranea* 14: 305–308.
- Campisi P., Dia M. G. & Aiello P. (2003): First record of *Anacolia webbii* (*Bartramiaceae, Musci*) in Italian peninsula. – *Flora Mediterranea* 13: 327–330.
- Cannone N. (2004): Minimum area assessment and different sampling approaches for the study of vegetation communities in Antarctica. – *Antarctic Science* 16: 157–164.
- Cannone N., Guglielmin M. & Gerdol R. (2004): Relationships between vegetation patterns and periglacial landforms in northwestern Svalbard. – *Polar Biology* 27: 562–571.
- Cano M. J. (2005): Proposal to conserve the name *Pseudocrossidium revolutum* (*Pottiaceae, Bryophyta*) with a conserved type. – *Taxon* 54: 189–190.
- Cano M. J., Jiménez J. A., Gallego M. T., Ros R. M. & Guerra J. (2004): Bryophyte check-list of Murcia Province (southeastern Spain). – *Annales de Biología* 2, Sección Especial 2, Universidad de Murcia 26: 117–155.
- Cano M. J., Werner O. & Guerra J. (2005): A morphometric and molecular study in *Tortula subulata* complex (*Pottiaceae, Bryophyta*). – *Botanical Journal of the Linnean Society* 147: 333–350.
- Cao T. & Koponen T. (2004): Musci in ‘Symbolae sinicae’; an annotated checklist of mosses collected by H. Handel-Mazzetti in China in 1914–1918, and described by V. F. Brotherus in 1922–1929. – *Bryobrothera* 8: 1–34.

- Carafa A., Duckett J. G., Knox J. P. & Ligrone R. (2005): Distribution of cell-wall xylans in bryophytes and tracheophytes: new insights into basal interrelationships of land plants. – *New Phytologist* 168: 231–240.
- Cargill D. C., Renzaglia K. S., Villarreal J. C. & Duff R. J. (2005): Generic concepts within hornworts: historical review, contemporary insights and future directions. – *Australian Systematic Botany* 18: 7–16.
- Casas C., Brugués M. & Cros R. M. (2004): Flora dels briòfits dels Països Catalans, Vol. II. Hepàtiques i Antocerotes. – Institut d’Estudis Catalans, Barcelona. [138 pp.]
- Casas-Garcia S., Burgaz A. R. & Escudero A. (2005): Effects of the holm-oak canopy on the distribution of terricolous bryophytes and lichens. – *Cryptogamie: Bryologie* 26: 67–77.
- Castells E., Peñuelas J. & Valentine D. W. (2005): Effects of plant leachates from four boreal understorey species on soil N mineralization, and white spruce (*Picea glauca*) germination and seedling growth. – *Annals of Botany* 95: 1247–1252.
- Cattaneo A., Cloutier L. & Méthot G. (2004): The response of invertebrates in moss and in gravel to water level fluctuations in a Québec stream. – *Archiv für Hydrobiologie* 161: 21–43.
- Chapin C. T., Bridgman S. D. & Pastor J. (2004): pH and nutrient effects on above-ground net primary production in a Minnesota, USA bog and fen. – *Wetlands* 24: 186–201.
- Chen H. Y., Légaré S. & Bergeron Y. (2004): Variation of the understory composition and diversity along a gradient of productivity in *Populus tremuloides* stands of northern British Columbia, Canada. – *Canadian Journal of Botany-Revue Canadienne de Botanique* 82: 1314–1323.
- Christensen M., Hahn K., Mountford E. P., Ódor P., Standovár T., Rozenbergar D., Daci J., Wijdeven S., Meyer P., Winter S. & Vrška T. (2005): Dead wood in European beech (*Fagus sylvatica*) forest reserves. – *Forest Ecology and Management* 210: 267–282.
- Chuyong G. B., Newbery D. M. & Songwe N. C. (2004): Rainfall input, throughfall and stemflow of nutrients in a central African rain forest dominated by ectomycorrhizal trees. – *Biogeochemistry* 67: 73–91.
- Clark K. L., Nadkarni N. M. & Gholz H. L. (2005): Retention of inorganic nitrogen by epiphytic bryophytes in a tropical montane forest. – *Biotropica* 37: 328–336.
- Cleavitt N. (2004): Comparative ecology of a lowland and a subalpine species of *Mnium* in the northern Rocky Mountains. – *Plant Ecology* 174: 205–216.
- Cleavitt N. (2004): Controls on the distribution of *Mnium arizonicum* along an elevation gradient in the Front Ranges of the Rocky Mountains, Alberta. – *Journal of the Torrey Botanical Society* 131: 150–160.
- Cleavitt N. (2005): How many bryophytes are there in New York and are any of them rare? – *Solidago* 6: 1–6.
- Cortini Pedrotti C. (2005): Flora dei muschi d’Italia. *Bryopsida* (II parte). – Antonio Delfino Editore, Roma. [432 pp.]
- Couto J. A., Aboal J. R., Fernández J. A. & Carballeira A. (2004): A new method for testing the sensitivity of active biomonitoring: an example of its application to a terrestrial moss. – *Chemosphere* 57: 303–308.
- Couto J. A., Fernández J. A., Aboal J. R. & Carballeira A. (2004): Active biomonitoring of element uptake with terrestrial mosses: a comparison of bulk and dry deposition. – *Science of the Total Environment* 324: 211–222.
- Couwenberg J. & Joosten H. (2005): Self-organization in raised bog patterning: the origin of microtopose zonation and mesotope diversity. – *Journal of Ecology* 93: 1238–1248.
- Crandall-Stotler B. J., Forrest L. L. & Stotler R. E. (2005): Evolutionary trends in the simple thalloid liverworts (*Marchantiophyta*, *Jungermanniopsida* subclass *Metzgeriidae*). – *Taxon* 54: 299–316.
- Croisetière L., Hare L., Tessier A. & Duchesne S. (2005): Modeling cadmium exchange by an aquatic moss (*Fontinalis dalecarlica*). – *Environmental Science & Technology* 39: 3056–3060.
- Cros R. M., Sáez L. & Rosselló J. A. (2005): *Plagiochasma appendiculatum* Lehm. & Lindenb. (*Marchantiales*, *Aytoniaceae*), a species new to the European bryophyte flora. – *Journal of Bryology* 27: 3–6. [Erratum p. 190]
- Crow S. E. & Wieder R. K. (2005): Sources of CO<sub>2</sub> emission from a northern peatland: Root respiration, exudation, and decomposition. – *Ecology* 86: 1825–1834.
- Crowley P. H., Stieha C. R. & McLetchie D. N. (2005): Overgrowth competition, fragmentation and sex-ratio dynamics: a spatially explicit, sub-individual-based model. – *Journal of Theoretical Biology* 233: 25–42.
- Csotonyi J. T. & Addicott J. F. (2004): Influence of trampling-induced microtopography on growth of the soil crust bryophyte *Ceratodon purpureus* in Jasper National Park. – *Canadian Journal of Botany – Revue Canadienne de Botanique* 82: 1382–1392.
- Cucu-Man S., Mocanu R., Culicov O., Steinnes E. & Frontasyeva M. (2004): Atmospheric deposition of metals in Romania studied by biomonitoring using the epiphytic moss *Hypnum cupressiforme*. – *International Journal of Environmental Analytical Chemistry* 84: 845–854.

- Cuny D., Denayer F.-O., de Foucault B., Schumacker R., Colein P. & Van Haluwyn C. (2004): Patterns of metal soil contamination and changes in terrestrial cryptogamic communities. – Environmental Pollution 129: 289–297.
- Curtis C. J., Emmett B. A., Grant H., Kernan M., Reynolds B. & Shilland E. (2005): Nitrogen saturation in UK moorlands: the critical role of bryophytes and lichens in determining retention of atmospheric N deposition. – Journal of Applied Ecology 42: 507–517.
- Czernyadjeva I. V. (2003): Rod *Hygrohypnum* (*Amblystegiaceae, Musci*) v Rossii [The genus *Hygrohypnum* (*Amblystegiaceae, Musci*) in Russia]. – Arctoa 12: 25–58.
- Czernyadjeva I. V. & Ignatova E. A. (2004): *Pohlia tundrae* Shaw (*Musci*) in Russia. – Arctoa 13: 29–32.
- Da Costa D. P., Imbassahy C. A. D. & Da Silva V. P. A. V. (2005): Checklist and distribution of mosses, liverworts and hornworts of Rio de Janeiro State, Brazil. – Journal of the Hattori Botanical Laboratory 98: 259–298.
- Daniëls F. J. A., Talbot S. S., Talbot S. L. & Schofield W. B. (2004): Phytosociological study of the dwarf shrub heath of Simeonof Wilderness, Shumagin Islands, Southwestern Alaska. – Phytocoenologia 34: 465–489.
- De Zuttere Ph. (2005): La mousse *Tortella bambengeri* (Schimp.) Broth. (*Pottiaceae*) doit être considérée comme une bonne espèce: sa répartition en Europe. – Nowellia Bryologica 29: 12–23.
- Demkiv O. T., Khorkavtsiv Ya. D., Kyiak N. Ya. & Kit N. A. (2005): Vplyv gravitacyji na fotomorfogenez protonemy *Pottia intermedia* (Turn.) Furnr., *Pottiales* [Effect of gravity on photomorphogenesis of protonemata *Pottia intermedia* (Turn.) Furnr., *Pottiales*]. – Ukrains’kyi Botanichnyi Zhurnal 62: 329–336.
- Desponts M., Brunet G., Bélanger L. & Bouchard M. (2004): The eastern boreal old-growth balsam fir forest: a distinct ecosystem. – Canadian Journal of Botany – Revue Canadienne de Botanique 82: 830–849.
- Dia M. G., Privitera M. & Puglisi M. (2003): New reports of *Pseudocrossidium replicatum* (*Pottiaceae, Musci*) from Sicily. – Flora Mediterranea 13: 297–301.
- Dickson J. H., Hofbauer W., Kofler W., Oegg K. & Platzgummer J. (2005): How to find the bogmoss, *Sphagnum imbricatum* s.l., in South Tyrol, Italy: Microscopically examine the Iceman’s colon contents. – Vegetation History and Archaeobotany 14: 207–210.
- Djingova R., Kuleff I. & Markert B. (2004): Chemical fingerprinting of plants. – Ecological Research 19: 3–11.
- Dorrepaal E., Cornelissen J. H. C., Aerts R., Wallén B. & van Logtestijn R. S. P. (2005): Are growth forms consistent predictors of leaf litter quality and decomposability across peatlands along a latitudinal gradient? – Journal of Ecology 93: 817–828.
- Dowdall M., Gwynn J. P., Moran C., O’Dea J., Davids C. & Lind B. (2005): Uptake of radionuclides by vegetation at a High Arctic location. – Environmental Pollution 133: 327–332.
- Draper I., Mazimpaka V., Alberto B., Garilleti R. & Lara F. (2005): A survey of the epiphytic bryophyte flora of the Rif and Tazzeka Mountains (northern Morocco). – Journal of Bryology 27: 23–34.
- Drehwald U. (2005): Biomonitoring of disturbance in neotropical rainforests using bryophytes as indicators. – Journal of the Hattori Botanical Laboratory 97: 117–126.
- Duff R. J. & Moore F. B. G. (2005): Pervasive RNA editing among hornwort *rbcL* transcripts except *Leiosporoceros*. – Journal of Molecular Evolution 61: 571–578.
- Eckstein J. (2004): *Acaulon casasianum* (*Musci, Pottiaceae*) – neu für die Flora von Mitteleuropa. – Haussknechtia 10: 103–112.
- Eckstein R. L., Danihelka J., Hözel N. & Otte A. (2004): The effects of management and environmental variation on population stage structure in three river-corridor violets. – Acta Oecologica – International Journal of Ecology 25: 83–91.
- Ederra A. & Guerra J. (2005): *Cinclidotus vivesii* sp. nov. (*Musci, Pottiaceae*) from the Iberian Peninsula. – Nova Hedwigia 81: 471–476.
- Ederra A. (2005): *Pottiaceae: Cinclidotus*. – In: Guerra J. & Cros R. M. (eds.), Flora Briofítica Ibérica, p. 16–22, Sociedad Española de Briología, Murcia.
- Ederra A. (2005): *Pottiaceae: Scopelophila*. – In: Guerra J. & Cros R. M. (eds.), Flora Briofítica Ibérica, p. 5–8, Sociedad Española de Briología, Murcia.
- Edwards D. & Richardson J. B. (2004): Silurian and Lower Devonian plant assemblages from the Anglo-Welsh Basin: a palaeobotanical and palynological synthesis. – Geological Journal 39: 375–402.
- Ellis C. J. & Rochefort L. (2004): Century-scale development of polygon-patterned tundra wetland, Bylot island (73° N, 80° W). – Ecology 85: 963–978.

- El-Saadawi W., Shabbara H. M., Refai S. M. & Abou-Salama U. Y. (2003): Mosses of different phytogeographical territories of Egypt. – *Bocconeia* 16: 133–146.
- Engel J. J. & Braggins J. E. (2005): Are *Mylia* and *Trabacellula* (*Hepaticae*) related? Unsuspected links revealed by cell wall morphology, with the transfer of *Mylia anomala* to a new genus (*Leiomylia* J.J. Engel & Braggins) of *Jungmanniaceae*. – *Taxon* 54: 665–680.
- Ericsson T. S., Berglund H. & Östlund L. (2005): History and forest biodiversity of woodland key habitats in south boreal Sweden. – *Biological Conservation* 122: 289–303.
- Ermakova E. V., Frontasyeva M. V., Pavlov S. S., Povtoreiko E. A., Steinnes E. & Cheremisina Y. N. (2004): Air pollution studies in Central Russia (Tver and Yaroslavl regions) using the moss biomonitoring technique and neutron activation analysis. – *Journal of Atmospheric Chemistry* 49: 549–561.
- Estébanez B., Ron E., Alfayate C., González A. & Bermejo J. (2004): The occurrence of lipidic storage compounds in *Grimmia* (*Bryopsida*). – *Bryologist* 107: 501–504.
- Eversman S. & Horton D. (2004): Recolonization of burned substrates by lichens and mosses in Yellowstone National Park. – *Northwest Science* 78: 85–92.
- Feldberg K., Groth H., Wilson R., Schäfer-Verwimp A. & Heinrichs J. (2004): Cryptic speciation in *Herbertus* (*Herbertaceae*, *Jungmanniopsida*): range and morphology of *Herbertus sendtneri* inferred from nrITS sequences. – *Plant Systematics and Evolution* 249: 247–261.
- Feldberg K. & Heinrichs J. (2005): On the identity of *Herbertus borealis* (*Jungmanniopsida*: *Herbertaceae*), with notes on the possible origin of *H. sendtneri*. – *Journal of Bryology* 27: 343–350.
- Fenton N. J. & Frego K. A. (2005): Bryophyte (moss and liverwort) conservation under remnant canopy in managed forests. – *Biological Conservation* 122: 417–430.
- Fenton N., Lecomte N., Légaré S. & Bergeron Y. (2005): Paludification in black spruce (*Picea mariana*) forests of eastern Canada: Potential factors and management implications. – *Forest Ecology and Management* 213: 151–159.
- Figueira R. & Ribeiro T. (2005): Transplants of aquatic mosses as biomonitor of metals released by a mine effluent. – *Environmental Pollution* 136: 293–301.
- Fleischbein K., Wilcke W., Goller R., Boy J., Valarezo C., Zech W. & Knoblich K. (2005): Rainfall interception in a lower montane forest in Ecuador: effects of canopy properties. – *Hydrological Processes* 19: 1355–1371.
- Fletcher B. J., Beerling D. J., Brentnall S. J. & Royer D. L. (2005): Fossil bryophytes as recorders of ancient CO<sub>2</sub> levels: Experimental evidence and a Cretaceous case study. – *Global Biogeochemical Cycles* 19/3: GB3012.
- Flint S. D., Ryel R. J. & Caldwell M. M. (2003): Ecosystem UV-B experiments in terrestrial communities: a review of recent findings and methodologies. – *Agricultural and Forest Meteorology* 120: 177–189.
- Forrest L. L. & Crandall-Stotler B. J. (2005): Progress towards a robust phylogeny for the liverworts, with particular focus on the simple thalloids. – *Journal of the Hattori Botanical Laboratory* 97: 127–159.
- Frahm J.-P. (2004): A guide to bryological hotspots in Europe. 1. The Azores. – *Archive for Bryology* 3: 1–14.
- Frahm J.-P. (2004): A new contribution to the moss flora of Baltic and Saxon amber. – *Review of Palaeobotany and Palynology* 129: 81–101.
- Frahm J.-P. (2005): Neue Laubmoosfunde in Baltischem Bernstein [New records of mosses from Baltic amber]. – *Palaeontographica Abteilung B, Paläophytologie* 271: 159–168.
- Frahm J.-P. & Wiethold J. (2004): Die Moosflora des Mittelalters und der Frühen Neuzeit in Mitteleuropa nach archäologischen Funden zusammengestellt. – *Herzogia* 17: 303–324.
- Frank W., Decker E. L. & Reski R. (2005): Molecular tools to study *Physcomitrella patens*. – *Plant Biology* 7: 220–227.
- Frank W., Ratnadewi D. & Reski R. (2005): *Physcomitrella patens* is highly tolerant against drought, salt and osmotic stress. – *Planta* 220: 384–394.
- Frey W. & Stech M. (2005): A morpho-molecular classification of the *Anthocerotophyta* (hornworts). – *Nova Hedwigia* 80: 541–545.
- Frey W. & Stech M. (2005): A morpho-molecular classification of the liverworts (*Hepaticophytina*, *Bryophytina*). – *Nova Hedwigia* 81: 55–78.
- Friedel A. & Müller F. (2004): Bryophytes and lichens as indicators for changes of air pollution in the Serrahn Natural Forest Reserve (Mueritz National Park). – *Herzogia* 17: 279–286.
- Fudali E. (2005): Bryophyte species diversity and ecology in the parks and cemeteries of selected Polish cities. – Agricultural University of Wrocław, Wrocław. [212 pp. + 4 Figs.]

- Fuertes E., Oliván G. & Acón M. (2004): *Drepanocladus* (*Amblystegiaceae*, *Bryopsida*) in the Iberian Peninsula. – Belgian Journal of Botany 136: 167–173.
- Furuki T. & Mizutani M. (2004): Checklist of Japanese *Hepaticae* and *Anthocerotae*, 2004. – Bryological Research 8: 296–316. [In Japanese with English abstract]
- Gabriel R. & Bates J. W. (2005): Bryophyte community composition and habitat specificity in the natural forests of Terceira, Azores. – Plant Ecology 177: 125–144.
- Gallego M. T. (2005): A taxonomic study of the genus *Syntrichia* Brid. (*Pottiaceae*, *Musci*) in the Mediterranean Region and Macaronesia. – Journal of the Hattori Botanical Laboratory 98: 47–122.
- Gallego M. T. (2005): *Pottiaceae: Bryoerythrophyllum*. – In: Guerra J. & Cros R. M. (eds.), Flora Briofítica Ibérica, p. 8–16, Sociedad Española de Briología, Murcia.
- Gallego M. T., Cano M. J. & Sérgio C. (2005): *Syntrichia bogotensis* (*Bryopsida*, *Pottiaceae*) new for Macaronesia. – Bryologist 108: 219–223.
- Gallego M. T., Werner O., Sérgio C. & Guerra J. (2005): A morphological and molecular study of the *Syntrichia laevipila* complex (*Pottiaceae*) in Portugal. – Nova Hedwigia 80: 301–322.
- Ganeva A. & Natcheva R. (2004): An overview of the Bulgarian bryophyte flora: past, present and future. – Phytologia Balcanica 10: 21–26.
- Gang Y.-Y., Du G.-S., Shi D.-J., Wang M.-Z., Li X.-D. & Hua Z.-L. (2003): Establishment of in vitro regeneration system of the *Atrichum* mosses. – Acta Botanica Sinica 45: 1475–1480.
- Gapon S. V. (2004): Stan vychennya mokhovoi roslynnosti v Ukrayini ta osoblyvosti jiji klasyfikaciji. – Ukrains’kyi Botanichnyi Zhurnal 61: 60–66.
- Garcia C., Sérgio C. & Sim-Sim M. (2005): Epiphytic bryophyte diversity in oak woodlands of centre and northern Portugal. First analyses from a national survey. – Journal of the Hattori Botanical Laboratory 97: 161–181.
- Garcia-Pausas J., Casals P. & Romanya J. (2004): Litter decomposition and faunal activity in Mediterranean forest soils: effects of N content and the moss layer. – Soil Biology & Biochemistry 36: 989–997.
- Gardiner A., Ignatov M., Huttunen S. & Troitsky A. (2005): On resurrection of the families *Pseudoleskeaceae* Schimp. and *Pylaisiaceae* Schimp. (*Musci*, *Hypnales*). – Taxon 54: 651–663.
- Garilletti R. (2005): *Pottiaceae: Barbula*. – In: Guerra J. & Cros R. M. (eds.), Flora Briofítica Ibérica, p. 11–18, Sociedad Española de Briología, Murcia.
- Gignac L. D. & Dale M. R. T. (2005): Effects of fragment size and habitat heterogeneity on cryptogam diversity in the low-boreal forest of western Canada. – Bryologist 108: 50–66.
- Giordano S., Adamo P., Sorbo S. & Vingiani S. (2005): Atmospheric trace metal pollution in the Naples urban area based on results from moss and lichen bags. – Environmental Pollution 136: 431–442.
- Gjerde I., Sætersdal M., Rolstad J., Storaunet K. O., Blom H. H., Gundersen V. & Heegaard E. (2005): Productivity-diversity relationships for plants, bryophytes, lichens, and polypore fungi in six northern forest landscapes. – Ecography 28: 705–720.
- Goffinet B. (2004): Bibliography of systematic and population genetic studies in bryophytes based on DNA data. III. – Cryptogamie: Bryologie 25: 385–390.
- Goffinet B., Wickett N. J., Shaw A. J. & Cox C. J. (2005): Phylogenetic significance of the *rpoA* loss in the chloroplast genome of mosses. – Taxon 54: 353–360.
- Goia I. & Schumacker R. (2004): The study of corticolous bryophytes communities from the Arieşul Mare Basin. – Contribuții Botanice 39: 105–114.
- Goldberg I. & Mogensen G. S. (2005): *Gyroweisia groenlandica* (Kindb.) I. Goldberg & Mogensen, comb. nov. for *Weissia groenlandica* Kindb. (*Pottiaceae*, *Bryophyta*). – Lindbergia 30: 39–42.
- Gombert S., Rausch de Traubenberg C., Losno R., Leblond S., Colin J. L. & Cossa D. (2004): Biomonitoring of element deposition using mosses in the 2000 French Survey: Identifying sources and spatial trends. – Journal of Atmospheric Chemistry 49: 479–502.
- Goncharova I. A. & Benkov A. V. (2004): Osobennosti usloviy proizrastaniya mkhov na oligotrofnom bolote [The features of moss habitat conditions in oligotrophic bogs]. Sibirskiy Ekologicheskiy Zhurnal 2: 22–234.
- González-Mancebo J. M., Romaguera F., Losada-Lima A. & Suarez A (2004): Epiphytic bryophytes growing on *Laurus azorica* (Seub.) Franco in three laurel forest areas in Tenerife (Canary Islands). – Acta Oecologica – International Journal of Ecology 25: 159–167.
- Gorecka H. (2003): Bioindicators of environmental pollution with metals. – Przemysł Chemiczny 82: 1081–1083.

- Gorelick R. (2005): Theory for why dioecious plants have equal length sex chromosomes. – American Journal of Botany 92: 979–984.
- Goremykin V. V. & Hellwig F. H. (2005): Evidence for the most basal split in land plants dividing bryophyte and tracheophyte lineages. – Plant Systematics and Evolution 254: 93–103. [Erratum p. 119]
- Gradstein S. R., Tan B. C., Zhu R. L., Ho B. C., King C. S. H., Drubert C. & Pitopang R. (2005): A catalogue of the Bryophytes of Sulawesi, Indonesia. – Journal of the Hattori Botanical Laboratory 98: 213–257.
- Gremmen N. J. M., Smith V. R. & van Tongeren O. F. R. (2003): Impact of trampling on the vegetation of subantarctic Marion Island. – Arctic, Antarctic and Alpine Research 35: 442–446.
- Grodzińska K., Szarek-Łukaszewska G., Frontasyeva M., Pavlov S. S. & Gudorina S. F. (2005): Multielement concentration in mosses in the forest influenced by industrial emissions (Niepolomice Forest, S Poland) at the end of the 20<sup>th</sup> century. – Polish Journal of Environmental Studies 14: 165–172.
- Groeneveld E. V. G. & Rochefort L. (2005): *Polytrichum strictum* as a solution to frost heaving in disturbed ecosystems: A case study with milled peatlands. – Restoration Ecology 13: 74–82.
- Grolle R., Long D. G. & McNeill J. (2005): Nomenclatural problems in *Pleurozia* (*Pleuroziaceae*): the lectotypification of *Pleurozia* Dumort., *Jungermannia sphagnoides* Schwägr., *Pleurozia sphagnoides* Dumort., *P. purpurea* Lindb., *Mnium jungermannia* L. and *Jungermannia undulata* L. (*Hepaticae*). – Taxon 54: 503–508.
- Grolle R. & Meister K. (2004): *Lophozia kutscheri*, a new hepatic (*Jungermanniales*) in Bitterfeld amber from central Germany. – Bryologist 107: 79–81.
- Grolle R. & Meister K. (2004): The liverworts in Baltic and Bitterfeld amber. – Weissdorn-Verlag, Jena. [91 pp.]
- Grolle R. & So M. L. (2004): A fossil *Porella* (*Porellaceae*, *Hepaticae*) in Eocene Baltic amber. – Botanical Journal of the Linnean Society 145: 485–488.
- Groth-Malonek M., Heinrichs J., Schneider H. & Gradstein S. R. (2004): Phylogenetic relationships in the *Lejeuneaceae* (*Hepaticae*) inferred using ITS sequences of nuclear ribosomal DNA. – Organisms Diversity & Evolution 4: 51–57.
- Groth-Malonek M. & Knoop V. (2005): Bryophytes and other basal land plants: the mitochondrial perspective. – Taxon 54: 293–297.
- Groth-Malonek M., Pruchner D., Grewe F. & Knoop V. (2005): Ancestors of *trans*-splicing mitochondrial introns support serial sister group relationships of hornworts and mosses with vascular plants. – Molecular Biology and Evolution 22: 117–125.
- Grünfeld K. (2005): Integrating spatio-temporal information in environmental monitoring data – a visualization approach applied to moss data. – Science of the Total Environment 347: 1–20.
- Guerra J. & Brugués M. (2005): *Pottiaceae: Anoectangium*. – In: Guerra J. & Cros R. M. (eds.), Flora Briofítica Ibérica, p. 8–10, Sociedad Española de Briología, Murcia.
- Guerra J. (2005): *Pottiaceae: Leptodontium*. – In: Guerra J. & Cros R. M. (eds.), Flora Briofítica Ibérica, p. 18–20, Sociedad Española de Briología, Murcia.
- Guerra J. (2005): *Pottiaceae: Triquetrella*. – In: Guerra J. & Cros R. M. (eds.), Flora Briofítica Ibérica, p. 20–22, Sociedad Española de Briología, Murcia.
- Gunnarsson U. (2005): Global patterns of *Sphagnum* productivity. – Journal of Bryology 27: 269–280.
- Gunnarsson U., Hassel K. & Söderström L. (2005): Genetic structure of the endangered peat moss *Sphagnum angermanicum* in Sweden: a result of historic or contemporary processes? – Bryologist 108: 194–203.
- Gustafsson L., Appelgren L. & Nordin A. (2005): Biodiversity value of potential forest fertilisation stands, as assessed by red-listed and ‘signal’ bryophytes and lichens. – Silva Fennica 39: 191–200.
- Gutman A. B. & Crawford G. H. (2005): Botanical briefs: Liverworts – *Frullania* species. – Cutis 75: 262–264.
- Haberl H., Schulz N. B., Plutzar C., Erb K. H., Krausmann F., Loibl W., Moser D., Sauberer N., Weisz H., Zechmeister H. G. & Zulka P. (2004): Human appropriation of net primary production and species diversity in agricultural landscapes. – Agriculture, Ecosystems & Environment 102: 213–218.
- Haeussler S. & Bergeron Y. (2004): Range of variability in boreal aspen plant communities after wildfire and clear-cutting. – Canadian Journal of Forest Research – Revue Canadienne de Recherche Forestière 34: 274–288.
- Hall B. D. & Louis V. L. St. (2004): Methylmercury and total mercury in plant litter decomposing in upland forests and flooded landscapes. – Environmental Science & Technology 38: 5010–5021.

- Hans F. (2004): Neue und seltene Arten aus der Familie *Orthotrichaceae (Musci)* für Luxemburg – Mögliche Indikatoren für einen Klimawechsel? – Bulletin de la Société des Naturalistes Luxembourgeois 105: 15–25.
- Harmens H., Buse A., Büker P., Norris D., Mills G., Williams B., Reynolds B., Ashenden T. W., Röhling Å. & Steinnes E. (2004): Heavy metal concentrations in European mosses: 2000/2001 survey. – Journal of Atmospheric Chemistry 49: 425–436.
- Hassel K., Pedersen B. & Söderström L. (2005): Age and size at maturity in mountain and lowland populations of the expanding moss *Pogonatum dentatum*. – Plant Ecology 179: 207–216.
- Hassel K., Pedersen B. & Söderström L. (2005): Changes in life-history traits in an expanding moss species: phenotypic plasticity or genetic differentiation? A reciprocal transplantation experiment with *Pogonatum dentatum*. – Ecography 28: 71–80.
- Hassel K., Såstad S. M., Gunnarsson U. & Söderström L. (2005): Genetic variation and structure in the expanding moss *Pogonatum dentatum (Polytrichaceae)* in its area of origin and in a recently colonized area. – American Journal of Botany 92: 1684–1690.
- Hassel K. & Söderström L. (2005): The expansion of the alien mosses *Orthodontium lineare* and *Campylopus introflexus* in Britain and continental Europe. – Journal of the Hattori Botanical Laboratory 97: 183–193.
- Hasselbach L., Ver Hoef J. M., Ford J., Neitlich P., Crecelius E., Berryman S., Wolk B. & Bohle T. (2005): Spatial patterns of cadmium and lead deposition on and adjacent to National Park Service lands in the vicinity of Red Dog Mine, Alaska. – Science of the Total Environment 348: 211–230.
- Haugland J. E. & Beatty S. W. (2005): Vegetation establishment, succession and microsite frost disturbance on glacier forelands within patterned ground chronosequences. – Journal of Biogeography 32: 145–153.
- Hautala H., Jalonen J., Laaka-Lindberg S. & Vanha-Majamaa I. (2004): Impacts of retention felling on coarse woody debris (CWD) in mature boreal spruce forests in Finland. – Biodiversity and Conservation 13: 1541–1554.
- Hébrard J. P. (2005): Contribution to the study of the bryoflora of the high valleys of the Verdon (Alpes-de-Haute-Provence) and Var (Alpes-Maritimes). – Cryptogamie: Bryologie 26: 183–207.
- Hébrard J.-P. (2005): New localities and ecology of *Buxbaumia viridis* (*Buxbaumiaceae, Musci*) in Corsica. – Nova Hedwigia 80: 241–246.
- Hedenäs L. (2005): Bryophyte flora of Uganda. 4. *Rhytidaceae, Hylocomiaceae* and *Hypnaceae* (Part 1). – Journal of Bryology 27: 55–66.
- Hedenäs L. (2005): *Onchophorus wahlenbergii* var. *elongatus* I. Hagen, an overlooked taxon in northern Europe. – Lindbergia 30: 32–38.
- Hedenäs L. & Bisang I. (2004): Key to the European *Dicranum* species. – Herzogia 17: 179–197.
- Hedenäs L., Bisang I., Tehler A., Hammarberg C., Hamnede M., Jaederfelt K. & Odelvik G. (2003): Hur våra herbarier kan användas för att bedöma om arter blivit vanligare eller mer sällsynta [A method for evaluating temporal frequency variation, based on herbarium material]. – Svensk Botanisk Tidskrift 97: 225–236.
- Hedenäs L., Oliván G. & Eldenäs P. (2005): Phylogeny of the *Calliergonaceae (Bryophyta)* based on molecular and morphological data. – Plant Systematics and Evolution 252: 49–61.
- Hedenäs L. & Watling M. C. (2005): Bryophyte flora of Uganda. 5. *Hypnaceae* (Part 2). – Journal of Bryology 27: 153–160.
- Hedges S. B., Blair J. E., Venturi M. L. & Shoe J. L. (2004): A molecular timescale of eukaryote evolution and the rise of complex multicellular life. – BMC Evolutionary Biology 4: 2.
- Heilmann-Clausen J., Aude E. & Christensen M. (2005): Cryptogam communities on decaying deciduous wood – does tree species diversity matter? – Biodiversity and Conservation 14: 2061–2078.
- Heinken T. & Zippel E. (2004): Natural re-colonization of experimental gaps by terricolous bryophytes in Central European pine forests. – Nova Hedwigia 79: 329–351.
- Heino J., Paavola R., Virtanen R. & Muotka T. (2005): Searching for biodiversity indicators in running waters: do bryophytes, macroinvertebrates, and fish show congruent diversity patterns? – Biodiversity and Conservation 14: 415–428.
- Heino J., Virtanen R., Vuori K. M., Saastamoinen J., Ohtonen A. & Muotka T. (2005): Spring bryophytes in forested landscapes: Land use effects on bryophyte species richness, community structure and persistence. – Biological Conservation 124: 539–545.
- Heinrichs J., Gradstein S. R., Wilson R. & Schneider H. (2005): Towards a natural classification of liverworts (*Marchantiophyta*) based on the chloroplast gene *rbcL*. – Cryptogamie: Bryologie 26: 131–150.

- Heinrichs J., Groth H., Lindner M., Feldberg K. & Rycroft D. S. (2004): Molecular, morphological, and phytochemical evidence for a broad species concept of *Plagiochila bifaria* (*Hepaticae*). – *Bryologist* 107: 28–40.
- Heinrichs J., Groth H., Lindner M., Renker C., Pócs T. & Pröschold T. (2004): Intercontinental distribution of *Plagiochila corrugata* (*Plagiochilaceae*, *Hepaticae*) inferred from nrDNA ITS sequences and morphology. – *Botanical Journal of the Linnean Society* 146: 469–481.
- Heinrichs J., Groth H. & Sauer M. (2004): New synonyms in *Plagiochila* (*Hepaticae*) III. – *Cryptogamie: Bryologie* 25: 35–37.
- Heinrichs J., Lindner M., Gradstein S. R., Groth H., Buchbender V., Solga A. & Fischer E. (2005): Origin and subdivision of *Plagiochila* (*Jungmanniidae*: *Plagiochilaceae*) in tropical Africa based on evidence from nuclear and chloroplast DNA sequences and morphology. – *Taxon* 54: 317–333.
- Heinrichs J., Lindner M. & Groth H. (2004): Sectional classification of the North American *Plagiochila* (*Hepaticae*, *Plagiochilaceae*). – *Bryologist* 107: 489–496.
- Heinrichs J., Lindner M., Groth H. & Renker C. (2005): Distribution and synonymy of *Plagiochila punctata* (Taylor) Taylor, with hypotheses on the evolutionary history of *Plagiochila* sect. *Arrectae* (*Plagiochilaceae*, *Hepaticae*). – *Plant Systematics and Evolution* 250: 105–117.
- Herrnstadt I. & Kidron G. J. (2005): Reproductive strategies of *Bryum dunense* in three microhabitats in the Negev desert. – *Bryologist* 108: 101–109.
- Hess S., Frahm J.-P. & Theisen I. (2005): Evidence of zoophagy in a second liverwort species, *Pleurozia purpurea*. – *Bryologist* 108: 212–218.
- Heylen O., Hermy M. & Schrevens E. (2005): Determinants of cryptogamic epiphyte diversity in a river valley (Flanders). – *Biological Conservation* 126: 371–382.
- Higuchi S., Kawamura M., Miyajima I., Akiyama H., Kosuge K., Kato M. & Nozaki H. (2003): Morphology and phylogenetic position of a mat-forming green plant from acidic rivers in Japan. – *Journal of Plant Research* 116: 461–467.
- Higuchi M. & Shan-Hsiung Lin (2004): Correlations between the moss floras of Japan and Taiwan. – *National Science Museum Monographs* 24: 227–240.
- Hock Zs., Szővényi P. & Tóth Z. (2004): Seasonal variation in the bryophyte diaspore bank of open grasslands on dolomite rock. – *Journal of Bryology* 26: 285–292.
- Hohe A. & Reski R. (2005): From axenic spore germination to molecular farming: One century of bryophyte in vitro culture. – *Plant Cell Reports* 23: 513–521.
- Hokkanen P. J. (2004): Bryophyte communities in herb-rich forests in Koli, eastern Finland: comparison of forest classifications based on bryophytes and vascular plants. – *Annales Botanici Fennici* 41: 331–365.
- Hollister R. D., Webber P. J., Tweedie C. E. (2005): The response of Alaskan arctic tundra to experimental warming: differences between short- and long-term responses. – *Global Change Biology* 11: 525–536.
- Holyoak D. T. (2004): Taxonomic notes on some European species of *Bryum* (*Bryopsida*: *Bryaceae*). – *Journal of Bryology* 26: 247–264.
- Holyoak D. T. & Bryan V. S. (2005): *Ephemereum hibernicum* sp. nov. (*Bryopsida*: *Ephemeraceae*) from Ireland. – *Journal of Bryology* 27: 89–96.
- Holz I. & Gradstein S. R. (2005): Cryptogamic epiphytes in primary and recovering upper montane oak forests of Costa Rica – species richness, community composition and ecology. – *Plant Ecology* 178: 89–109.
- Holz I. & Gradstein S. R. (2005): Phytogeography of the bryophyte floras of oak forests and páramo of the Cordillera de Talamanca, Costa Rica. – *Journal of Biogeography* 32: 1591–1609.
- Hotes S., Poschlod P., Takahashi H., Grootjans A. P. & Adema E. (2004): Effects of tephra deposition on mire vegetation: a field experiment in Hokkaido, Japan. – *Journal of Ecology* 92: 624–634.
- Huang J.-X., Qu L.-J., Yang J., Yin H. & Gu H.-Y. (2004): A preliminary study on the origin and evolution of chalcone synthase (CHS) gene in angiosperms. – *Acta Botanica Sinica* 46: 10–19.
- Hugonnot V. (2004): Contribution to the chorology and ecology of *Ricciaceae* in the Massif Central (France), chiefly in the Ardèche and Haute-Loire departments. – *Cryptogamie: Bryologie* 25: 39–67.
- Hugonnot V. (2004): *Sphagnum obtusum* Warnst. in France. – *Cryptogamie: Bryologie* 25: 357–365.
- Huttunen S., Lappalainen N. M. & Turunen J. (2005): UV-absorbing compounds in subarctic herbarium bryophytes. – *Environmental Pollution* 133: 303–314.
- Huttunen S., Taipale T., Lappalainen N. M., Kubin E., Lakkala K. & Kaurola J. (2005): Environmental specimen bank samples of *Pleurozium schreberi* and *Hylocomium splendens* as indicators of the radiation environment at the surface. – *Environmental Pollution* 133: 315–326.

- Hylander K. (2005): Aspect modifies the magnitude of edge effects on bryophyte growth in boreal forests. – *Journal of Applied Ecology* 42: 518–525.
- Hylander K., Dynesius M., Jonsson B. G. & Nilsson C. (2005): Substrate form determines the fate of bryophytes in riparian buffer strips. – *Ecological Applications* 15: 674–688.
- Ignatov M. S., Huttunen S. & Koponen T. (2005): Bryophyte flora of Hunan Province, China. 5. *Brachytheciaceae* (*Musci*), with an overview of *Eurhynchadelphus* and *Rhynchostegiella* in SE Asia. – *Acta Botanica Fennica* 178: 1–56.
- Ignatov M., Tsegmed T., Tan B. I., Bai X. L. & Zolotov V. (2004): Mosses of Gobi in Mongolia. – *Journal of the Hattori Botanical Laboratory* 96: 183–210.
- Ignatova E. & Muñoz J. (2004): The genus *Grimmia* Hedw. (*Grimmiaceae*, *Musci*) in Russia. – *Arctoa* 13: 101–182.
- Infante M. & Heras P. (2005): *Ephememerum cohaerens* (Hedw.) Hampe and *E. spinulosum* Bruch & Schimp. (*Ephemeraceae*, *Bryopsida*), new to the Iberian Peninsula. – *Cryptogamie: Bryologie* 26: 327–333.
- Ingerpuu N., Liira J. & Pärtel M. (2005): Vascular plants facilitated bryophytes in a grassland experiment. – *Plant Ecology* 180: 69–75.
- Jägerbrand A. K. (2004): Patterns of species richness and vegetative performance in heath ecosystems at Thingvellir, southwest Iceland. – *Icelandic Agricultural Sciences* 16–17: 29–38.
- Jägerbrand A. K. (2005): Subarctic bryophyte ecology: phenotypic variation and responses to simulated environmental change. – Ms. [PhD thesis, Göteborg University]
- Jankowiak K., Lesicka J., Pacak A., Rybarczyk A. & Szwejkowska-Kulińska Z. (2004): A comparison of group II introns of plastid tRNA<sup>Lys</sup><sub>UUU</sub> genes encoding maturase protein. – *Cellular & Molecular Biology Letters* 9: 239–251.
- Jankowiak K., Rybarczyk A., Wyatt R., Odrzykoski I., Pacak A. & Szwejkowska-Kulińska Z. (2005): Organellar inheritance in the allopolyploid moss *Rhizomnium pseudopunctatum*. – *Taxon* 54: 383–388.
- Jiménez J. A. (2004): *Pottiaceae: Didymodon*. – In: Guerra J. & Cros R. M. (eds.), *Flora Briofítica Ibérica*, p. 1–35, Sociedad Española de Briología, Murcia.
- Jiménez J. A., Ros R. M., Cano M. J. & Guerra J. (2004): New distribution data on *Didymodon anserinocapitatus* (X.-J. Li) R. H. Zander, *D. maschalogenia* (Renauld & Cardot) Broth. and *D. sicculus* M. J. Cano et al. (*Bryophyta*, *Pottiaceae*). – *Cryptogamie: Bryologie* 25: 91–97.
- Jiménez J. A., Ros R. M., Cano M. J. & Guerra J. (2005): A new evaluation of the genus *Trichostomopsis* (*Pottiaceae*, *Bryophyta*). – *Botanical Journal of the Linnean Society* 147: 117–127.
- Jiménez J. A., Ros R. M., Cano M. J. & Guerra J. (2005): A revision of *Didymodon* section *Fallaces* (*Musci*, *Pottiaceae*) in Europe, North Africa, Macaronesia, and Southwest and Central Asia. – *Annals of the Missouri Botanical Garden* 92: 225–247.
- Johri R. M., Lata S. & Tyagi K. (2004): A textbook of bryophyta. – New Delhi: Dominant Publishers and Distributors. [324 pp.]
- Jónsdóttir I. S., Magnússon B., Gudmundsson J., Elmarsdóttir A. & Hjartarson H. (2005): Variable sensitivity of plant communities in Iceland to experimental warming. – *Global Change Biology* 11: 553–563.
- Juslén A. (2004): Phylogeny of isophyllous liverworts. – *Cladistics* 20: 597–598.
- Kantvilas G. & Jarman S. J. (2004): Lichens and bryophytes on *Eucalyptus obliqua* in Tasmania: management implications in production forests. – *Biological Conservation* 117: 359–373.
- Kato M. & Akiyama H. (2005): Interpolation hypothesis for origin of the vegetative sporophyte of land plants. – *Taxon* 54: 443–450.
- Keçeli T. & Çetin B. (2005): *Ptilidium pulcherrimum* (*Ptilidiaceae*, *Hepaticae*) new to South-West Asia. – *Cryptogamie: Bryologie* 26: 313–317.
- Kelly D. L., O'Donovan G., Feehan J., Murphy S., Drangeid S. O. & Marcano-Berti L. (2004): The epiphyte communities of a montane rain forest in the Andes of Venezuela: patterns in the distribution of the flora. – *Journal of Tropical Ecology* 20: 643–666.
- Kimmerer R. W. (2005): Patterns of dispersal and establishment of bryophytes colonizing natural and experimental treefall mounds in northern hardwood forests. – *Bryologist* 108: 391–401.
- Kis G. (2004): Non European Bryophyta types and list of exsiccata in the Eger Cryptogamic Herbarium (EGR). – *Folia Historico-Naturalia Musei Matraensis* 28: 5–52.
- Klama H. 2004. *Scapania parvifolia* (*Marchantiopsida*, *Scapaniaceae*) – nowy gatunek wątrobowca dla flory Karkonoszy [*Scapania parvifolia* (*Marchantiopsida*, *Scapaniaceae*) – a new liverwort of the flora of Karkonosze Mts.]. – *Fragmenta Floristica et Geobotanica Polonica* 11: 213–215.

- Klanderud K. & Totland Ø. (2004): Habitat dependent nurse effects of the dwarf-shrub *Dryas octopetala* on alpine and arctic plant community structure. – *Ecoscience* 11: 410–420.
- Klanderud K. & Totland Ø. (2005): Simulated climate change altered dominance hierarchies and diversity of an alpine biodiversity hotspot. – *Ecology* 86: 2047–2054.
- Klazenga N. (2005): Generic concepts in Australian mosses. – *Australian Systematic Botany* 18: 17–23.
- Koch K., Neinhuis C., Ensikat H.-J. & Barthlott W. (2004): Self assembly of epicuticular waxes on living plant surfaces imaged by atomic force microscopy (AFM). – *Journal of Experimental Botany* 55: 711–718.
- Köckinger H., Kučera J. & Stebel A. (2005): *Pohlia nutans* subsp. *schimperi* (Müll.Hal.) Nyholm, a neglected Nordic moss in Central Europe. – *Journal of Bryology* 27: 351–355.
- Koponen T. (2005): Invited review: Viktor Ferdinand Brotherus (1849–1929) and *Musci* in Engler & Prantl's *Die Natürlichen Pflanzenfamilien*. – *Bryologist* 108: 345–362.
- Koponen T., Cao T., Huttunen S., Juslén A., Peng C., Piippo S., Rao P., Váňa J. & Virtanen V. (2004): Bryophyte flora of Hunan Province, China. 3. Bryophytes from Taoyuandong and Yankou Nature Reserves and Badagongshan and Hupingshan National Nature Reserves, with additions to floras of Mangshan Nature Reserve and Wulingyuan Cultural Heritage Area. – *Acta Botanica Fennica* 177: 1–47.
- Korpelainen H., Pohjamo M. & Laaka-Lindberg S. (2005): How efficiently does bryophyte dispersal lead to gene flow? – *Journal of the Hattori Botanical Laboratory* 97: 195–205.
- Kosiba P. (2004): Biomonitoring of air and reactions of mosses in conditions of pollution with industry emissions. – Wydawnictwo Uniwersytetu Wrocławskiego, Prace Botaniczne, 80. [110 pp.]
- Kottke I. & Nebel M. (2005): The evolution of mycorrhiza-like associations in liverworts: an update. – *New Phytologist* 167: 330–334.
- Krayesky D. M., Crandall-Stotler B. & Stotler R. E. (2005): A revision of the genus *Fossombronia* Raddi in East Asia and Oceania. – *Journal of the Hattori Botanical Laboratory* 98: 1–45.
- Kremer C. L. & Drinnan A. N. (2004): Secondary walls in hyaline cells of *Sphagnum*. – *Australian Journal of Botany* 52: 243–256.
- Krisai R. & Strobl W. (2005): Zur Verbreitung des Laubmooses *Hookeria lucens* in Österreich und dem angrenzenden Südbayern. – *Herzogia* 18: 211–218.
- Krishna M. V. B., Karunasagar D. & Arunachalam J. (2004): Sorption characteristics of inorganic, methyl and elemental mercury on lichens and mosses: Implication in biogeochemical cycling of mercury. – *Journal of Atmospheric Chemistry* 49: 317–328.
- Kubešová S. & Chytrý M. (2005): Diversity of bryophytes on treeless cliffs and talus slopes in a forested central European landscape. – *Journal of Bryology* 27: 35–46.
- Kučera J. & Holyoak D. T. (2005): Lectotypification of *Bryum moravicum* Podp. (*Bryopsida*: *Bryaceae*). – *Journal of Bryology* 27: 161–162.
- Kürschner H. (2004): Life strategies and adaptations in bryophytes from the Near and Middle East. – *Turkish Journal of Botany* 28: 73–84.
- Kürschner H. (2004): Phytosociology, ecology and life strategies of a remarkable epiphytic bryophyte community of Socotra Island, Yemen. – *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 125: 377–395.
- Kürschner H. & Ochyra R. (2004): Remarkable new records to the bryophyte flora of Yemen (al-Mahra and Socotra Island). Additions to the Bryophyte Flora of the Arabian Peninsula and Socotra 6. – *Cryptogamie: Bryologie* 25: 69–81.
- Kürschner H. & Parolly G. (2004): Phytomass and water-storing capacity of epiphytic rain forest bryophyte communities in S Ecuador. Ecosociological studies in Ecuadorian bryophyte communities IV. – *Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie* 125: 489–504.
- Kürschner H. & Parolly G. (2005): Ecosociological studies in Ecuadorian bryophyte communities III. Life forms, life strategies and ecomorphology of the submontane and montane epiphytic vegetation of S Ecuador. – *Nova Hedwigia* 80: 89–113.
- Kürschner H. & Wagner D. (2005): Phytosociology and life strategies of a new loess slope bryophyte community from N China (Gansu), including *Crossidium laxofilamentosum* new to China. – *Nova Hedwigia* 81: 229–246.
- Kushida K., Kim Y., Tanaka N. & Fukuda M. (2004): Remote sensing of net ecosystem productivity based on component spectrum and soil respiration observation in a boreal forest, interior Alaska. – *Journal of Geophysical Research – Atmospheres* 109: D06108.
- Kusumi J. & Tachida H. (2005): Compositional properties of green-plant plastid genomes. – *Journal of Molecular Evolution* 60: 417–425.

- Kutschera U. & Koopmann V. (2005): Growth in liverworts of the *Marchantiales* is promoted by epiphytic methylobacteria. – *Naturwissenschaften* 92: 347–349.
- Kuusk A., Lang M. & Nilson T. (2004): Simulation of the reflectance of ground vegetation in sub-boreal forests. – *Agricultural and Forest Meteorology* 126: 33–46.
- Laaka-Lindberg S. (2005): Reproductive phenology in the leafy hepatic *Lophozia silvicola* Buch in southern Finland. – *Journal of Bryology* 27: 253–260.
- Laaka-Lindberg S., Pohjamäo M. & Korpelainen H. (2005): Niche breadth and niche overlap in three epixylic hepatics in a boreal old-growth forest, southern Finland. – *Journal of Bryology* 27: 119–127.
- Lachance D., Lavoie C. & Desrochers A. (2005): The impact of peatland afforestation on plant and bird diversity in southeastern Québec. – *Ecoscience* 12: 161–171.
- Lamentowicz M. & Mitchell E. A. D. (2005): The ecology of testate amoebae (protists) in *Sphagnum* in North-Western Poland in relation to peatland ecology. – *Microbial Ecology* 50: 48–63.
- Lang D., Eisinger J., Reski R. & Rensing S. A. (2005): Representation and high-quality annotation of the *Physcomitrella patens* transcriptome demonstrates a high proportion of proteins involved in metabolism in mosses. – *Plant Biology* 7: 238–250.
- Lara F. (2005): *Pottiaceae: Dialytrichia*. – In: Guerra J. & Cros R. M. (eds.), *Flora Briofítica Ibérica*, p. 22–27. Sociedad Española de Briología, Murcia.
- Lara F., Puntillo D., Garilleti R. & Mazimpaka V. (2004): *Orthotrichum gymnostomum* Brid. and other interesting *Orthotrichaceae* from Calabria (Italy). – *Cryptogamie: Bryologie* 25: 349–356.
- Lariguet P. & Dunand C. (2005): Plant photoreceptors: Phylogenetic overview. – *Journal of Molecular Evolution* 61: 559–569.
- Latorre A. V. P., Cano M. J., Cabezudo B. & Guerra J. (2005): Phytocoenological behaviour, distribution and conservation of *Trichomanes speciosum* Willd. (*Pteridophyta*) in the south of the Iberian Peninsula. – *Cryptogamie: Bryologie* 26: 249–261.
- Lauer H. (2005): Die Moose der Pfalz. – Pollichia, Bad Dürkheim – Buch Nr. 46. [1219 pp.]
- Leblond S., Gombert S., Colin J. L., Losno R. & Rausch de Traubenberg C. (2004): Biological and temporal variations of trace element concentrations in the moss species *Scleropodium purum* (Hedw.) Limpr. – *Journal of Atmospheric Chemistry* 49: 95–110.
- Lee C. S. L., Li X.-D., Zhang G., Peng X.-Z. & Zhang L. (2005): Biomonitoring of trace metals in the atmosphere using moss (*Hypnum plumaeforme*) in the Nanling Mountains and the Pearl River Delta, Southern China. – *Atmospheric Environment* 39: 397–407.
- Lee K. J. D., Knight C. D. & Knox J. P. (2005): *Physcomitrella patens*: A moss system for the study of plant cell walls. – *Plant Biosystems* 139: 16–19.
- Leitch I. J., Soltis D. E., Soltis P. S. & Bennett M. D. (2005): Evolution of DNA amounts across land plants (*Embryophyta*). – *Annals of Botany* 95: 207–217.
- Lewis Smith R. I. (2005): Bryophyte diversity and ecology of two geologically contrasting Antarctic islands. – *Journal of Bryology* 27: 195–206.
- Lewis Smith R. I. (2005): The thermophilic bryoflora of Deception Island: unique plant communities as a criterion for designating an Antarctic Specially Protected Area. – *Antarctic Science* 17: 17–27.
- Liiv S. & Kaasik M. (2004): Trace metals in mosses in the Estonian oil shale processing region. – *Journal of Atmospheric Chemistry* 49: 563–578.
- Limpens J., Berendse F. & Klees H. (2004): How phosphorus availability affects the impact of nitrogen deposition on *Sphagnum* and vascular plants in bogs. – *Ecosystems* 7: 793–804.
- Lobachevska O., Kyjak N., Khorkavtsiv O., Dovgalyuk A., Kit N., Klyuchivska O., Stoika R., Ripetsky R. & Cove D. (2005): Influence of metabolic stress on the inheritance of cell determination in the moss, *Pottia intermedia*. – *Cell Biology International* 29: 181–186.
- Löbel S., Dengler J. & Hobohm C. (2004): Beziehungen zwischen der Artenvielfalt von Gefäßpflanzen, Moosen und Flechten in Trockenrasen der Insel Öland (Schweden). – *Kieler Notizen zur Pflanzenkunde in Schleswig-Holstein und Hamburg* 32: 9–13.
- Locky D. A., Bayley S. E. & Vitt D. H. (2005): The vegetational ecology of black spruce swamps, fens, and bogs in southern boreal Manitoba, Canada. – *Wetlands* 25: 564–582.
- Long D. G. (2005): Studies on the genus *Asterella* (Acytiaceae) VI: Infrageneric classification in *Asterella*. – *Journal of the Hattori Botanical Laboratory* 97: 249–261.
- Lucaciu A., Timofte L., Culicov O., Frontasyeva M. V., Oprea C., Cucu-Man S., Mocanu R. & Steinnes E. (2004): Atmospheric deposition of trace elements in Romania studied by the moss biomonitoring technique. – *Journal of Atmospheric Chemistry* 49: 533–548.

- Luís L., Draper D. & Sim-Sim M. (2005): The distribution of the genus *Radula* in mainland Portugal and the Madeira Archipelago. – *Lindbergia* 30: 3–10.
- Lüth M. (2003): *Uloota hutchinsiae* – Ökologie und Vergesellschaftung einer fast verschollenen Art. – *Herzogia* 16: 207–220.
- Lüth M. & Schäfer-Verwimp A. (2004): Additions to the bryophyte flora of the Neotropics. – *Tropical Bryology* 25: 7–17.
- Maier E. & Price M. J. (2005): In the footsteps of Lantzius-Beninga: investigating the peristome characters of mosses. – *Bryologist* 108: 36–46.
- Maksimov A. I., Potemkin A. D., Hokkainen T. J. & Maksimova T. A. (2003): Bryophytes of fragmented old-growth spruce forest stands of the North Karelian Biosphere Reserve and adjacent areas of Finland. – *Arctoa* 12: 9–23.
- Malmer N. & Wallén B. (2005): Nitrogen and phosphorus in mire plants: variation during 50 years in relation to supply rate and vegetation type. – *Oikos* 109: 539–554.
- Manzke W. (2004): Zur Verbreitung und Bestandssituation von *Notothylas orbicularis*, *Anthoceros neesii*, *Anthoceros agrestis*, *Phaeoceros carolinianus* und *Riccia ciliata* auf Stoppelfeldern im Vogelsberg (Hessen). – *Hessische Floristische Briefe* 53: 53–65.
- Manzke W. (2005): Zur Verbreitung und Gefährdung von *Notothylas orbicularis* und *Anthoceros neesii* im Vogelsberg (Hessen). – *Bryologische Rundbriefe* 86: 1–3.
- Martin P. H., Sherman R. E. & Fahey T. J. (2004): Forty years of tropical forest recovery from agriculture: Structure and floristics of secondary and old-growth riparian forests in the Dominican Republic. – *Biotropica* 36: 297–317.
- Martínez-Abaigar J., Núñez-Olivera E., Matcham H. W. & Duckett J. G. (2005): Interactions between parasitic fungi and mosses: pegged and swollen-tipped rhizoids in *Funaria* and *Bryum*. – *Journal of Bryology* 27: 47–53.
- Mayaba N. & Beckett R. P. (2003): Increased activities of superoxide dismutase and catalase are not the mechanism of desiccation tolerance induced by hardening in the moss *Atrichum androgynum*. – *Journal of Bryology* 25: 281–286.
- McDaniel S. F. & Shaw A. J. (2005): Selective sweeps and intercontinental migration in the cosmopolitan moss *Ceratodon purpureus* (Hedw.) Brid. – *Molecular Ecology* 14: 1121–1132.
- McGee G. G. & Kimmerer R. W. (2004): Size of *Acer saccharum* hosts does not influence growth of mature bryophyte gametophytes in Adirondack northern hardwood forests. – *Bryologist* 107: 302–311.
- Meister K. & Liebert H. P. (2004): Eine neue Methode zur Ökosystembewertung eines Hochmoors anhand der Erfassung der Torfmoos-, „Vitalität“. – *Herzogia* 17: 287–302.
- Mekuria T., Steiner U., Hindorf H., Frahm J.-P. & Dehme H. W. (2005): Bioactivity of bryophyte extracts against *Botrytis cinerea*, *Alternaria solani* and *Phytophthora infestans*. – *Journal of Applied Botany and Food Quality – Angewandte Botanik* 79: 89–93.
- Melosik I., Odrzykoski I. I. & Śliwińska E. (2005): Delimitation of taxa of *Sphagnum subsecundum* s.l. (*Musci, Sphagnaceae*) based on multienzyme phenotype and cytological characters. – *Nova Hedwigia* 80: 397–412.
- Melosik I. & Såstad S. M. (2005): In vitro propagation of selected *Sphagnum* species (section *Subsecunda*). – *Lindbergia* 30: 21–31.
- Mills S. E. & Macdonald S. E. (2004): Predictors of moss and liverwort species diversity of microsites in conifer-dominated boreal forest. – *Journal of Vegetation Science* 15: 189–198.
- Mills S. E. & Macdonald S. E. (2005): Factors influencing bryophyte assemblage at different scales in the western Canadian boreal forest. – *Bryologist* 108: 86–100.
- Minami A., Nagao M., Ikegami K., Koshiba T., Arakawa K., Fujikawa S. & Takezawa D. (2005): Cold acclimation in bryophytes: low-temperature-induced freezing tolerance in *Physcomitrella patens* is associated with increases in expression levels of stress-related genes but not with increase in level of endogenous abscisic acid. – *Planta* 220: 414–423.
- Mishler B. (2004): The evolution and ecology of desiccation tolerance in mosses. – *Integrative and Comparative Biology* 44: 608–608.
- Mitchell R. J., Truscott A. M., Leith I. D., Cape J. N., van Dijk N., Tang Y. S., Fowler D. & Sutton M. A. (2005): A study of the epiphytic communities of Atlantic oak woods along an atmospheric nitrogen deposition gradient. – *Journal of Ecology* 93: 482–492.
- Moret J.-L. (2005): Les publications de Charles Meylan (1868 – 1941). – *Meylania* 31: 13–23.

- Mulholland P. J. (2004): The importance of in-stream uptake for regulating stream concentrations and outputs of N and P from a forested watershed: evidence from long-term chemistry records for Walker Branch Watershed. – *Biogeochemistry* 70: 403–426.
- Nagy J., Németh N. & Figecky G. (2003): Dynamics of *Sphagnum* cushions in the willow carr of Benceótó mire (NE Hungary) and its nature conservation aspects. – *Polish Botanical Journal* 48: 163–169.
- Nakata M., Watanabe Y., Sakurai Y., Hashimoto Y., Matsuzaki M., Takahashi Y. & Satoh T. (2004): Germin-like protein gene family of a moss, *Physcomitrella patens*, phylogenetically falls into two characteristic new clades. – *Plant Molecular Biology* 56: 381–395.
- Naszradi T., Badacsonyi A., Németh N., Tuba Z. & Batić F. (2004): Zinc, lead and cadmium content in meadow plants and mosses along the M3 Motorway (Hungary). – *Journal of Atmospheric Chemistry* 49: 593–603.
- Natcheva R. & Cronberg N. (2004): What do we know about hybridization among bryophytes in nature? – *Canadian Journal of Botany – Revue Canadienne de Botanique* 82: 1687–1704.
- Natcheva R. & Ganeva A. (2005): Check-list of the bryophytes of Bulgaria. II. Musci. – *Cryptogamie: Bryologie* 26: 209–232.
- Nebel M. & Philippi G. (eds.) (2005): Die Moose Baden-Württembergs 3. *Bryophyta: Sphagnopsida, Marchantiophyta, Anthocerotophyta*. – Verlag Eugen Ulmer: Stuttgart, Germany. [487 pp.]
- Nelson C. R. & Halpern C. B. (2005): Short-term effects of timber harvest and forest edges on ground-layer mosses and liverworts. – *Canadian Journal of Botany – Revue Canadienne de Botanique* 83: 610–620.
- Newmaster S. G., Belland R. J., Arsenault A., Vitt D. H. & Stephens T. R. (2005): The ones we left behind: Comparing plot sampling and floristic habitat sampling for estimating bryophyte diversity. – *Diversity and Distributions* 11: 57–72.
- Newton A. E. & De Luna E. (2004): *Climacium dendroides* from Cofre de Perote, a high-elevation tropical montane site in Veracruz, Mexico. – *Bryologist* 107: 368–372.
- Nifontova M. G. (2005): Use of lichens and mosses for on-line testing of environmental radioactive contamination. – *Russian Journal of Nondestructive Testing* 41: 60–63.
- Niklas K. J. (2004): The cell walls that bind the tree of life. – *Bioscience* 54: 831–841.
- Nikodemus O., Brūmelis G., Tabors G., Lapia L. & Pope S. (2004): Monitoring of air pollution in Latvia between 1990 and 2000 using moss. – *Journal of Atmospheric Chemistry* 49: 521–531.
- Nilsson M. C. & Wardle D. A. (2005): Understory vegetation as a forest ecosystem driver: evidence from the northern Swedish boreal forest. – *Frontiers in Ecology and the Environment* 3: 421–428.
- Nishiyama T., Wolf P. G., Kugita M., Sinclair R. B., Sugita M., Sugiura C., Wakasugi T., Yamada K., Yoshinaga K., Yamaguchi K., Ueda K. & Hasebe M. (2004): Chloroplast phylogeny indicates that bryophytes are monophyletic. – *Molecular Biology and Evolution* 21: 1813–1819.
- Nordin A., Strengbom J., Witzell J., Näsholm T. & Ericson L (2005): Nitrogen deposition and the biodiversity of boreal forests: Implications for the nitrogen critical load. – *Ambio* 34: 20–24.
- Núñez-Olivera E., Arróniz-Crespo M., Martínez-Abaigar J., Tomás R. & Beaucourt N. (2005): Assessing the UV-B tolerance of sun and shade samples of two aquatic bryophytes using short-term tests. – *Bryologist* 108: 435–448.
- Núñez-Olivera E., Martínez-Abaigar J., Tomás R., Beaucourt N. & Arróniz-Crespo M. (2004): Influence of temperature on the effects of artificially enhanced UV-B radiation on aquatic bryophytes under laboratory conditions. – *Photosynthetica* 42: 201–212.
- O'Shea B. J. (2004): The mosses described by Carl Miller from Georg Schweinfurth's collections in Sudan and DR Congo in 1869–1871, with a new checklist of Sudan mosses. – *Cryptogamie: Bryologie* 25: 107–116.
- Ochyra R. (2004): *Tortula hoppeana*, the correct name for *Desmatodon latifolius* in *Tortula* (*Bryopsida, Pottiaceae*). – *Bryologist* 107: 497–500.
- Ochyra R. & Bednarek-Ochyra H. (2004): *Streptocolea atrata* (*Bryopsida, Grimmiaceae*), newly found in western North America, with a review of its global distribution. – *Bryologist* 107: 542–549.
- Ochyra R. & Smith V. R. (2004): On the occurrence of *Ceratodon purpureus* (Hedw.) Brid. (*Ditrichaceae*) on Subantarctic Marion Island. – *Cryptogamie: Bryologie* 25: 131–135.
- Ochyra R. & Zijlstra G. (2005): The basionym of *Eucladium verticillatum* (*Pottiaceae*). – *Taxon* 54: 808–810.
- Oesau A. & Augustin B. (2004): Side effects of herbicides and fungicides on the diversity of mosses in orchards and vineyards. – *Zeitschrift für Pflanzenkrankheiten Und Pflanzenschutz – Journal of Plant Diseases and Protection*, Sp. Iss. 19: 935–942.
- Offner K. (2004): Neufunde von *Andreaea heinemannii* in der bayerischen Rhön. – *Herzogia* 17: 329–331.

- Økland R. H. & Bakkestuen V. (2004): Fine-scale spatial patterns in populations of the clonal moss *Hylocomium splendens* partly reflect structuring processes in the boreal forest floor. – *Oikos* 106: 565–575.
- Økland T., Bakkestuen V., Økland R. H. & Eilertsen O. (2004): Changes in forest understorey vegetation in Norway related to long-term soil acidification and climatic change. – *Journal of Vegetation Science* 15: 437–448.
- Oliván G. & Newton A. M. (2005): Lectotypification of the name *Hypnum ochraceum* (*Bryopsida: Amblystegiaceae*). – *Journal of Bryology* 27: 303–309.
- Oliver M. J. (2004): Desiccation tolerance in bryophytes; Is tolerance the primitive condition in plants? – *Integrative and Comparative Biology* 44: 615–615.
- Oliver M. J., Dowd S. E., Zaragoza J., Mauget S. A. & Payton P. R. (2004): The rehydration transcriptome of the desiccation-tolerant bryophyte *Tortula ruralis*: transcript classification and analysis. – *BMC Genomics* 5: 89.
- Opelt K. & Berg G. (2004): Diversity and antagonistic potential of bacteria associated with bryophytes from nutrient-poor habitats of the Baltic Sea coast. – *Applied and Environmental Microbiology* 70: 6569–6579.
- Orrego O. (2005): Ecological indexes and mathematical models as tools for evaluating the epiphytic bryophyte diversity in Colombian forest patches in the Departamento del Quindío. – *Journal of the Hattori Botanical Laboratory* 97: 271–280.
- Otnyukova T. N., Ignatova E. A., Ignatov M. S. & Fedosov V. E. (2004): New records of *Tortella alpicola* Dix. in Eurasia. – *Arctoa* 13: 197–201.
- Otnyukova T. N. & Ochyra R. (2003): On *Dicranum dispersum* Engelmark and *Dicranum orientale* Otnyukova. – *Arctoa* 12: 115–116.
- Paciolla C. & Tommasi F. (2003): The ascorbate system in two bryophytes: *Brachythecium velutinum* and *Marchantia polymorpha*. – *Biologia Plantarum* 47: 387–393.
- Pannowitz S., Alan Green T. G., Maysek K., Schlensog M., Seppelt R., Sancho L. G., Türk R. & Schroeter B. (2005): Photosynthetic responses of three common mosses from continental Antarctica. – *Antarctic Science* 17: 341–352.
- Parolly G. & Kürschner H. (2004): Ecosociological studies in Ecuadorian bryophyte communities. II. Syntaxonomy of the submontane and montane epiphytic vegetation of S Ecuador. – *Nova Hedwigia* 79: 377–424.
- Parolly G. & Kürschner H. (2005): Ecosociological studies in Ecuadorian bryophyte communities. V. Syntaxonomy, life forms and life strategies of the bryophyte vegetation on decaying wood and tree bases in S Ecuador. – *Nova Hedwigia* 81: 1–35.
- Parolly G., Kürschner H., Schäfer-Verwimp A. & Gradstein S. R. (2004): Cryptogams of the Reserva Biológica San Francisco (Province Zamora-Chinchipe, Southern Ecuador) - III. Bryophytes - Additions and new species. – *Cryptogamie: Bryologie* 25: 271–289.
- Pärtel M., Helm A., Ingerpuu N., Reier Ü. & Tuvi E.-L. (2004): Conservation of Northern European plant diversity: the correspondence with soil pH. – *Biological Conservation* 120: 525–531.
- Paulissen M. P. C. P., Besalú L. E., de Bruijn H., van der Ven P. J. M. & Bobbink R. (2005): Contrasting effects of ammonium enrichment on fen bryophytes. – *Journal of Bryology* 27: 109–117.
- Paulissen M. P. C. P., van der Ven P. J. M., Dees A. J. & Bobbink R. (2004): Differential effects of nitrate and ammonium on three fen bryophyte species in relation to pollutant nitrogen input. – *New Phytologist* 164: 451–458.
- Pedersen N. & Hedenäs L. (2005): Taxonomic and nomenclatural implications of phylogenetic studies of the *Bryaceae* based on molecular data and morphology. – *Bryologist* 108: 123–128.
- Pedersen N. (2005): Validation of *Imbribryum* (*Bryaceae*). – *Bryologist* 108: 449.
- Peng-cheng Wu, Crosby M. R. & He Si (2005): Moss flora of China, English version, Vol. 8: *Sematophyllaceae – Polytrichaceae*. – Missouri Botanical Garden Press, St. Louis, U. S. A. [385 pp.]
- Pénzes-Kónya E. (2004): Comparative analysis of the bryophyte vegetation in acidophilous forest communities in the Bükk Mts (NE Hungary). – *Acta Botanica Hungarica* 46: 373–384.
- Perold S. M. (2003): *Anthocerotophyta*. – In: Germishuizen G. & Meyer N. L. (eds.), *Plants of southern Africa: an annotated checklist*, *Strelitzia* 14: 63–64.
- Perold S. M. (2003): *Hepatophyta*. – In: Germishuizen G. & Meyer N. L. (eds.), *Plants of southern Africa: an annotated checklist*, *Strelitzia* 14: 39–61.

- Peterson R. L. & Massicotte H. B. (2004): Exploring structural definitions of mycorrhizas, with emphasis on nutrient-exchange interfaces. – Canadian Journal of Botany – Revue Canadienne de Botanique 82: 1074–1088.
- Pfeiffer T., Zippel E., Fritz S. & Stech M. (2005): Application of the nonradioactive biotin-streptavidin system to visualize AFLP fragments. – Molecular Ecology Notes 5: 673–675.
- Pharo E. J., Kirkpatrick J. B., Gilfedder L., Mendel L. & Turner P. A. M. (2005): Predicting bryophyte diversity in grassland and eucalypt-dominated remnants in subhumid Tasmania. – Journal of Biogeography 32: 2015–2024.
- Pistarino A., Miserere L., Schumacker R., D'Andrea S. & Soldán Z. (2005): Brionite del Piemonte: la collezione della Val Sangone (Alpi occidentali, Torino). – Catalogo XV, Museo Regionale di Scienze Naturali, Torino. [458 pp.]
- Poikolainen J., Kubin E., Piispanen J. & Karhu J. (2004): Estimation of the long-range transport of mercury, cadmium, and lead to Northern Finland on the basis of moss surveys. – Arctic, Antarctic and Alpine Research 36: 292–297.
- Popper Z. A., Sadler I. H. & Fry S. C. (2004): 3-O-Methylrhamnose in lower land plant primary cell walls. – Biochemical Systematics and Ecology 32: 279–289.
- Potemkin A. D. (2004): Hepatic from North Karelia Biosphere Reserve and adjacent territories: rare and new species for Finland and Karelia borealis. – Memoranda Societatis pro Fauna et Flora Fennica 80: 69–73.
- Pressel S., Matcham H. W. & Duckett J. G. (2005): Studies of protonemal morphogenesis in mosses. X. *Ephemeraceae* revisited; new dimensions underground. – Journal of Bryology 27: 311–318.
- Proctor M. C. F. (2004): Light and desiccation responses of *Weymouthia mollis* and *W. cochlearifolia*, two pendulous rainforest epiphytes from Australia and New Zealand. – Journal of Bryology 26: 167–173.
- Proctor M. C. F. (2005): Why do *Polytrichaceae* have lamellae? – Journal of Bryology 27: 221–230.
- Przywara L., Kuta E. & Kłos J. (2003): Karyology of *Plagiomnium*. II. *Plagiomnium undulatum* (Hedw.) T. Kop. – Acta Biologica Cracoviensia, Ser. Botanica 45/2: 105–110.
- Puche F. (2004): *Pottiaceae: Tortella*. – In: Guerra J. & Cros R. M. (eds.), Flora Briofítica Ibérica, p. 5–15, Sociedad Española de Briología, Murcia.
- Qiu G.-L., Feng X.-B., Wang S.-F. & Shang L.-H. (2005): Mercury and methylmercury in riparian soil, sediments, mine-waste calcines, and moss from abandoned Hg mines in east Guizhou province, southwestern China. – Applied Geochemistry 20: 627–638.
- Quandt D. & Stech M. (2005): Molecular evolution of the *trnL<sub>UAA</sub>* intron in bryophytes. – Molecular Phylogenetics and Evolution 36: 429–443.
- Raghoebarsing A. A., Smolders A. J. P., Schmid M. C., Rijpstra W. I. C., Wolters-Arts M., Derkken J., Jetten M. S. M., Schouten S., Sinnighe Damsté J. S., Lamers L. P. M., Roelofs J. G. M., Op den Camp H. J. & Strous M. (2005): Methanotrophic symbionts provide carbon for photosynthesis in peat bogs. – Nature 436: 1153–1156.
- Rams S., Ros R. M., Werner O. & Shaw A. J. (2004): *Pohlia bolanderi* from Sierra Nevada, Spain, new to the European bryophyte flora. – Bryologist 107: 312–315.
- Rascio N. & La Rocca N. (2005): Resurrection plants: The puzzle of surviving extreme vegetative desiccation. – Critical Reviews in Plant Sciences 24: 209–225.
- Rausch de Traubenberg C. & Ah-Peng C. (2004): A procedure to purify and culture a clonal strain of the aquatic moss *Fontinalis antipyretica* for use as a bioindicator of heavy metals. – Archives of Environmental Contamination and Toxicology 46: 289–295.
- Reimann M. & Dürhammer O. (2004): *Gymnomitrion coralliooides* Nees (*Bryophyta, Hepaticopsida*) neu für Deutschland. – Hoppea, Denkschriften der Regensburgischen Botanischen Gesellschaft 65: 705–710.
- Renner M. A. M. & Braggins J. E. (2004): The sterile gametophyte as a source of informative characters in the genus *Radula* (*Radulaceae: Hepaticae*). – Nova Hedwigia 78: 243–268.
- Renner M. A. M. & Braggins J. E. (2005): Systematically relevant characters of the *Radula* sporophyte (*Radulaceae: Hepaticae*). – Nova Hedwigia 81: 271–300.
- Rensing S. A., Fritzowsky D., Lang D. & Reski R. (2005): Protein encoding genes in an ancient plant: analysis of codon usage, retained genes and splice sites in a moss, *Physcomitrella patens*. – BMC Genomics 6: 43.
- Rensing S. A., Kiessling J., Reski R. & Decker E. L. (2004): Diversification of *ftsZ* during early land plant evolution. – Journal of Molecular Evolution 58: 154–162.
- Rice S. K., Gutman C. & Krouglicof N. (2005): Laser scanning reveals bryophyte canopy structure. – New Phytologist 166: 695–704.

- Richard O., Paquet N., Haudecoeur E. & Charrier B. (2005): Organization and expression of the GSK3/Shaggy kinase gene family in the moss *Physcomitrella patens* suggest early gene multiplication in land plants and an ancestral response to osmotic stress. – Journal of Molecular Evolution 61: 99–113.
- Rico R. & Pócs T. (2004): Bryophytes from high lands of Venezuelan Guyana: Hepatics of Roraima-tepui I. – Cryptogamie: Bryologie 25: 249–269.
- Riedel S. M., Epstein H. E., Walker D. A., Richardson D. L., Calef M. P., Edwards E. & Moody A. (2005): Spatial and temporal heterogeneity of vegetation properties among four tundra plant communities at Ivotuk, Alaska, USA. – Arctic, Antarctic and Alpine Research 37: 25–33.
- Roberts A. W. & Roberts E. (2004): Cellulose synthase (*CesA*) genes in algae and seedless plants. – Cellulose 11: 419–435.
- Roberts N. R., Dalton P. J. & Jordan G. J. (2005): Epiphytic ferns and bryophytes of Tasmanian tree-ferns: A comparison of diversity and composition between two host species. – Austral Ecology 30: 146–154.
- Robinson S. A., Turnbull J. D. & Lovelock C. E. (2005): Impact of changes in natural ultraviolet radiation on pigment composition, physiological and morphological characteristics of the Antarctic moss, *Grimmia antarctica*. – Global Change Biology 11: 476–489.
- Robinson S. A., Wasley J. & Tobin A. K. (2003): Living on the edge - plants and global change in continental and maritime Antarctica. – Global Change Biology 9: 1681–1717.
- Ron E., Gómez D. & Fernández-Mendoza F. (2004): *Pottiaceae: Pleurochaete*. – In: Guerra J. & Cros R. M. (eds.), Flora Briofítica Ibérica, p. 15–17, Sociedad Española de Briología, Murcia.
- Ros R. M. & Werner O. (2005): *Pottiaceae: Stegonia, Pottia, Microbryum*. – In: Guerra J. & Cros R. M. (eds.), Flora Briofítica Ibérica, p. 5–35, Sociedad Española de Briología, Murcia.
- Ros R. M., Werner O. & Rams S. (2005): New taxonomical data on the genus *Microbryum* Schimp. (*Pottiaceae, Musci*). – Cryptogamie: Bryologie 26: 167–172.
- Rowntree J. K., Sheffield E. & Burch J. (2005): Growth and development of mosses are inhibited by the common herbicide Asulam. – Bryologist 108: 287–294.
- Rozema J., Boelen P. & Blokker P. (2005): Depletion of stratospheric ozone over the Antarctic and Arctic: Responses of plants of polar terrestrial ecosystems to enhanced UV-B, an overview. – Environmental Pollution 137: 428–442.
- Russell J. & Bulman S. (2005): The liverwort *Marchantia foliacea* forms a specialized symbiosis with arbuscular mycorrhizal fungi in the genus *Glomus*. – New Phytologist 165: 567–579.
- Ryall K., Whitton J., Schofield W., Ellis S. & Shaw A. J. (2005): Molecular phylogenetic study of interspecific variation in the moss *Isothecium* (*Brachytheciaceae*). – Systematic Botany 30: 242–247.
- Rycroft D. S., Groth H. & Heinrichs J. (2004): Reinstatement of *Plagiochila maderensis* (*Jungermanniopsida: Plagiochilaceae*) based on chemical evidence and nrDNA ITS sequences. – Journal of Bryology 26: 37–45.
- Rykovskiy G. F. & Maslovskiy O. M. (2004): Flora Belarusi: *Bryophyta* [Flora of Belarus]. – Technalohija, Minsk. [436 pp.]
- Sabovljević A., Sabovljević M., Grubišić D. & Konjević R. (2005): The effect of sugars on development of two moss species (*Bryum argenteum* and *Atrichum undulatum*) during in vitro culture. – Belgian Journal of Botany 138: 79–84.
- Sætersdal M., Gjerde I. & Blom H. H. (2005): Indicator species and the problem of spatial inconsistency in nestedness patterns. – Biological Conservation 122: 305–316.
- Samecka-Cymerman A., Kolon K. & Kempers A. J. (2005): A comparison of native and transplanted *Fontinalis antipyretica* Hedw. as biomonitor of water polluted with heavy metals. – Science of the Total Environment 341: 97–107.
- Samecka-Cymerman A., Kolon K. & Kempers A. J. (2005): Differences in concentration of heavy metals between native and transplanted *Plagiothecium denticulatum*: A case study of soils contaminated by oil well exudates in South East Poland. – Archives of Environmental Contamination and Toxicology 49: 317–321.
- Samecka-Cymerman A., Kolon K., Kempers A. J., Jansen J. & Boonen B. (2005): Bioaccumulation of elements in bryophytes from Serra da Estrela, Portugal, and Veluwezoom, the Netherlands. – Environmental Science and Pollution Research 12: 71–79.
- Sanderska A., Filipiak E. & Pisarek W. (2003): Bryophytes in the Bolimów Landscape Park. – Monographiae Botanicae 9: 197–231.
- Sari H., Mendil D., Tüzen M., Hasdemir E. & Özdemir T. (2005): AAS determination of trace metals in some moss samples from Trabzon, Turkey. – Fresenius Environmental Bulletin 14: 473–477.

- Sauberer N., Zulka K. P., Abensperg-Traun M., Berg H.-M., Bieringer G., Milasowszky N., Moser D., Plutzar C., Pollheimer M., Storch C., Tröstl R., Zechmeister H. & Grabherr G. (2004): Surrogate taxa for biodiversity in agricultural landscapes of eastern Austria. – Biological Conservation 117: 181–190.
- Sawai Y. & Nasu H. (2005): A 4500-year record of emergence events at Onnetoh, Hokkaido, northern Japan, reconstructed using plant macrofossils. – Marine Geology 217: 49–65.
- Schaumann F. (2005): Terricolous bryophyte vegetation of Chilean temperate rain forests. Communities, adaptive strategies and divergence patterns (Studies in austral temperate rain forest bryophytes 26). – Bryophytorum Bibliotheca 62: 1–154.
- Schaumann F., Frey W., Pfeiffer T. & Stech M. (2005): Molecular circumscription, intrafamilial relationships and biogeography of the Gondwanan liverwort family *Pallaviciniaceae* (*Hepaticophytina*, *Bryophyta*) – Studies in austral temperate rain forest bryophytes 27. – Plant Systematics and Evolution 252: 27–48.
- Schintu M., Cogoni A., Durante L., Cantaluppi C. & Contu A (2005): Moss (*Bryum radiculosum*) as a bioindicator of trace metal deposition around an industrialised area in Sardinia (Italy). – Chemosphere 60: 610–618.
- Schlensog M., Pannewitz S., Green T. G. A. & Schroeter B. (2004): Metabolic recovery of continental Antarctic cryptogams after winter. – Polar Biology 27: 399–408.
- Schmull M., Heinrichs J., Baier R., Ullrich D., Wagenitz G., Groth H., Hourticolon S. & Gradstein S. R. (2005): The type database at Göttingen (GOET) – a virtual herbarium online. – Taxon 54: 251–254.
- Schröder W. & Pesch R. (2004): Integrative monitoring analysis aiming at the detection of spatial and temporal trends of metal accumulation in mosses. – Journal of Atmospheric Chemistry 49: 23–38.
- Schröder W. & Pesch R. (2004): Spatial analysis and indicator building for metal accumulation in mosses. – Environmental Monitoring and Assessment 98: 131–155.
- Schröder W. & Pesch R. (2005): Time series of metals in mosses and their correlation with selected sampling site-specific and ecoregional characteristics in Germany. – Environmental Science and Pollution Research 12: 159–167.
- Schween G., Schulte J., Reski R. & Hohe A. (2005): Effect of ploidy level on growth, differentiation, and morphology in *Physcomitrella patens*. – Bryologist 108: 27–35.
- Sedia E. G. & Ehrenfeld J. G. (2005): Differential effects of lichens, mosses and grasses on respiration and nitrogen mineralization in soils of the New Jersey Pinelands. – Oecologia 144: 137–147.
- Seifert E. (2004): Epiphyten im Wandel: Zur Vorkommen epiphytischer Moose im Erzgebirge. – Beiträge zum Naturschutz im Mittleren Erzgebirgskreis 3: 77–86.
- Selosse M. A. (2005): Are liverworts imitating mycorrhizas? – New Phytologist 165: 345–349.
- Senn H. (2000): Die Moose des Fürstentums Liechtenstein. – Naturkundliche Forschung im Fürstentum Liechtenstein 17: 1–248.
- Sérgio C. & Carvalho S. (2003): Annotated catalogue of Portuguese bryophytes. – Portugaliae Acta Biologica 21: 5–230.
- Sérgio C., Draper D. & Garcia C. (2005): Modelling the distribution of *Cryptothallus mirabilis* Malmb. (*Aneuraceae*, *Hepaticopsida*) in the Iberian Peninsula. – Journal of the Hattori Botanical Laboratory 97: 309–316.
- Sérgio C., Sim-Sim M., Jansen J., Garcia C. & Carvalho P. (2003): Bryophyte diversity and impact of land-use in semi-natural grasslands in the Serra da Estrela (Portugal). – Bocconea 16: 1002–1019.
- Shaw A. J. & Cox C. J. (2005): Variation in “biodiversity value” of peatmoss species in *Sphagnum* section *Acutifolia* (*Sphagnaceae*). – American Journal of Botany 92: 1774–1783.
- Shaw A. J., Cox C. J. & Boles S. B. (2004): Phylogenetic relationships among *Sphagnum* sections: *Hemitheca*, *Isocladus*, and *Subsecunda*. – Bryologist 107: 189–196.
- Shaw A. J., Cox C. J. & Goffinet B. (2005): Global patterns of moss diversity: taxonomic and molecular inferences. – Taxon 54: 337–352.
- Shaw A. J., Melosik I., Cox C. J. & Boles S. B. (2005): Divergent and reticulate evolution in closely related species of *Sphagnum* section *Subsecunda*. – Bryologist 108: 363–376.
- Shaw J. & Renzaglia K. (2004): Phylogeny and diversification of bryophytes. – American Journal of Botany 91: 1557–1581.
- Shevock J. R. & Shaw A. J. (2005): *Pohlia robertsonii* and *P. rabunbaldensis* (*Bryopsida*, *Mniaceae*), two new species from the western and eastern United States. – Bryologist 108: 177–182.
- Shimamura M., Brown R. C., Lemmon B. E., Akashi T., Mizuno K., Nishihara N., Tomizawa K. I., Yoshimoto K., Deguchi H., Hosoya H., Horio T. & Mineyuki Y. (2004):  $\gamma$ -tubulin in basal land plants:

- characterization, localization, and implication in the evolution of acentriolar microtubule organizing centers. – *Plant Cell* 16: 45–59.
- Shimamura M., Furuki T. & Deguchi H. (2005): Sporophyte anatomy of *Cavicularia densa* (*Blasiaceae*). – *Bryologist* 108: 420–426.
- Sim-Sim M., Carvalho P., Sérgio C., Garcia C. & Rego F. (2004): Recolonisation and changes in bryophyte and lichen biodiversity in burned areas from the Serra da Estrela (Portugal). – *Cryptogamie: Bryologie* 25: 367–384.
- Sim-Sim M., Esquível M. da G., Fontinha S. & Stech M. (2005): The genus *Plagiochila* (Dumort.) Dumort. (*Plagiochilaceae*, *Hepaticophytina*) in Madeira Archipelago – molecular relationships, ecology, and biogeographic affinities. – *Nova Hedwigia* 81: 449–462.
- Sim-Sim M., Esquível M. G., Fontinha S. & Carvalho S. (2004): *Plagiochila stricta* Lindenb. new to Madeira Morphological and molecular evidence. – *Nova Hedwigia* 79: 497–505.
- Sim-Sim M., Stech M., Esquivel M. G., Figueiredo A. C., Costa M. M., Barroso J. G., Pedro L. G., Fontinha S. & Lobo C. (2005): *Plagiochila spinulosa* (Dicks.) Dumort. (*Plagiochilaceae*, *Hepaticophytina*) in Madeira Island – morphological, phytochemical, and molecular evidence. – *Journal of the Hattori Botanical Laboratory* 98: 131–147.
- Singh M. K., Juhasz A., Csintalan Z., Kaligaric M., Marek M. V., Urban O. & Tuba Z. (2005): Preliminary estimation of bryophyte biomass and carbon pool from three contrasting different vegetation types. – *Cereal Research Communications* 33: 267–270.
- Skotnicki M. L., Mackenzie A. M., Clements M. A. & Selkirk P. M. (2005): DNA sequencing and genetic diversity of the 18S-26S nuclear ribosomal internal transcribed spacers (ITS) in nine Antarctic moss species. – *Antarctic Science* 17: 377–384.
- Skrzypczak R. & Pierrot R. B. (2004): *Desmatodon wilczekii* Meyl. retrouvé en Savoie. – *Bulletin de la Société Botanique du Centre-Ouest, Nouvelle Série* 35: 359–362.
- Skrzypczak R. (2004): Contribution à la bryoflore de Savoie. – *Bulletin de la Société Botanique du Centre-Ouest, Nouvelle Série* 35: 297–336.
- Skrzypczak R. (2004): Presence de *Didymodon glaucus* Ryan en France. – *Bulletin de la Société Botanique du Centre-Ouest, Nouvelle Série* 35: 353–357.
- Slavik K., Peterson B. J., Deegan L. A., Bowden W. B., Hershey A. E. & Hobbie J. E. (2004): Long-term responses of the Kuparuk River ecosystem to phosphorus fertilization. – *Ecology* 85: 939–954.
- Sliwinski M. K. & Goodman R. M. (2004): Comparison of crenarchaeal consortia inhabiting the rhizosphere of diverse terrestrial plants with those in bulk soil in native environments. – *Applied and Environmental Microbiology* 70: 1821–1826.
- Sloover J. L. (2003): Illustrations de mousses Africaines. – *Scripta Botanica Belgica* 28: 1–223.
- Smagin V. A. (2004): Travyanaya i travyanomokhovaya rastitel'nost' evtrofnykh bolot taezhnoj zony evropejskoj Rossii i ee sintaksonomija [Rich fen vegetation of the taiga zone of the European Russia and its syntaxonomy]. – *Botanicheskiy Zhurnal* 89: 1302–1319.
- Smaw S. G. & Petersen R. L. (2005): Effects of simulated rain and cement amendments on protonematal growth of the urban moss, *Bryum capillare*, in culture. – *Bryologist* 108: 236–240.
- Smirnov L. I., Frontas'eva M. V., Steinnes E., Lyapunov S. M., Cherchintsev V. D., Romanov S. A. & Samosadnyi V. T. (2004): Multidimensional statistical analysis of the concentration of heavy metals and radionuclides in moss and soil in southern Urals. – *Atomic Energy* 97: 510–515.
- Snäll T., Hagström A., Rudolphi J. & Rydin H. (2004): Distribution pattern of the epiphyte *Neckera pennata* on three spatial scales - importance of past landscape structure, connectivity and local conditions. – *Ecography* 27: 757–766.
- So M. L. (2004): *Metzgeria* (*Metzgeriaceae*, *Marchantiophyta*) in Africa. – *New Zealand Journal of Botany* 42: 271–292.
- So M. L. (2005): A synopsis of *Radula* (*Radulaceae*, *Marchantiophyta*) in New Zealand and Tasmania. – *Journal of the Hattori Botanical Laboratory* 98: 149–174.
- Söderström L. & During H. J. (2005): Bryophyte rarity viewed from the perspectives of life history strategy and metapopulation dynamics. – *Journal of Bryology* 27: 261–268.
- Solga A., Burkhardt J., Zechmeister H. G. & Frahm J.-P. (2005): Nitrogen content, <sup>15</sup>N natural abundance and biomass of two pleurocarpous mosses *Pleurozium schreberi* (Brid.) Mitt. and *Scleropodium purum* (Hedw.) Limpr. in relation to atmospheric nitrogen deposition. – *Environmental Pollution* 134: 465–473.
- Soria A., Gómez D., Ron E. & Guerra J. (2005): *Pottiaceae: Timmiella*. – In: Guerra J. & Cros R. M. (eds.), *Flora Briofítica Ibérica*, p. 5–8, Sociedad Española de Briología, Murcia.

- Sotiaux A. & Sotiaux O. (2005): *Tortula mucronifolia* (Pottiaceae, Musci), espèce boréo-montagnarde nouvelle pour la bryoflore belge dans la région de Soignies (province du Hainaut). – Dumortiera 84: 15–17.
- Sotiaux A. & Vanderpoorten A. (2004): Catalogue atlas commenté et mesures de conservation des bryophytes du bassin hydrographique de la Semois (Belgique, France). – Lejeunia, Nouvelle série 175: 1–116.
- Spence J. R. (2005): New genera and combinations in *Bryaceae* (*Bryales, Musci*) for North America. – Phytologia 87: 15–28.
- Springer S. (2004): Die Vegetation des Thalhamer Mooses. – Berichte der Bayerischen Botanischen Gesellschaft zur Erforschung der Heimischen Flora 73/74: 157–169.
- Staniaszek M. (2005): Epixyloous mosses and liverworts in Belgium: a list based on literature reports. – Nowellia Bryologica 28: 2–17.
- Staples G. W., Imada C. T., Hoe W. J. & Smith C. W. (2004): A revised checklist of Hawaiian mosses. – Tropical Bryology 25: 35–70.
- Stark L. R. (2005): Phenology of patch hydration, patch temperature and sexual reproductive output over a four-year period in the desert moss *Crossidium crassinerve*. – Journal of Bryology 27: 231–240.
- Stark L. R., McLetchie D. N., & Mishler B. D. (2005): Sex expression, plant size, and spatial segregation of the sexes across a stress gradient in the desert moss *Syntrichia caninervis*. – Bryologist 108: 183–193.
- Stark L. R., Nichols L., McLetchie D. N. & Bonine M. L. (2005): Do the sexes of the desert moss *Syntrichia caninervis* differ in desiccation tolerance? A leaf regeneration assay. – International Journal of Plant Sciences 166: 21–29.
- Stebel A. & Fojcik B. (2003): Atlas rozmieszczenia mchów chronionych Polski w województwie Śląskim [Distribution atlas of Polish legally protected mosses in the Silesia province]. – In: Materiały Opracowania, Tom 7, p. 1–112, Centrum Dziedzictwa Przyrody Górnego Śląska, Katowice.
- Stebel A. & Ochyra R. (eds.) (2004): Bryological studies in Western Carpathians. – Sorus, Poznań [160 pp.]
- Stebel A., Ochyra R., Stuchlik L. & Parusel J. B. (2004): Mosses of the Polica range (Polish Western Carpathians). – Sorus, Poznań [121 pp.]
- Stotler R. E. & Crandall-Stotler B. (2005): A revised classification of the *Anthocerotophyta* and a checklist of the hornworts of North America, north of Mexico. – Bryologist 108: 16–26.
- Streiff A. (2005): Morphological study of the genus *Grimmia* Hedw. (*Grimmiaceae, Bryopsida*). – Journal of the Hattori Botanical Laboratory 97: 317–338.
- Suchara I. & Sucharová J. (2004): Current atmospheric deposition loads and their trends in the Czech Republic determined by mapping the distribution of moss element contents. – Journal of Atmospheric Chemistry 49: 503–519.
- Suetsugu N., Mittmann F., Wagner G., Hughes J. & Wada M. (2005): A chimeric photoreceptor gene, NEOCHROME, has arisen twice during plant evolution. – Proceedings of the National Academy of Sciences of the United States of America 102: 13705–13709.
- Sugita M., Sugiura C., Arikawa T. & Higuchi M. (2004): Molecular evidence of a *rpoA* gene in the basal moss chloroplast genomes: *rpoA* is a useful molecular marker for phylogenetic analysis of mosses. – Hikobia 14: 171–175.
- Sugiura C. & Sugita M. (2004): Plastid transformation reveals that moss tRNA<sup>Arg</sup>-CCG is not essential for plastid function. – Plant Journal 40: 314–321.
- Sundberg S. (2005): Larger capsules enhance short-range spore dispersal in *Sphagnum*, but what happens further away? – Oikos 108: 115–124.
- Szańkowski M. & Kłosowski S. (2004): Distribution and habitat conditions of the phytocoenoses of *Sphagnum denticulatum* Bridel and *Warnstorffia exannulata* (B., S. & G.) Loeske in Polish *Lobelia* lakes. – Acta Societatis Botanicorum Poloniae 73: 255–262.
- Szwejkowski J., Buczkowska K. & Odrzykoski I. J. (2005): *Conocephalum salebrosum* (*Marchantiopsida, Conocephalaceae*) – a new Holarctic liverwort species. – Plant Systematics and Evolution 253: 133–158.
- Tabors G., Brūmelis G., Lapina L., Pospelova G. & Nikodemus O. (2004): Changes in element concentrations in moss segments after cross-transplanting between a polluted and non-polluted site. – Journal of Atmospheric Chemistry 49: 191–197.
- Tahvanainen T. & Tuomaala T. (2003): The reliability of mire water pH measurements – a standard sampling protocol and implications to ecological theory. – Wetlands 23: 701–708.
- Takezawa D. & Minami A. (2004): Calmodulin-binding proteins in bryophytes: identification of abscisic acid-, cold-, and osmotic stress-induced genes encoding novel membrane-bound transporter-like proteins. – Biochemical and Biophysical Research Communications 317: 428–436.

- ter Braak C. J. F. & Schaffers A. P. (2004): Co-correspondence analysis: A new ordination method to relate two community compositions. – *Ecology* 85: 834–846.
- Thiel H. & Preussing M. (2004): *Dicranum viride* (Sull. & Lesq.) Lindb. in Thüringen – Lebensraum, Vergesellschaftung, Verbreitung, Bestandsentwicklung, Schutz. – *Haussknechtia* 10: 69–102.
- Thompson D. B., Walker L. R., Landau F. H. & Stark L. R. (2005): The influence of elevation, shrub species, and biological soil crust on fertile islands in the Mojave Desert, USA. – *Journal of Arid Environments* 61: 609–629.
- Thormann M. N., Bayley S. E. & Currah R. S. (2004): Microcosm tests of the effects of temperature and microbial species number on the decomposition of *Carex aquatilis* and *Sphagnum fuscum* litter from southern boreal peatlands. – *Canadian Journal of Microbiology* 50: 793–802.
- Thornton L. E., Keren N., Ohad I. & Pakrasi H. B. (2005): *Physcomitrella patens* and *Ceratodon purpureus*, mosses as model organisms in photosynthesis studies. – *Photosynthesis Research* 83: 87–96.
- Tomassen H. B. M., Smolders A. J. P., Lamers L. P. M. & Roelofs J. G. M. (2005): How bird droppings can affect the vegetation composition of ombrotrophic bogs. – *Canadian Journal of Botany – Revue Canadienne de Botanique* 83: 1046–1056.
- Tømmervik H., Johansen B., Tombre I., Thannheiser D., Høgda K. A., Gaare E. & Wielgolaski F. E. (2004): Vegetation changes in the Nordic mountain birch forest: The influence of grazing and climate change. – *Arctic, Antarctic and Alpine Research* 36: 323–332.
- Townsend C. C. (2005): Mosses from the Caucasian region and eastern Turkey. – *Journal of Bryology* 27: 143–152.
- Tremper A. H., Burton M. A. S. & Higgs D. E. B. (2004): Field and laboratory exposures of two moss species to low level metal pollution. – *Journal of Atmospheric Chemistry* 49: 111–120.
- Tsikritzis L. I. (2005): Chemometrics of the distribution and origin of  $^{226}\text{Ra}$ ,  $^{228}\text{Ra}$ ,  $^{40}\text{K}$  and  $^{137}\text{Cs}$  in plants near the West Macedonia Lignite Center (Greece). – *Journal of Radioanalytical and Nuclear Chemistry* 264: 651–656.
- Tsubota H., De Luna E., González E., Ignatov M. S. & Deguchi H. (2004): Molecular phylogenetics and ordinal relationships based on analyses of a large-scale data set of 600 *rbcL* sequences of mosses. – *Hikobia* 14: 149–169.
- Turetsky M. R. (2004): Decomposition and organic matter quality in continental peatlands: The ghost of permafrost past. – *Ecosystems* 7: 740–750.
- Turner P. A. M. & Pharo E. J. (2005): Influence of substrate type and forest age on bryophyte species distribution in Tasmanian mixed forest. – *Bryologist* 108: 67–85.
- Tyler T. (2005): The Bryophyte flora of Scanian sand-steppe vegetation and its relation to soil pH and phosphate availability. – *Lindbergia* 30: 11–20.
- Ugur A., Özden B., Saç M. M., Yener G., Altınbaş Ü., Kurucu Y. & Bolca M. (2004): Lichens and mosses for correlation between trace elements and  $^{210}\text{Po}$  in the areas near coal-fired power plant at Yatağan, Turkey. – *Journal of Radioanalytical and Nuclear Chemistry* 259: 87–92.
- Uotila A., Hotanen J.-P. & Kouki J. (2005): Succession of understory vegetation in managed and seminatural Scots pine forests in eastern Finland and Russian Karelia. – *Canadian Journal of Forest Research – Revue Canadienne de Recherche Forestière* 35: 1422–1441.
- Uotila A. & Kouki J. (2005): Understorey vegetation in spruce-dominated forests in eastern Finland and Russian Karelia: Successional patterns after anthropogenic and natural disturbances. – *Forest Ecology and Management* 215: 113–137.
- Urban K. E. (2005): Plant species dynamics during restoration of heath ponds in northwestern Germany. – *Phytocoenologia* 35: 511–532.
- van der Wal R., Bardgett R. D., Harrison K. A. & Stien A. (2004): Vertebrate herbivores and ecosystem control: cascading effects of faeces on tundra ecosystems. – *Ecography* 27: 242–252.
- van der Wal R., Pearce I. S. K. & Brooker R. W. (2005): Mosses and the struggle for light in a nitrogen-polluted world. – *Oecologia* 142: 159–168.
- van Rooy J. (2003): *Bryophyta*. – In: Germishuizen G. & Meyer N. L. (eds.), *Plants of southern Africa: an annotated checklist*, *Strelitzia* 14: 1–37.
- van Tooren B. & Bruin K. (2004): Bryophytes of the sand dunes of the West Frisian Islands. – *Lindbergia* 29: 51–63.
- van Wijk M. T., Clemmensen K. E., Shaver G. R., Williams M., Callaghan T. V., Chapin F. S., Cornelissen J. H. C., Gough L., Hobbie S. E., Jonasson S., Lee J. A., Michelsen A., Press M. C., Richardson S. J. & Rueth H. (2004): Long-term ecosystem level experiments at Toolik Lake, Alaska, and at Abisko,

- Northern Sweden: generalizations and differences in ecosystem and plant type responses to global change. – *Global Change Biology* 10: 105–123.
- van Zanten B. O. (2005): *Philonotis hastata* (Duby) Wijk & Marg., een tropisch mos gevonden in een onverwarmde kas te Eelde (Drenthe). – *Buxbaumiella* 70: 23–27.
- Váňa J. (2004): Bryophyte flora of Uganda. 1. *Jungermanniaceae*. – *Journal of Bryology* 26: 147–151.
- Váňa J. (2004): Bryophyte flora of Uganda. 2. *Gymnomitriaceae*. – *Journal of Bryology* 26: 199–202.
- Váňa J. (2004): Bryophyte flora of Uganda. 3. *Lophoziaceae*. – *Journal of Bryology* 26: 293–299.
- Váňa J. & Gremmen N. (2005): Hepatics of Heard Island. – *Cryptogamie: Bryologie* 26: 79–90.
- Váňa J., Piippo S. & Koponen T. (2005): Bryophyte flora of Hunan Province, China. 6. *Jungermanniaceae* and *Gymnomitriaceae (Hepaticae)*. – *Acta Botanica Fennica* 178: 57–78.
- Vanderpoorten A., Ignatov M. S., Huttunen S. & Goffinet B. (2005): A molecular and morphological recircumscription of *Brachytheciastrum* (*Brachytheciaceae, Bryopsida*). – *Taxon* 54: 369–376.
- Vanderpoorten A., Sotiaux A. & Engels P. (2005): A GIS-based survey for the conservation of bryophytes at the landscape scale. – *Biological Conservation* 121: 189–194.
- Vasilevich V. I. & Bibikova T. V. (2004): Sfagnovye el'niki evropeyskoy Rossii [Sphagnum-spruce forests in European Russia]. – *Botanicheskiy Zhurnal* 89: 734–748.
- Veinotte C., Freedman B., Maass W. & Kirstein F. (2003): Comparison of the ground vegetation in spruce plantations and natural forest in the Greater Fundy Ecosystem, New Brunswick. – *Canadian Field-Naturalist* 117: 531–540.
- Vellak K. & Ingerpuu N. (2005): Management effects on bryophytes in Estonian forests. – *Biodiversity and Conservation* 14: 3255–3263.
- Virchenko V. M. (2004): Novi znakhidky ridkisnykh dlya Ukrayiny mokhopodibnykh [New findings of rare bryophytes for Ukraine]. – *Ukrain's'kyi Botanichnyi Zhurnal* 61: 106–110.
- Votintseva A. A. & Mukhin V. A. (2004): Effect of extractive compounds from lichens and mosses on the development of basidiospores and mycelium of tinder fungus. – *Russian Journal of Ecology* 35: 283–289.
- Wahren C.-H. A., Walker M. D. & Bret-Harte M. S. (2005): Vegetation responses in Alaskan arctic tundra after 8 years of a summer warming and winter snow manipulation experiment. – *Global Change Biology* 11: 537–552.
- Wall D. P. (2005): Origin and rapid diversification of a tropical moss. – *Evolution* 59: 1413–1424.
- Wanek W. & Pörtl K. (2005): Phyllosphere nitrogen relations: reciprocal transfer of nitrogen between epiphyllous liverworts and host plants in the understorey of a lowland tropical wet forest in Costa Rica. – *New Phytologist* 166: 577–588.
- Wang Y.-D., Mosbrugger V. & Zhang H. (2005): Early to Middle Jurassic vegetation and climatic events in the Qaidam Basin, northwest China. – *Palaeogeography, Palaeoclimatology, Palaeoecology* 224: 200–216.
- Weibull H. & Rydin H. (2005): Bryophyte species richness on boulders: relationship to area, habitat diversity and canopy tree species. – *Biological Conservation* 122: 71–79.
- Werner O. & Guerra J. (2004): Molecular phylogeography of the moss *Tortula muralis* Hedw. (*Pottiaceae*) based on chloroplast *rps4* gene sequence data. – *Plant Biology* 6: 147–157.
- Werner O., Jiménez J. A. & Ros R. M. (2004): The systematic position of the moss *Kingiobryum paramicola* (*Pottiaceae*) based on molecular and morphological data. – *Bryologist* 107: 215–221.
- Werner O., Jiménez J. A., Ros R. M., Cano M. J. & Guerra J. (2005): Preliminary investigation of the systematics of *Didymodon* (*Pottiaceae, Musci*) based on nrITS sequence data. – *Systematic Botany* 30: 461–470.
- Werner O., Rams S. & Ros R. M. (2005): Genetic diversity of *Pohlia bolanderi* (*Mniaceae*), a rare and threatened moss in Sierra Nevada (Spain), estimated by ISSR molecular markers. – *Nova Hedwigia* 81: 413–420.
- Werner O., Ros R. M. & Grundmann M. (2005): Molecular phylogeny of *Trichostomoidae* (*Pottiaceae, Bryophyta*) based on nrITS sequence data. – *Taxon* 54: 361–368.
- Whitehouse H. E. & Bayley S. E. (2005): Vegetation patterns and biodiversity of peatland plant communities surrounding mid-boreal wetland ponds in Alberta, Canada. – *Canadian Journal of Botany – Revue Canadienne de Botanique* 83: 621–637.
- Wibraham J. & Long D. G. (2005): *Zygodon* Hook. & Taylor and *Bryomaltaea* Goffinet (*Bryopsida: Orthotrichaceae*) in the Sino-Himalaya. – *Journal of Bryology* 27: 329–342.
- Wigginton M.J. (2004): Checklist and distribution of the liverworts and hornworts of sub-Saharan Africa, including the East African Islands (edition 2, September 2004). – *Tropical Bryology Research Reports* 5: 1–102.

- Wiklund K. & Rydin H. (2004): Ecophysiological constraints on spore establishment in bryophytes. – *Functional Ecology* 18: 907–913.
- Wilkinson C. E., Hocking M. D. & Reimchen T. E. (2005): Uptake of salmon-derived nitrogen by mosses and liverworts in coastal British Columbia. – *Oikos* 108: 85–98.
- Wood A. J., Oliver M. J. & Cove D. J. (eds.) (2004): New frontiers in bryology: physiology, molecular biology and functional genomics. – Kluwer Academic Publishers, Dordrecht. [xi + 203 pp.]
- Wood A. J., Reski R. & Frank W. (2004): Isolation and characterization of *ALDH11A5*, a novel non-phosphorylating GAPDH cDNA from *Physcomitrella patens*. – *Bryologist* 107: 385–387.
- Wu Q.-L., Riise G., Pflugmacher S., Greulich K. & Steinberg C. E. W. (2005): Combined effects of the fungicide propiconazole and agricultural runoff sediments on the aquatic bryophyte *Vesicularia dubyana*. – *Environmental Toxicology and Chemistry* 24: 2285–2290.
- Wyatt R., Odrzykoski I. J. & Cronberg N. (2005): High levels of genetic variation in the haploid leafy liverwort *Porella platyphylla* from the southeastern United States. – *Journal of Bryology* 27: 247–252.
- Yang R.-D., Mao J.-R., Zhang W.-H., Jiang L.-J. & Gao H. (2004): Bryophyte-like fossil (*Parafunaria sinensis*) from Early-Middle Cambrian Kaili Formation in Guizhou Province, China. – *Acta Botanica Sinica* 46: 180–185.
- Zackrisson O., DeLuca T. H., Nilsson M. C., Sellstedt A. & Berglund L. M. (2004): Nitrogen fixation increases with successional age in boreal forests. – *Ecology* 85: 3327–3334.
- Zanazzi A. & Mora G. (2005): Paleoclimatic implications of the relationship between oxygen isotope ratios of moss cellulose and source water in wetlands of Lake Superior. – *Chemical Geology* 222: 281–291.
- Zechmeister H. G. (2005): Bryophytes of continental salt meadows in Austria. – *Journal of Bryology* 27: 297–302.
- Zechmeister H. G., Hohenwallner D., Riss A. & Hanus-Illar A. (2005): Estimation of element deposition derived from road traffic sources by using mosses. – *Environmental Pollution* 138: 238–249.
- Zechmeister H. G., Riss A. & Hanus-Illar A. (2004): Biomonitoring of atmospheric heavy metal deposition by mosses in the vicinity of industrial sites. – *Journal of Atmospheric Chemistry* 49: 461–477.
- Zheleznova G. V., Kuznetsova E. G., Evdokimova T. V. & Turubanova L. P. (2005): Monitoring of vegetation development in technogenically disturbed areas of the Usinskoe oil field. – *Russian Journal of Ecology* 36: 243–248.
- Zheleznova G. V. & Shubina T. P. (2005): Vidovoy sostav mkhov tekhnogenno narushennykh landshaftov Respubliki Komi. – *Botanicheskiy Zhurnal* 90: 215–221.
- Zhuravleba E. N. & Ipatov V. S. (2005): Vzaimootnosheniya vidov rasteniy v zabolochennykh sosnovykh lesakh severo-zapada Rossii. 1. Vliyaniye ekologicheskikh faktorov, formiruemnykh drevostoyem, na vidy mokhovogo i travyano-kustarnichkovogo yarusov. – *Botanicheskiy Zhurnal* 90: 702–711.
- Zielke M., Solheim B., Spjelkavik S. & Olsen R. A. (2005): Nitrogen fixation in the high arctic: Role of vegetation and environmental conditions. – *Arctic, Antarctic and Alpine Research* 37: 372–378.
- Zobell O., Coupland G. & Reiss B. (2005): The family of CONSTANS-like genes in *Physcomitrella patens*. – *Plant Biology* 7: 266–275.
- Zoghbi M. G. B., Andrade E. H. A., Lobato R. C. L., Tavares A. C. C., Souza A. P. S., Conceição C. C. C. & Guimarães E. F. (2005): *Peperomia circinnata* Link and *Peperomia rotundifolia* (L.) Kunth growing on different host-trees in Amazon: volatiles and relationship with bryophytes. – *Biochemical Systematics and Ecology* 33: 269–274.