

## NOVÁ BRYOLOGICKÁ LITERATURA XV.

### New bryological literature, XV

Jan Kučera & Táňa Štechová

Jihočeská Univerzita, Biologická fakulta, Branišovská 31, CZ–370 05 České Budějovice, e-mail:  
kucera@bf.jcu.cz

Počínaje tímto dílem Nové bryologické literatury dochází k částečné změně v koncepcí excerptí a formálního uspořádání rubriky. Vzhledem ke stále rostoucímu počtu ve světě vycházejících bryologických prací budeme výběr ze světové bryologické literatury i naší bibliografii přinášet v každém čísle. Dosud vycházející rubrika „Česká a slovenská bryologická bibliografie“ bude transformována do několika podrubrik Nové bryologické literatury a budou v ní publikovány pouze práce vycházející v České nebo Slovenské republice.

#### Výběr ze světové bryologické literatury [Selection from the world bryological literature]

- Aboal J. R., Couto J. A., Fernández J. A. & Carballeira A. (2006): Definition and number of subsamples for using mosses as biomonitoring of airborne trace elements. – Archives of Environmental Contamination and Toxicology 50: 88–96.
- Aboal J. R., Real C., Fernández J. A. & Carballeira A. (2006): Mapping the results of extensive surveys: The case of atmospheric biomonitoring and terrestrial mosses. – Science of the Total Environment 356: 256–274.
- Afonina O. M., Potemkin A. D. & Czernyadjeva I. V. (eds) (2005): Actual Problems in Bryology: Proceedings of the International Meeting Devoted to the 90<sup>th</sup> Anniversary of A. L. Abramova (Saint Petersburg, November 22–25, 2005). – V. L. Komarov Botanical Institute, St. Petersburg. [204 pp.]
- Ah-Peng C. & Bardat J. (2005): Check list of the bryophytes of Réunion Island (France). – Tropical Bryology 26: 89–118.
- Allen B. (2005): Maine mosses. *Sphagnaceae–Timmiaeae*. – Memoirs of the New York Botanical Garden 93: i–xii, 1–420.
- Åström M., Dynesius M., Hylander K. & Nilsson C. (2005): Effects of slash harvest on bryophytes and vascular plants in southern boreal forest clear-cuts. – Journal of Applied Ecology 42: 1194–1202.
- Bakalin V. A. (2005): Monograficheskaya obrabotka roda *Lophozia* (Dumort.) Dumort. s. str. – Nauka, Moskva. [240 pp.]
- Bakalin V. A. (2006): Ecological features and trends in the formation of the flora of liverworts under the impact of current volcanic and glacial activities: The example of Ushkovskii volcano, Kamchatka. – Russian Journal of Ecology 37: 13–18.
- Bao W. K. (2005): Structural features of *Polytrichum formosum* Hedw. populations along a habitat sequence of cutover restoration in the eastern Tibetan Plateau. – Ecological Research 20: 701–707.
- Bartholomew-Began S. & Jones N. L. (2005): Juvenile gametophyte development in the *Blasiales* (*Marchantiophyta*; *Metzgeriidae*). 1. Receptacular gemma/gemmaling ontogeny of *Blasia pusilla*. – Bryologist 108: 497–505.
- Bassi P., Basile A., Ferraro M., Masi M., Migliaccio D., Morelli G. & Napolitano E. (2006): Plasticity of repetitive DNA in response to metal stress in bryophytes. – Plant Biosystems 140: 80–86.
- Baumann M. & Müller F. (2006): Bryologische Beobachtungen im Erzgebirge II. – Limprichtia 29: 155–160.
- Bednarek-Ochyra H. & Ochyra R. (2005): Typification of *Codriophorus fascicularis* (Grimmiaceae). – Taxon 54: 1065–1074.
- Bennici A. (2005): A fundamental plant evolutionary problem: The origin of land-plant sporophyte; is a new hypothesis possible? – Rivista di Biologia – Biology Forum 98: 469–480.
- Bertram J. (2005): Herkunft und Bedeutung der Gattungsnamen der in Deutschland, in der Schweiz und in Österreich vorkommenden Moose. – Meylania 32 & 33: 1–76.
- Bharali B. & Bates J. W. (2006): Detoxification of dissolved SO<sub>2</sub> (bisulfite) by terricolous mosses. – Annals of Botany 97: 257–263.

- Bischler H., Boisselier-Dubayle M. C., Fontinha S. & Lambourdière J. (2006): Species boundaries in European and Macaronesian *Porella* L. (*Jungermanniales*, *Porellaceae*). – Cryptogamie Bryologie 27: 35–57.
- Bischler-Causse H., Gradstein S. R., Jovet-Ast S., Long D. G. & Salazar Allen N. (2005): *Marchantiidae*. – Flora Neotropica Monograph 97: 1–267.
- Blockeel T. L., Chlebicki A., Hájková P., Hájek M., Hradílek Z., Kürschner H., Ochyra R., Parolly G., Plášek V., Quandt D., Townsend C. C. & Vanderpoorten A. (2006): New national and regional bryophyte records, 12. – Journal of Bryology 28: 68–70.
- Booth R. K., Hotchkiss S. C. & Wilcox D. A. (2005): Discoloration of polyvinyl chloride (PVC) tape as a proxy for water-table depth in peatlands: validation and assessment of seasonal variability. – Functional Ecology 19: 1040–1047.
- Borges P. A. V., Cunha R., Gabriel R., Martins A. F., Silva L. & Vieira V. (eds.) (2005): Listagem da fauna e flora (*Mollusca* e *Arthropoda*) (*Bryophyta*, *Pteridophyta* e *Spermatophyta*) terrestres dos Açores [A list of terrestrial fauna (*Mollusca* and *Arthropoda*) and flora (*Bryophyta*, *Pteridophyta* e *Spermatophyta*) from the Azores. – Direcção Regional do Ambiente and Universidade dos Açores, Horta, Angra do Heroísmo and Ponta Delgada. [317 pp.]
- Bosanquet S. D. S. (2006): *Tortella bambergeri* (Schimp.) Broth. in the British Isles. – Journal of Bryology 28: 5–10.
- Botting R. S. & Fredeen A. L. (2006): Contrasting terrestrial lichen, liverwort, and moss diversity between old-growth and young second-growth forest on two soil textures in central British Columbia. – Canadian Journal of Botany 84: 120–132.
- Bradfield G. E. & Sadler K. D. (2006): Transient assemblage dynamics of terrestrial bryophytes in a subalpine forest. – Bryologist 109: 18–25.
- Bruggeman-Nannenga M. A. (2006): Bryophyte flora of Uganda. 6. *Fissidentaceae* (part 1). – Journal of Bryology 28: 53–62.
- Bruun H. H., Moen J., Virtanen R., Grytnes J. A., Oksanen L. & Angerbjörn A. (2006): Effects of altitude and topography on species richness of vascular plants, bryophytes and lichens in alpine communities. – Journal of Vegetation Science 17: 37–46.
- Buczkowska K., Adamczak M., Chudzińska E., Wachowiak W. & Baczkiewicz A. (2006): In vitro propagation of cryptic species of *Aneura pinguis* (*Hepaticae*, *Metzgeriales*). – Cryptogamie Bryologie 27: 241–251.
- Burgisser L. & Price M. J. (2005): Inventory of the moss flora of the Canton of Geneva (Switzerland). Bibliographic catalogue (1791–2001). – Candollea 60: 513–562.
- Bykowszczenko N., Baranowska-Bosiacka I., Bosiacka B., Kaczmarek A. & Chlubek D. (2006): Determination of heavy metal concentration in mosses of Słowiński National Park using atomic absorption spectrometry and neutron activation analysis methods. – Polish Journal of Environmental Studies 15: 41–46.
- Cano M. J., Werner O. & Guerra J. (2005): A morphometric and molecular study in *Tortula subulata* complex (*Pottiaceae*, *Bryophyta*). – Botanical Journal of the Linnean Society 149: 333–350.
- Cantonati M., Gerecke R. & Bertuzzi E. (2006): Springs of the Alps – sensitive ecosystems to environmental change: From biodiversity assessments to long-term studies. – Hydrobiologia 562: 59–96.
- Cao T., Zhu R. L., Tan B. C., Guo S. L., Gao C., Wu P. C. & Li X. J. (2006): A report of the first national red list of Chinese endangered bryophytes. – Journal of the Hattori Botanical Laboratory 99: 275–295.
- Casas C., Brugués M., Cros R. M., Saez L. & Balaguer P. (2004): Referencias bibliográficas sobre la flora briofítica de las Islas Baleares. – Boletín de la Sociedad Española de Briología 25: 33–40.
- Chung S. J., Takechi K., Sakai A., Ono K. & Takano H. (2006): Detection of genes differentially expressed in cultured cells of *Marchantia polymorpha*, but not in *Arabidopsis thaliana*, using an *Arabidopsis* cDNA macroarray. – Journal of the Hattori Botanical Laboratory 99: 245–257.
- Chung S. J., Takechi K., Sakai A., Ono K. & Takano H. (2006): Profiling of bryophyte gene expression by hybridization of an *Arabidopsis* cDNA array with bryophyte cDNA. – Journal of the Hattori Botanical Laboratory 99: 233–244.
- Cleavitt N. (2005): How many bryophytes are there in New York and are any of them rare? – Solidago 6: 1–6.
- Cleavitt N. L. (2005): Patterns, hypotheses and processes in the biology of rare bryophytes. – Bryologist 108: 554–566.
- Clerc P. & Price M. J. (2005): La cryptogamie aux Conservatoire et Jardin botaniques de la Ville de Genève (CJB). – Saussurea 35: 53–72.

- Cove D. (2005): The moss *Physcomitrella patens*. – Annual Review of Genetics 39: 339–358.
- Crandall-Stotler B. J., Stotler R. E. & Doyle W. T. (2006): On *Anthoceros phymatodes* M. Howe and the hornwort genus *Phymatoceros* Stotler, W.T. Doyle & Crand.-Stotl. (*Anthocerotophyta*). – Cryptogamie Bryologie 27: 59–73.
- Cronberg N., Rydgren K. & Økland R. H. (2006): Clonal structure and genet-level sex ratios suggest different roles of vegetative and sexual reproduction in the clonal moss *Hylocomium splendens*. – Ecography 29: 95–103.
- Crosby M. R. & Magill R. E. (2005): Index of Bryophytes 2001–2004. – Missouri Botanical Garden St. Louis [31 pp.]
- Cykowska B. (2005): *Tetraplodon mnioides* (Bryopsida, Splachnaceae) in the Tatra National Park (Poland). 159. – Polish Botanical Journal 50: 159–161.
- Dauphin G. (2005): Catalogue of Costa Rican Hepaticae and Anthocerotae. – Tropical Bryology 26: 141–218.
- Dia M. G. & Campisi P. (2006): First record of *Pseudocrossidium obtusulum* (Lindb.) H.A. Crum & L.E. Anderson (Pottiaceae, Musci) in Italy. – Nova Hedwigia 82: 205–208.
- Dierßen K. & Dierßen B. (2005): Studies on the vegetation of fens, springs and snow fields in West Greenland. – Phytocoenologia 35: 849–885.
- Dorrepaal E., Aerts R., Cornelissen J. H. C., van Logtestijn R. S. P. & Callaghan T. V. (2006): *Sphagnum* modifies climate-change impacts on subarctic vascular bog plants. – Functional Ecology 20: 31–41.
- Drea S. C., Lao N. T., Wolfe K. H. & Kavanagh T. A. (2006): Gene duplication, exon gain and neofunctionalization of *OEP16*-related genes in land plants. – Plant Journal 46: 723–735.
- Duckett J. G. & Ligrone R. (2006): Novel features of the plastids in some deep-shade, antipodean thalloid liverworts. – Cryptogamie Bryologie 27: 75–83.
- Duff R. J. (2006): Divergent RNA editing frequencies in hornwort mitochondrial *nad5* sequences. – Gene 366: 285–291.
- Dulger B., Yayintas O. T. & Gonuz A. (2005): Antimicrobial activity of some mosses from Turkey. – Fitoterapia 76: 730–732.
- Dürhammer O., Köckinger H. & Reimann M. (2005): Beiträge zur Kryptogamenflora im Gebiet der Neuen Regensburger Hütte (Stubaier Alpen, Österreich). Teil III: Moose. – Hoppea 66: 615–627.
- Eggers J. (2005): Ergänzungsband zu J.-P. Frahm & J. Eggers „Lexikon deutschsprachiger Bryologen“. – Limprichtia 27: 1–245.
- Engel J. J. (2006): Austral Hepaticae 40. *Tritomaria* (Jungermanniaceae subfam. *Lophozioideae*) new to the south temperate, together with a new subspecies, *T. exsecta* subsp. *novaehollandiae* subsp. nov. – Bryologist 109: 60–67.
- Erzberger P. (2005): The bulbiliferous species of *Pohlia* (Bryaceae, Musci) in Hungary. – Studia Botanica Hungarica 36: 67–75.
- Eskelinen A. & Oksanen J. (2006): Changes in the abundance, composition and species richness of mountain vegetation in relation to summer grazing by reindeer. – Journal of Vegetation Science 17: 245–254.
- Estébanez B., Yamaguchi T. & Deguchi H. (2006): Ultrastructure of the spore in four Japanese species of *Ptychomitrium* Fürnr. (Musci). – Grana 45: 61–70.
- Euler A., Fujiwara K. & Saraiva R. (2005): Species richness, community structure and other characteristics of a tropical rainforest in the Southwestern Amazon, Acre, Brazil. – Hikobia 14: 273–291.
- Fenton N. J. & Bergeron Y. (2006): Facilitative succession in a boreal bryophyte community driven by changes in available moisture and light. – Journal of Vegetation Science 17: 65–76.
- Fenton N. J. & Bergeron Y. (2006): *Sphagnum* spore availability in boreal forests. – Bryologist 109: 173–181.
- Fernandez C. C., Shevock J. R., Glazer A. N. & Thompson J. N. (2006): Cryptic species within the cosmopolitan desiccation-tolerant moss *Grimmia laevigata*. – Proceedings of the National Academy of Sciences of the United States of America 103: 637–642.
- Fernández J. A., Vázquez M. D., López J. & Carballeira A. (2006): Modelling the extra and intracellular uptake and discharge of heavy metals in *Fontinalis antipyretica* transplanted along a heavy metal and pH contamination gradient. – Environmental Pollution 139: 21–31.
- Fife A. J. & Dalton P. J. (2005): A reconsideration of *Pleurophascum* (Musci: Pleurophascaceae) and specific status for a New Zealand endemic, *Pleurophascum ovalifolium* stat. et nom. nov. – New Zealand Journal of Botany 43: 871–884.
- Figueiredo A. C., Barroso J. G., Pedro L. G., Fontinha S. S., Sim-Sim M., Sérgio C., Luis L. & Scheffer J. J. C. (2006): *Asterella africana* (Mont.) A. Evans grown on Madeira and in mainland Portugal: morphological data and composition of the essential oil. – Flavour and Fragrance Journal 21: 534–538.

- Fisher K. M. (2006): Rank-free monography: A practical example from the moss clade *Leucophanella* (*Calymperaceae*). – Systematic Botany 31: 13–30.
- Flatberg K. I., Thingsgaard K. & Såstad S. M. (2006): Interploidal gene flow and introgression in bryophytes: *Sphagnum girgensohnii* × *Sphagnum russowii*, a case of spontaneous neotriploidy. – Journal of Bryology 28: 27–37.
- Frahm J.-P. (2004): Guide to the bryological hotspots of the world Part 1. Azores. – Archive for Bryology 3: 1–14.
- Frahm J.-P. & Newton A. E. (2005): A new contribution to the moss flora of Dominican amber. – Bryologist 108: 526–536.
- Franzle O. (2006): Complex bioindication and environmental stress assessment. – Ecological Indicators 6: 114–136.
- Garcia C., Lara F., Sergio C., Sim-Sim M., Garilleti R. & Mazimpaka V. (2006): *Zygodon catarinoi* (*Orthotrichaceae, Bryopsida*), a new epiphytic species from the Western Mediterranean Basin. – Nova Hedwigia 82: 247–256.
- Germano S. R. & Porto K. C. (2006): Bryophyte communities in an Atlantic forest remnant, state of Pernambuco, Brazil. – Cryptogamie Bryologie 27: 153–163.
- Glime J. M. (2006): Bryophytes and herbivory. – Cryptogamie Bryologie 27: 191–203.
- Goncharova I. A. (2005): On the sod structure and productivity of *Sphagnum* mosses on oligotrophic swamps. – Sibirskiy Ekologicheskiy Zhurnal 12: 131–134.
- Gradstein S. R. (2006): The lowland cloud forest of French Guiana – a liverwort hotspot. – Cryptogamie Bryologie 27: 141–152.
- Gramatica P., Battaini F., Giani E., Papa E., Jones R. J. A., Preatoni D. & Cenci R. M. (2006): Analysis of mosses and soils for quantifying heavy metal concentrations in Sicily: A multivariate and spatial analytical approach. – Environmental Science and Pollution Research 13: 28–36.
- Green T. G. A., Kulle D., Pannewitz S., Sancho L. G. & Schroeter B. (2005): UV-A protection in mosses growing in continental Antarctica. – Polar Biology 28: 822–827.
- Grundmann M., Schneider H., Russell S. J. & Vogel J. C. (2006): Phylogenetic relationships of the moss genus *Pleurochaete* Lindb. (*Bryales: Pottiaceae*) based on chloroplast and nuclear genomic markers. – Organisms Diversity & Evolution 6: 33–45.
- Guerra J. & Cros R.M. (eds.) (2006): Flora Briofitica Ibérica. Vol. III, *Pottiales: Pottiaceae, Encalyptales: Encalyptaceae*. – Universidad de Murcia, SEB, Murcia. [305 pp.]
- Haig D. & Wilczek A. (2006): Sexual conflict and the alternation of haploid and diploid generations. – Philosophical Transactions of the Royal Society B – Biological Sciences 361: 335–343.
- Harris A., Bryant R. G. & Baird A. J. (2006): Mapping the effects of water stress on *Sphagnum*: Preliminary observations using airborne remote sensing. – Remote Sensing of Environment 100: 363–378.
- Harrison S., Davies K. F., Safford H. D. & Viers J. H. (2006): Beta diversity and the scale-dependence of the productivity-diversity relationship: a test in the Californian serpentine flora. – Journal of Ecology 94: 110–117.
- Hasse T. & Daniëls F. J. A. (2006): Species responses to experimentally induced habitat changes in a *Corynephorus* grassland. – Journal of Vegetation Science 17: 135–146.
- He-Nygrén X., Juslén A., Ahonen I., Glenny D. & Piippo S. (2006): Illuminating the evolutionary history of liverworts (*Marchantiophyta*) – Towards a natural classification. – Cladistics 22: 1–31.
- Heber U., Lange O. L. & Shuvalov V. A. (2006): Conservation and dissipation of light energy as complementary processes: homoiohydric and poikilohydric autotrophs. – Journal of Experimental Botany 57: 1211–1223.
- Heinrichs J., Lindner M., Groth H., Hentschel J., Feldberg K., Renker C., Engel J. J., Konrat M. von, Long D. G. & Schneider H. (2006): Goodbye or welcome Gondwana? – insights into the phylogenetic biogeography of the leafy liverwort *Plagiochila* with a description of *Proskauera*, gen. nov. (*Plagiochilaceae, Jungermanniales*). – Plant Systematics and Evolution 258: 227–250.
- Hentschel J., Wilson R., Burghardt M., Zündorf H.-J., Schneider H. & Heinrichs J. (2006): Reinstatement of *Lophocoleaceae* (*Jungermanniopsida*) based on chloroplast gene *rbcL* data: exploring the importance of female involucres for the systematics of *Jungermanniales*. – Plant Systematics and Evolution 258: 211–226.
- Hirakawa N., Nagashima K. & Nakagoshi N. (2005): Stand structure and homogeneity of an evergreen broad-leaved coppice forest. – Hikobia 14: 293–303.

- Hugonnot V. & Schill D. B. (2006): *Mannia californica* (Gottsche ex Underw.) L.C. Wheeler (*Aytoniaceae, Marchantiales*) in Ardèche (France), new to Europe. – *Cryptogamie Bryologie* 27: 181–189.
- Ignatov M. S., Huttunen S. & Koponen T. (2005): Bryophyte flora of Hunan Province, China. 5. *Brachytheciaceae (Musci)*, with an overview of *Eurhynchiadelphus* and *Rhynchostegiella* in SE Asia. – *Acta Botanica Fennica* 178: 1–56.
- Ignatov M. S., Ignatova E. A., Konstantinova N. A., Pronkina G. A., Urbanavichus G. P. & Urbanavichene I. N. (2004): The present-day state of biological diversity within protected areas in Russia. Issue 3. Lichens and Bryophytes. – Ministry of Natural Resources of the Russian Federation, Moscow [369 pp.]
- Iida K. & Go M. (2006): Survey of conserved alternative splicing events of mRNAs encoding SR proteins in land plants. – *Molecular Biology and Evolution* 23: 1085–1094.
- Iijima H., Shibuya M., Saito H. & Takahashi K. (2006): The water relation of seedlings of *Picea jezoensis* on fallen logs. – *Canadian Journal of Forest Research* 36: 664–670.
- Infante M. & Heras P. (2005): Bryophytes in the Spanish law. – *Boletín de la Sociedad Española de Briología* 26–27: 5–16.
- Iwatsuki Z. & Suzuki T. (2006): A taxonomic revision of *Trematodon asanoi* and its related species (*Dicranaceae, Musci*). – *Journal of the Hattori Botanical Laboratory* 99: 259–269.
- Jones P. R. & Rosentreter R. (2006): Gametophyte fragment growth of three common desert mosses on artificial and natural substrates. – *Bryologist* 109: 166–172.
- Kade A., Walker D. A. & Raynolds M. K. (2005): Plant communities and soils in cryoturbated tundra along a bioclimate gradient in the Low Arctic, Alaska. – *Phytocoenologia* 35: 761–820.
- Kamisugi Y., Cuming A. C. & Cove D. J. (2005): Parameters determining the efficiency of gene targeting in the moss *Physcomitrella patens*. – *Nucleic Acids Research* 33: e173.
- Kapila S., Kaur R. & Saxena P. (2005): Effect of nickel and cadmium on regeneration of *Hyophila involuta* and *Funaria hygrometrica*. – *Hikobia* 14: 245–249.
- Khorkavciv O. Ya., Demkiv O. T. & Ripetskyi R. T. (2006): Hormonal'na reghuliatsia rozvytku aposporychnogho ghametofita *Pottia intermedia* (Furnr.) Turn. – *Ukrains'kyi Botanichnyi Zhurnal* 63: 70–80.
- Kobayashi F., Kofuji R., Yamashita Y. & Nakamura Y. (2006): A novel treatment system of wastewater contaminated with copper by a moss. – *Biochemical Engineering Journal* : 28: 295–298.
- Kürschner H., Parolly G. & Erdağ A. (2006): Life forms and life strategies of epiphytic bryophytes in *Quercus vulcanica* forests of Turkey. – *Nova Hedwigia* 82: 331–347.
- Le Roux G., Aubert D., Stille P., Krachler M., Kober B., Cheburkin A., Bonani G. & Shotyk W. (2005): Recent atmospheric Pb deposition at a rural site in southern Germany assessed using a peat core and snowpack, and comparison with other archives. – *Atmospheric Environment* 39: 6790–6801.
- Leonardia A. A. P., Kumar P. P. & Tan. B. C. (2006): Development of microsatellite markers for the tropical moss, *Acanthorrhynchium papillatum*. – *Molecular Ecology Notes* 6: 396–398.
- León-Vargas Y., Engwald S. & Proctor M. C. F. (2006): Microclimate, light adaptation and desiccation tolerance of epiphytic bryophytes in two Venezuelan cloud forests. – *Journal of Biogeography* 33: 901–913.
- Lo Giudice R. & Cristaldo A. (2004): Chorological and ecological survey on the vascular and bryophytic flora in Etna territory (Erei Mountains, C Sicily). – *Flora Mediterranea* 14: 357–417.
- Lobel S., Snäll T. & Rydin H. (2006): Species richness patterns and metapopulation processes – evidence from epiphyte communities in boreo-nemoral forests. – *Ecography* 29: 169–182.
- Machida M., Takechi K., Sato H., Chung S. J., Kuroiwa H., Takio S., Seki M., Shinozaki K., Fujita T., Hasebe M. & Takano H. (2006): Genes for the peptidoglycan synthesis pathway are essential for chloroplast division in moss. – *Proceedings of the National Academy of Sciences of the United States of America* 103: 6753–6758.
- Mallón R., Reinoso J., Rodríguez-Oubiña J. & González M. L. (2006): In vitro development of vegetative propagules in *Splachnum ampullaceum*: brood cells and chloronematal bulbils. – *Bryologist* 109: 215–223.
- Markham K., Chalk T. & Stewart C. N. (2006): Evaluation of fern and moss protein-based defenses against phytophagous insects. – *International Journal of Plant Sciences* 167: 111–117.
- Marques J., Hespanhol H., Vieira C. & Séneca A. (2005): Comparative study of the bryophyte epiphytic vegetation in *Quercus pyrenaica* and *Quercus robur* woodlands from northern Portugal. – *Boletín de la Sociedad Española de Briología* 26–27: 75–84.
- Maslovsky O. (2005): Rare and threatened bryophytes and a proposal for an eastern European Red Book. – *Boletín de la Sociedad Española de Briología* 26–27: 47–54.

- McDaniel S. F. (2005): Genetic correlations do not constrain the evolution of sexual dimorphism in the moss *Ceratodon purpureus*. – Evolution 59: 2353–2361.
- McLetchie D. N. & Stark L. R. (2006): Sporophyte and gametophyte generations differ in their thermotolerance response in the moss *Microbryum*. – Annals of Botany 97: 505–511.
- Meier E., Paal J., Liira J. & Jüriado I. (2005): Influence of tree stand age and management on the species diversity in Estonian eutrophic alvar and boreo-nemoral *Pinus sylvestris* forests. – Scandinavian Journal of Forest Research 20, Suppl. 6: 135–144.
- Miller N. G., Fryday A. M. & Hinds J. W. (2005): Bryophytes and lichens of a calcium-rich spring seep isolated on the granitic terrain of Mt. Katahdin, Maine, USA. – Rhodora 107: 339–358.
- Muir P. S., Norman K. N. & Sikes K. G. (2006): Quantity and value of commercial moss harvest from forests of the Pacific Northwest and Appalachian regions of the U.S. – Bryologist 109: 197–214.
- Nair M. C., Rajesh K. P. & Madhusoodanan P. V. (2005): Bryophytes of Wayanad in Western Ghats. – Malabar Natural History Society, Kerala, India. [iv + 284 pp.]
- Natcheva R. (2005): Three new species for the bryophyte flora of Bulgaria. – Phytologia Balcanica 11: 33–34.
- Navrátilová J. & Hájek M. (2005): Recording relative water table depth using PVC tape discolouration: Advantages and constraints in fens. – Applied Vegetation Science 8: 21–26.
- Niinemets Ü. & Kull K. (2005): Co-limitation of plant primary productivity by nitrogen and phosphorus in a species-rich wooded meadow on calcareous soils. – Acta Oecologica – International Journal of Ecology 28: 345–356.
- Nimmo D. W. R., Herrmann S. J., Romine J. A., Meyer K. W., Johnson R. W., Davies P. H., Hansen D. L. & Brinkman S. F. (2006): Cadmium and zinc accumulation in aquatic bryophytes immersed in the Arkansas River, Colorado: Comparison of fall versus spring. – Journal of Freshwater Ecology 21: 331–339.
- Nordin A., Strengbom J. & Ericson L. (2006): Responses to ammonium and nitrate additions by boreal plants and their natural enemies. – Environmental Pollution 141: 167–174.
- Ochyra R. & Bednarek-Ochyra H. (2006): *Platyhypnidium lusitanicum* versus *P. alopecuroides* (*Brachytheciaceae*). – Journal of Bryology 28: 64–66.
- Odland A. (2005): Oligotrophic and mesotrophic vegetation in southern Scandinavian mountains. Gradients in species and community distribution extracted by numerical analyses of earlier published vegetation descriptions. – Phytocoenologia 35: 985–1018.
- Ódor P., van Dort K., Aude E., Heilmann-Clausen J. & Christensen M. (2005): Diversity and composition of dead wood inhabiting bryophyte communities in European beech forests. – Boletín de la Sociedad Española de Briología 26–27: 85–102.
- Oliver M. J., Velten J. & Mishler B. D. (2005): Desiccation tolerance in bryophytes: A reflection of the primitive strategy for plant survival in dehydrating habitats? – Integrative and Comparative Biology 45: 788–799.
- Olofsson J. (2006): Short- and long-term effects of changes in reindeer grazing pressure on tundra heath vegetation. – Journal of Ecology 94: 431–440.
- Paavola R., Muotka T., Virtanen R., Heino J., Jackson D. & Mäki-Petäys A. (2006): Spatial scale affects community concordance among fishes, benthic macroinvertebrates, and bryophytes in streams. – Ecological Applications 16: 368–379.
- Papp B. & Erzberger P. (2005): The bryophyte flora of Golija-Studenica Biosphere Reserve and some adjacent sites (SW Serbia, Serbia-Montenegro). – Studia Botanica Hungarica 36: 101–116.
- Papp B., Ódor P. & Szurdoki E. (2005): Methodological overview and a case study of the Hungarian bryophyte monitoring program. – Boletín de la Sociedad Española de Briología 26–27: 23–32.
- Peck J. E. (2006): Towards sustainable commercial moss harvest in the Pacific Northwest of North America. – Biological Conservation 128: 289–297.
- Peck J. E. & Christy J. A. (2006): Putting the stewardship concept into practice: Commercial moss harvest in Northwestern Oregon, USA. – Forest Ecology and Management 225: 225–233.
- Pedersen N., Russell S. J., Newton A. E. & Ansell S. W. (2006): A novel molecular protocol for the rapid extraction of DNA from bryophytes and the utility of direct amplification of DNA from a single dwarf male. – Bryologist 109: 257–264.
- Pedrotti C. C. (2005): Flora dei muschi d’Italia. *Bryopsida* (II parte). – Antonio Delfino Editore, Roma. [432 pp.]

- Pence V. C., Dunford S. S. & Redella S. (2005): Differential effects of abscisic acid on desiccation tolerance and carbohydrates in three species of liverworts. – *Journal of Plant Physiology* 162: 1331–1337.
- Pentecost A. & Zhang Z. H. (2006): Response of bryophytes to exposure and water availability on some European travertines. – *Journal of Bryology* 28: 21–26.
- Pesch R. & Schröder W. (2006): Mosses as bioindicators for metal accumulation: Statistical aggregation of measurement data to exposure indices. – *Ecological Indicators* 6: 137–152.
- Pesch R. & Schröder W. (2006): Spatiotemporal variability of metal accumulation in mosses. Analysis of measurement data and metadata by statistics and GIS. – *Nova Hedwigia* 82: 447–466.
- Peshkova N. V. & Andreyashkina N. I. (2006): Indicator aspect of the geographic analysis of plant communities with respect to floristic composition on mountain slopes differing in exposure (the Polar Urals). – *Russian Journal of Ecology* 37: 103–108.
- Pierrot R. B. (2005): Les bryophytes du Centre-Ouest: Classification, détermination, répartition. Deuxième tirage. – *Bulletin de la Société Botanique du Centre-Ouest, Nouvelle Série*, 5: 1–123.
- Pfeiffer T., Fritz S., Stech M. & Frey W. (2006): Vegetative reproduction and clonal diversity in *Rhytidium rugosum* (*Rhytidaceae, Bryopsida*) inferred by morpho-anatomical and molecular analyses. – *Journal of Plant Research* 119: 125–135.
- Pokorný L., Lara F. & Mazimpaka V. (2006): The bryophyte flora of the city of Trento (North Italy). – *Cryptogamie Bryologie* 27: 265–284.
- Porley R. D. (2005): The conservation of *Didymodon glaucus* Ryan in Britain. – *Boletín de la Sociedad Española de Briología* 26–27: 55–58.
- Price M. J. (2005): Catalogue of the Hedwig-Schwägrichen Herbarium (G). Part 1. Type material and a review of typifications for the Hedwig moss names. – *Boissiera* 61: 5–388.
- Puche F., Casas C. & Brugués M. (2006): *Didymodon eckeliae* (*Pottiaceae*), new to Europe. – *Bryologist* 109: 239–241.
- Rixen C. & Mulder C. P. H. (2005): Improved water retention links high species richness with increased productivity in arctic tundra moss communities. – *Oecologia* 146: 287–299.
- Romero C., Putz F. E. & Kitajima K. (2006): Ecophysiology in relation to exposure of pendant epiphytic bryophytes in the canopy of a tropical montane oak forest. – *Biotropica* 38: 35–41.
- Rydgren K., Cronberg N. & Økland R. H. (2006): Factors influencing reproductive success in the clonal moss, *Hylocomium splendens*. – *Oecologia* 147: 445–454.
- Ryoma R. & Laaka-Lindberg S. (2005): Bryophyte recolonization on burnt soil and logs. – *Scandinavian Journal of Forest Research* 20, Suppl. 6: 5–16.
- Saavedra L., Svensson J., Carballo V., Izmendi D., Welin B. & Vidal S. (2006): A dehydrin gene in *Physcomitrella patens* is required for salt and osmotic stress tolerance. – *Plant Journal* 45: 237–249.
- Sabovljević A., Soković M., Sabovljević M. & Grubišić D. (2006): Antimicrobial activity of *Bryum argenteum*. – *Fitoterapia* 77: 144–145.
- Sass-Gyarmati A., Pócs T. & Orbán S. (2005): Contribution to the knowledge of the bryoflora of Detunata Nature Reserve (Metaliferi Mountains, Transylvania). – *Studia Botanica Hungarica* 36: 123–130.
- Saxena A. (2006): Seasonal pattern of metal bioaccumulation and their toxicity on *Sphagnum squarrosum*. – *Journal of Environmental Biology* 27: 71–75.
- Scarlett P. & O'Hare M. (2006): Community structure of in-stream bryophytes in English and Welsh rivers. – *Hydrobiologia* 553: 143–152.
- Schlüsslmayr G. (2005): Soziologische Moosflora des südöstlichen Oberösterreich. – *Stapfia* 84: i–xi, 1–695. [Biology Centre of the Upper Austrian Museums, Linz]
- Schmidt I., Zerbe S., Betzin J. & Weckesser M. (2006): An approach to the identification of indicators for forest biodiversity – The Solling mountains (NW Germany) as an example. – *Restoration Ecology* 14: 123–136.
- Schröck C. (2004): Bryofloristische Untersuchungen im Wildgerlostal (Nationalpark Hohe Tauern, Salzburg, Österreich). – *Sauteria* 13: 365–428.
- Schulte J., Erxleben A., Schween G. & Reski R. (2006): High throughput metabolic screen of *Physcomitrella* transformants. – *Bryologist* 109: 247–256.
- Schumacker R. & Váňa J. (2005): Identification keys to the liverworts and hornworts of Europe and Macaronesia (distribution and status). 2<sup>nd</sup> ed. – Sorus, Poznań. [210 pp.]
- Seaward M. R. D., Ellis L. T., Pócs T. & Wigginton M. J. (2006): Bryophyte flora of the Chagos Archipelago. – *Journal of Bryology* 28: 11–19.

- Sérgio C. (2006): A review of the *Gymnostomum calcareum* Nees & Hornsch complex (*Bryopsida* : *Pottiaceae*) in southern Europe and the Macaronesian Islands, including *Gymnostomum calcareum* var. *atlanticum* var. nov. – *Journal of Bryology* 28: 38–45.
- Shigyo M., Hasebe M. & Ito M. (2006): Molecular evolution of the AP2 subfamily. – *Gene* 366: 256–265.
- Smith J. L. (2006): A bryological assessment of the Flint Woods Preserve, Delaware. – *Northeastern Naturalist* 13: 117–130.
- Solga A. & Frahm J.-P. (2006): Nitrogen accumulation by six pleurocarpous moss species and their suitability for monitoring nitrogen deposition. – *Journal of Bryology* 28: 46–52.
- Sollman P. (2005): A taxonomic revision of the genus *Chionoloma* Dixon (*Musci, Pottiaceae*) in Africa. – *Tropical Bryology* 26: 35–38.
- Stark L. R. & McLetchie D. N. (2006): Gender-specific heat-shock tolerance of hydrated leaves in the desert moss *Syntrichia caninervis*. – *Physiologia Plantarum* 126: 187–195.
- Streiff A. (2006): Phylogenetic study of *Grimmia* (*Grimmiaceae*) based on plastid DNA sequences (*trnL-trnF* and *rps4*) and on morphological characters. – *Bryologist* 109: 224–235.
- Sugita H. & Nagaike T. (2005): Microsites for seedling establishment of subalpine conifers in a forest with moss-type undergrowth on Mt. Fuji, central Honshu, Japan. – *Ecological Research* 20: 678–685.
- Suleiman M., Akiyama H. & Tan. B. C. (2006): A revised catalogue of mosses reported from Borneo. – *Journal of the Hattori Botanical Laboratory* 99: 107–183.
- Tanskanen H., Granström A., Venäläinen A. & Puttonen P. (2006): Moisture dynamics of moss-dominated surface fuel in relation to the structure of *Picea abies* and *Pinus sylvestris* stands. – *Forest Ecology and Management* 226: 189–198.
- Tews J. (2004): Hummock vegetation at the arctic tree-line near Churchill, Manitoba. – *Canadian Field-Naturalist* 118: 590–594.
- Turner P. A. M., Kirkpatrick J. B. & Pharo E. J. (2006): Bryophyte relationships with environmental and structural variables in Tasmanian old-growth mixed eucalypt forest. – *Australian Journal of Botany* 54: 239–247.
- Váňa J. & Gremmen N. (2006): Checklist of the hepatic flora of sub-Antarctic Iles Kerguelen. – *Cryptogamie Bryologie* 27: 131–139.
- Vanderpoorten A., Sotiaux A. & Engels P. (2006): A GIS-based model of the distribution of the rare liverwort *Aneura maxima* at the landscape scale for an improved assessment of its conservation status. – *Biodiversity and Conservation* 15: 829–838.
- Vieira C., Sérgio C. & Séneca A. (2005): Threatened bryophytes occurrence in Portuguese stream habitat. – *Boletín de la Sociedad Española de Briología* 26–27: 103–118.
- Villarreal J. C. & Renzaglia K. S. (2006): Structure and development of *Nostoc* strands in *Leiosporoceros dussii* (*Anthocerotophyta*): A novel symbiosis in land plants. – *American Journal of Botany* 93: 693–705.
- Virchenko V. M. (2005): Rid *Zygodon* Hook. et Taylor (*Orthotrichaceae, Bryophyta*) v Ukraini. – *Ukrajins'kyi Botanichnyi Zhurnal* 62: 715.
- Wasley J., Robinson S. A., Lovelock C. E. & Popp M. (2006): Some like it wet – biological characteristics underpinning tolerance of extreme water stress events in Antarctic bryophytes. – *Functional Plant Biology* 33: 443–455.
- Werner J., Schneider T., Schneider C. & Mahevas T. (2005): The bryophytes of Lorraine (France), outside the Vosges mountains, a critical annotated list. – *Cryptogamie Bryologie* 26: 347–402.
- Whinam J. & Copson G. (2006): *Sphagnum* moss: an indicator of climate change in the sub-Antarctic. – *Polar Record* 42: 43–49.
- Yamada K. & Iwatsuki Z. (2006): Catalog of the hepatics of Japan. – *Journal of the Hattori Botanical Laboratory* 99: 1–106.
- Yazaki T., Urano S. & Yabe K. (2006): Water balance and water movement in unsaturated zones of *Sphagnum* hummocks in Fuhrengawa Mire, Hokkaido, Japan. – *Journal of Hydrology* 319: 312–327.
- Zander R. H., Jiménez J. A. & Sagar T. (2005): *Didymodon bistratosus* (*Pottiaceae*) in the New World. – *Bryologist* 108: 540–543.
- Zartman C. E. & Shaw A. J. (2006): Metapopulation extinction thresholds in rain forest remnants. – *American Naturalist* 167: 177–189.
- Zechmeister H. G. (2005): Die Moosflora der Serpentinrasen im Burgenland. – *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Österreich* 142: 9–15.
- Zhang B.-H., Pan X.-P., Cannon C. H., Cobb G. P. & Anderson T. A. (2006): Conservation and divergence of plant microRNA genes. – *Plant Journal* 46: 243–259.

## Bryologické publikace z České a Slovenské republiky [Bryological publications issued in the Czech Republic and Slovakia]

### 1. Články a knižní publikace [Papers and books]

- Duda J. & Duda J. (2004): Mechiorsty okolí obce Donovaly na Slovensku. – Časopis Slezského Zemského Muzea 53: 283–287.
- Duda J. & Duda J. (2004): Doplněk k poznání mechiorstů Lubovnianské vrchoviny [Complement to the knowledge of bryophytes of Lubovnianská highlands (Slovak Republic)]. – Naturaе Tutela 8: 203–207.
- Franklová H. & Kolbek J. (2005): Mechiorsty Přírodních rezervací Na Babě a Brdatka v CHKO a BR Křivoklátsko [Bryophytes of the nature reserves Na Babě and Brdatka in the Křivoklátsko Protected Landscape Area and Biosphere Reserve (central Bohemia)]. – Zprávy České Botanické Společnosti 40: 151–158.
- González M. L., Mallón R., Reinoso J. & Rodríguez-Oubiña J. (2006): In vitro micropropagation and long-term conservation of the endangered moss *Splachnum ampullaceum*. – Biologia Plantarum 50: 339–345.
- Hofhanzlová E., Ekrt L. & Štechová T. (2005): Floristický a vegetační průzkum rašelinště Na Klátově [Floristic and vegetation research in the „Na Klátově“ peat bog]. – Acta Rerum Naturalium 1: 45–52.
- Horáková V., Kubešová S. & Šumberová K. (2005): Mechiorsty rybníků a sádek v Českobudějovické páni [Bryophytes of fishpond bottoms and storage ponds in the Českobudějovická pánev basin (Czech Republic)]. – Bryonora 35: 9–17.
- Hradilek Z. (2005): Mechiorsty zaznamenané v průběhu 11. Bryologicko-lichenologických dnů v Březníku u Náměště nad Oslavou [Bryophytes recorded during the 11th Bryological and Lichenological Days in Březník near the town of Náměšť nad Oslavou (south-western Moravia, Czech Republic)]. – Bryonora 36: 11–14.
- Jakšičová T. (2004): Mech *Hamatocaulis vernicosus* (Mitt.) Hedenäs v PR Bažiny. – Orchis, Zpravodaj Botanické Sekce 23: 2–6.
- Jansová I. & Soldán Z. (2006): The habitat factors that affect the composition of bryophyte and lichen communities on fallen logs. – Preslia 78: 67–86.
- Kalina T. & Váňa J. (2005): Sinice, řasy, houby mechiorsty a podobné organismy v současné biologii. – Karolinum, Praha. [608 pp.]
- Karofeld E. & Pajula R. (2005): Distribution and development of necrotic *Sphagnum* patches in two Estonian raised bogs. – Folia Geobotanica 40: 357–366.
- Kresáňová K., Mišiková-Janovicová K. & Kubinská A. (2005): Diversity of bryophytes in agrocoenoses of Slovakia. – Biologia, Sect. Botany, 60: 9–15.
- Kučera J. (2005): Nová bryologická literatura XIV. [New bryological literature, XIV]. – Bryonora 36: 45–68.
- Kučera J. (ed.) (2005): Zajímavé bryofloristické nálezy V. [Interesting bryofloristic records, VI]. – Bryonora 35: 32–35. [*Frullania inflata*, *Scapania lingulata*, *Scapania aspera*, *Aloina aloides* var. *ambigua*, *Bryum gemmiferum*, *Buxbaumia aphylla*, *Campylophyllum calcareum*, *Didymodon cordatus*, *Didymodon glaucus*, *Didymodon sinuosus*, *Fissidens arnoldii*, *Orthotrichum patens*, *Pohlia lescuriana*, *Rhynchostegiella tenella*, *Rhynchostegium confertum*, *Rhynchostegium megapolitanum*, *Splachnum ampullaceum*]
- Kučera J. (ed.) (2005): Zajímavé bryofloristické nálezy VI. [Interesting bryofloristic records, VII]. – Bryonora 36: 26–34. [*Anastrophyllum hellerianum*, *Cololejeunea calcarea*, *Geocalyx graveolens*, *Harpanthus scutatus*, *Hygrobiella laxifolia*, *Lophozia heterocolpos*, *Riccardia chamedryfolia*, *Barbula crocea*, *Bryum gemmiferum*, *Buxbaumia viridis*, *Campylophyllum calcareum*, *Conardia compacta*, *Dicranum viride*, *Didymodon cordatus*, *Didymodon glaucus*, *Drepanocladus sendtneri*, *Helodium blandowii*, *Isopterygiopsis muelleriana*, *Isopterygiopsis pulchella*, *Mnium lycopodioides*, *Orthotrichum patens*, *Orthotrichum stellatum*, *Paludella squarrosa*, *Physcomitrella patens*, *Plagiopus oederianus*, *Plagiothecium latebricola*, *Pohlia ludwigii*, *Rhynchostegium confertum*, *Scorpidium revolvens*, *Serpolleskea subtilis*, *Zygodon dentatus*]
- Kučera J., Müller F. & Plášek V. (2005): Mechiorsty zaznamenané v průběhu 12. jarního setkání Bryologicko-lichenologické Sekce v CHKO Křivoklátsko [Bryophytes recorded during the 12th Spring Meeting of the Bryological and Lichenological Section in the Křivoklátsko region (Central Bohemia)]. – Bryonora 35: 21–31.
- Kučera J. & Váňa J. (2006) ‘2005’: Seznam a červený seznam mechiorstů České republiky (2005) [Check-and Red list of bryophytes of the Czech Republic 2005]. – Příroda 23: 1–104.

- Loskotová E. (2004): Mechiorsty Povydří a dolního toku Křemelné. – In: Dvořák L. & Šustr P. (eds.), Aktuality Šumavského Výzkumu II, Sborník z konference, Srní 4.-7. října 2004, p. 121–123, Správa Národního parku a CHKO Šumava, Vimperk.
- Loskotová E. (2005): Bryoflóra vybraných území CHKO Blaník [Bryoflora of selected areas in the Protected Landscape Area Blaník (Central Bohemia)]. – Bryonora 35: 17–21.
- Loskotová E. (2006): Interesting records of *Aneura maxima* (Schiffn.) Steph. (*Metzgeriales*) in the Czech Republic and Slovakia. – Silva Gabreta 12: 15–18.
- Müller F. & Rätzel S. (2005): The bryophyte flora of the sandstone region Hradčanské stěny near Doksy in Central North Bohemia. – Bryonora 36: 1–8.
- Navrátilová J. & Navrátil J. (2004): Rašeliniště vegetace v severní části třeboňsko-jindřichohradeckého pomezí. – Sborník Jihočeského Muzea v Českých Budějovicích, Přírodní Vědy, 44: 45–58.
- Navrátilová J. & Navrátil J. (2005): Stanoviště nároky některých ohrožených a vzácných rostlin rašelinišť Třeboňska [Environmental factors of some endangered and rare plants in Třeboň's mires]. – Zprávy České Botanické Společnosti 40: 279–299.
- Němcová L. (2005): Moosgewächse des Naturschutzgebietes Březina im Böhmischem Mittelgebirge. – Severočeskou Přírodou (Litoměřice) 36–37: 123–126.
- Novotný I. & Kubešová S. (2004): Mechiorsty známé neznámé. – Moravské zemské muzeum, Brno. [42 pp.]
- Peksa O. & Loskotová E. (2005): Lišeňíky a mechiorsty na sutích. – Šumava 10: 14–16.
- Plášek V. (2004): Mechiorsty Hlučínska. – In: Koutecká V. a kol., Příroda Hlučínska, p. 19–20 & 31–36, Město Hlučín & Sdružení obcí Hlučínska.
- Plášek V. (2005): Mechiorsty Přírodní památky Prameny Zrzávky (Hostašovice, Moravskoslezský kraj). – Práce a Studie Muzea Beskyd, Přírodní Vědy, 15: 157–162.
- Plášek V. (2005): Mechiorsty. – In: Ševčík J., Pečinka A. & Plášek V. (eds.), Rostliny Národní přírodní památky Landek, p. 4–9, Biologicko-ekologický klub, Ostrava.
- Plášek V. (2005): Mechiorsty NPP Rešovské vodopády (Nízký Jeseník). – Časopis Slezského Zemského Muzea, Ser. A, 54: 1–10.
- Plášek V. (2005): Základy bryologie (Systém, fylogeneze a ekologie mechiorstů). – Ostravská univerzita v Ostravě, Ostrava. [Skriptum, 79 pp.]
- Plášek V., Stebel A. & Wierzcholska S. (eds.) (2005): Mechiorsty zaznamenané v průběhu 18. podzimního setkání Bryologicko-lichenologické sekce v Beskydech [Bryophytes recorded during the 18<sup>th</sup> Autumnal Meeting of the Bryological and Lichenological Section in Beskydy Mts. (northern Moravia & Silesia)]. – Bryonora 36: 14–24.
- Plášek V. & Wierzcholska S. (2005): Mechiorsty NPR Pulčín – Hradisko (Javorníky, Česká republika). – Časopis Slezského Zemského Muzea, Ser. A, 54: 215–222.
- Poulíčková A., Hájek M. & Rybníček K. (eds.) (2005): Ecology and palaeoecology of spring fens of the West Carpathians. – Palacký University, Olomouc. [209 pp.]
- Soldán Z. (2005): Česká a slovenská bryologická bibliografie XVII. – Bryonora 35: 48–51.
- Suchara I. & Sucharová J. (2005): Mechový archiv. – Vesmír 84: 598–603.
- Šoltés R. (2004): Zaniknuté lokality glaciálnych reliktov machorastov v rašeliniskách Slovenska [Old localities of glacial relic moss species in peatbogs of Slovakia]. – Naturae Tutela 8: 37–46.
- Šoltés R. & Mačáková M. (2005): The second record of *Oreas martiana* (*Bryophyta*) in the Carpathians (the Tatra Mts., Slovakia). – Thaiszia – Journal of Botany 15: 31–34.
- Váňa J. (2006): Obecná bryologie. – Karolinum, Praha. [Skriptum PřF UK Praha, 188 pp.]
- Váňa J., Zmrhalová M., Kučera J., Plášek V. & Shaw B. (2005): Vzácné a ohrožené druhy mechiorstů sudetských karů v ČR. – In: Campanula – sborník referátů z konference k 35. výročí Chráněné krajinné oblasti Jeseníky, Karlova Studánka a Malá Morávka 11.–12.11.2004, p. 113–114, SOP – Správa CHKO Jeseníky, Jeseník.
- Zmrhalová M., Kučera J., Plášek V., Váňa J. & Shaw B. (2005): Metodika bryologického výzkumu Vysokých Sudet. – In: Campanula – sborník referátů z konference k 35. výročí Chráněné krajinné oblasti Jeseníky, Karlova Studánka a Malá Morávka 11.–12.11.2004, p. 111–112, SOP – Správa CHKO Jeseníky, Jeseník.

## 2. Recenze [Reviews]

Kučera J. (2005): [Frahm J.-P. & Frey W. (2004): Moosflora, 4. Auflage]. – Bryonora 35: 44–45.

- Kučera J. (2005): [Ignatov M. S. & Ignatova E. A. (2004): Flora mchov sredněj časti evropského Rossii. Tom 2. *Fontinalaceae – Amblystegiaceae*.] – Bryonora 35: 42–44.
- Kučera J. (2005): [Müller F. (2004): Verbreitungsatlas der Moose Sachsen]. – Bryonora 35: 46–47.
- Kučera J. (2005): [Pistarino A. & al. (2005), Briosite del Piemonte: la collezione della Val Sangone]. – Bryonora 36: 40–41.
- Kučera J. (2005): [Smith A. J. E. (2004): The Moss Flora of Britain and Ireland, 2nd ed.]. – Bryonora 35: 45–46.
- Pohlová R. (2005): [Novotný I. & Kubešová S. (2004): Mechiorosty známé neznámé]. – Bryonora 36: 42.
- Váňa J. (2004): [Ignatov M. S. & Ignatova E. A. (2003): Flora mchov sredněj časti evropského Rossii. Tom 1. *Sphagnaceae – Hedwigiaceae*.] – Bryonora 34: 36–38.
- Zmrhalová M. (2005): [Plášek V. (2005): Základy bryologie]. – Bryonora 36: 41.

### 3. Bryologické rukopisné práce [Bryological manuscripts]

- Berka T. (2006): Mechiorosty Brtnické vrchoviny. – Ms., 113 pp. [Mgr. thesis, Palacký University, Olomouc]
- Buryová B. (2004): Morphological and genetic variation in selected species of the genus *Philonotis* (Bartramiaceae, Bryopsida). - Ms., 101 pp. [Ph.D. thesis, Charles University, Prague]
- Hájková P. (2004): Ekologie a vývoj prameništních slatiníšť v západní části Karpat. – Ms., 164 pp. [Ph.D. thesis, Masaryk University, Brno]
- Holá E. (2006): Bryoflóra horního toku Křemelné na Šumavě. – Ms., 50 pp. [Bc. thesis, University of South Bohemia, České Budějovice]
- Horodyská E. (2006): Minerotrofní rašeliniště Českomoravské vrchoviny: srovnání historických a současných dat. – Ms., 47 pp. [Mgr. thesis, Masaryk University, Brno]
- Jašík M. (2006): Vliv přítomnosti mechového patra na složení taxocenáz epigeických brouků. – Ms., 65 pp. [Bc. thesis, University of Ostrava]
- Koloničná M. (2005): Diverzifikace nik slatiných zástupců rodu *Drepanocladus* s. l. v Západních Karpatech: *Scorpidium cossonii*, *S. revolvens*, *Hamatocaulis vernicosus*. – Ms., 53 pp. [Mgr. thesis, Masaryk University, Brno]
- Košnar J. (2004): Bryofloristicko-ekologická studie Přírodní rezervace Maštale. – Ms., 67 pp. [Bc. thesis, University of South Bohemia, České Budějovice]
- Králová Š. (2005): Vegetační a stanovištní studie NPR Rašeliniště Jizery. – Ms., 71 pp. [Mgr. thesis, Masaryk University, Brno]
- Kubešová S. (2004): Materiály k bryoflóře Moravy. – Ms., 57 pp. [RNDr. thesis, Masaryk University, Brno]
- Loskotová E. (2004): Biodiverzita a ekologie mechiorostů ve vybraných I. zónách NP Šumava. – Ms., 127 pp. [Mgr. thesis, Charles University, Prague]
- Motyka O. (2006): Ekologická studie vybraných druhů epifytních mechiorostů v pobřežní vegetaci meandrů řeky Lučiny. – Ms., 51 pp. [Bc. thesis, University of Ostrava]
- Musil Z. (2005): Mechiorosty nivy spojeného toku Orlice na území přírodního parku Orlice. – Ms. [Mgr. thesis, Palacký University, Olomouc]
- Navrátilová J. (2006): Ekologie vegetace rašeliných okrajů rybníků Třeboňské pánve. – Ms. [Ph.D. thesis, Masaryk University, Brno]
- Rozbrojová Z. (2005): Indikace stanovištních podmínek prameništních rašelinišť pomocí chemického složení rostlin. – Ms., 77 pp. [Mgr. thesis, Masaryk University, Brno]
- Štechová T. (2005): Ekologická studie druhu *Hamatocaulis vernicosus* (*Amblystegiaceae*, *Bryopsida*) a návrh managementu na jeho lokalitách. – Ms., 51 pp. [Mgr. thesis, University of South Bohemia, České Budějovice]

### 4. Nebryologické práce s údaji o výskytu mechiorostů na území ČR a SR [Non-bryological publications with bryophyte records from the territory of CR and SR]

- Dimopoulos P., Sýkora K., Gilissen C., Wiecherink D & Georgiadis T. (2005): Vegetation Ecology of Kalodiki fen (NW Greece). – Biologia, Sect. Botany, 60: 69–82.
- Dítě D. (ed.) (2004): Zaujmavejší floristické nálezy. – Bulletin Slovenskej Botanickej Spoločnosti 26: 219–227. [*Helodium blandowii*, *Meesia triquetra*, *Paludella squarrosa*]

- Hrvnák R. & Ujházy K. (2005): Changes of the mountain grassland vegetation after abandonment and colonization by norway spruce. – *Ekológia* (Bratislava) 24: 231–253.
- Kanka R., Kollár J. & Barančok P. (2005): Monitoring of climatic change impacts on alpine vegetation in the Tatry Mts – First approach. – *Ekológia* (Bratislava) 24: 411–418.
- Kliment J. (2005): *Bellidiastro michelii-Seslerietum calcariae* – spoločenstvo zväzu *Astero-Seslerion calcariae* alebo *Cystopteridion fragilis?* [*Bellidiastro michelii-Seslerietum calcariae* – community of the alliance *Astero-Seslerion calcariae* or *Cystopteridion fragilis?*]. – *Bulletin Slovenskej Botanickej Spoločnosti* 27: 171–180.
- Kliment J., Jarolímek I., Šibík J. & Valachovič M. (2004): Syntaxonomy and nomenclature of the communities of the orders *Calamagrostietalia villosae* and *Adenostyletalia* in Slovakia. – *Thaiszia – Journal of Botany* 14: 93–157.
- Kliment J., Bernátová D., Jarolímek I. & Uhlířová J. (2005): Floristic composition and syntaxonomy of the plant communities with *Carex sempervirens* subs. *tatrorum* in the West Carpathians. – *Biologia, Section Botany*, 60: 37–56.
- Lepší M., Lepší P. & Štech M. (eds.) (2005): Výsledky floristického kurzu ČBS v Českých Budějovicích 2001 (1.-7. července 2001). – Zprávy České Botanické Společnosti 40, Příloha 2: 71–137.
- Malovcová-Staníková M. (2004): Communities of the suballiance *Caricenion gracilis* in the Borská nížina lowland. – *Zborník Slovenského Národného Muzea, Prírodné Vedy*, 53: 283–287.
- Melichar V. (2005): Poznámky o floristicky zajímavé lokalitě Jezero Odeř v Krušných horách. – *Calluna* 10: 13–15.
- Michálková D. (2005): Flóra vrchu Rohatín v Strážovských vrchoch [Flora of Mt. Rohatín in the Strážovské vrchy Mts]. – *Bulletin Slovenskej Botanickej Spoločnosti* 27: 121–129.
- Mikuška B. (2005): Syntaxonómia dubovo-borovicových kultúrnych lesov na Borskej nížine [Syntaxonomy of the cultural oak-pine forests in the Borská nížina Lowland]. – *Bulletin Slovenskej Botanickej Spoločnosti* 27: 157–169.
- Navrátilová J. & Navrátil J. (2005): Vegetation gradients in fishpond mires in relation to seasonal fluctuations in environmental factors. – *Preslia* 77: 405–418.
- Navrátilová J. & Navrátil J. (2005): Hlavní typy nelesní rašeliniskové vegetace Třeboňské pánve. – *Sborník Jihočeského Muzea v Českých Budějovicích, Přírodní Vědy*, 45: 45–56.
- Neuhäuslová Z. & Sofron J. (2005): Montane Fichten-(Tannen)-Buchenwälder (*Calamagrostio villosae-Fagetum*) im Böhmerwald. – *Sborník Západočeského Muzea v Plzni, Příroda*, 42: 1–19.
- Šibík J., Petrík A., Krajčiová-Šibíková I. & Dúbravcová Z. (2005): Asociácia *Dryado octopetalae-Caricetum firmae* Sillinger 1933 v Západných Karpatoch [The association *Dryado octopetalae-Caricetum firmae* Sillinger 1933 in the Western Carpathians]. – *Bulletin Slovenskej Botanickej Spoločnosti* 27: 121–129.

## NOVÁ LICHENOLOGICKÁ LITERATURA XV.

### New lichenological literature, XV

Zdeněk Palice

*Botanický ústav AV ČR, CZ-252 43 Průhonice, e-mail: palice@ibot.cas.cz*

- Aguirre-Hudson B., Farkas E. & Lőkös L. (2005): New records of *Leptorhaphis* and other ascomycete genera from the Carpathian basin. – *Herzogia* 17: 47–50.
- Ahti T. & DePriest P. (2005): Proposals to conserve eight names and reject one species name in *Cladoniaceae (Fungi)*. – *Taxon* 54: 183–188.
- Ahti T. & Hawksworth D. L. (2005): *Xanthoparmelia stenophylla*, the correct name for *X. somloënsis*, one of the most widespread usnic acid containing species of the genus. – *Lichenologist* 37: 363–366.
- Alexeeva N. (2005): Lichens from islands of the Russian part of the Gulf of Finland. – *Folia Cryptogamica Estonica* 41: 5–12.