

- Hrvnák R. & Ujházy K. (2005): Changes of the mountain grassland vegetation after abandonment and colonization by norway spruce. – *Ekológia* (Bratislava) 24: 231–253.
- Kanka R., Kollár J. & Barančok P. (2005): Monitoring of climatic change impacts on alpine vegetation in the Tatry Mts – First approach. – *Ekológia* (Bratislava) 24: 411–418.
- Kliment J. (2005): *Bellidiastro michelii-Seslerietum calcariae* – spoločenstvo zväzu *Astero-Seslerion calcariae* alebo *Cystopteridion fragilis?* [*Bellidiastro michelii-Seslerietum calcariae* – community of the alliance *Astero-Seslerion calcariae* or *Cystopteridion fragilis?*]. – *Bulletin Slovenskej Botanickej Spoločnosti* 27: 171–180.
- Kliment J., Jarolímek I., Šibík J. & Valachovič M. (2004): Syntaxonomy and nomenclature of the communities of the orders *Calamagrostietalia villosae* and *Adenostyletalia* in Slovakia. – *Thaiszia – Journal of Botany* 14: 93–157.
- Kliment J., Bernátová D., Jarolímek I. & Uhlířová J. (2005): Floristic composition and syntaxonomy of the plant communities with *Carex sempervirens* subs. *tatrorum* in the West Carpathians. – *Biologia, Section Botany*, 60: 37–56.
- Lepší M., Lepší P. & Štech M. (eds.) (2005): Výsledky floristického kurzu ČBS v Českých Budějovicích 2001 (1.-7. července 2001). – Zprávy České Botanické Společnosti 40, Příloha 2: 71–137.
- Malovcová-Staníková M. (2004): Communities of the suballiance *Caricenion gracilis* in the Borská nížina lowland. – *Zborník Slovenského Národného Muzea, Prírodné Vedy*, 53: 283–287.
- Melichar V. (2005): Poznámky o floristicky zajímavé lokalitě Jezero Odeř v Krušných horách. – *Calluna* 10: 13–15.
- Michálková D. (2005): Flóra vrchu Rohatín v Strážovských vrchoch [Flora of Mt. Rohatín in the Strážovské vrchy Mts]. – *Bulletin Slovenskej Botanickej Spoločnosti* 27: 121–129.
- Mikuška B. (2005): Syntaxonómia dubovo-borovicových kultúrnych lesov na Borskej nížine [Syntaxonomy of the cultural oak-pine forests in the Borská nížina Lowland]. – *Bulletin Slovenskej Botanickej Spoločnosti* 27: 157–169.
- Navrátilová J. & Navrátil J. (2005): Vegetation gradients in fishpond mires in relation to seasonal fluctuations in environmental factors. – *Preslia* 77: 405–418.
- Navrátilová J. & Navrátil J. (2005): Hlavní typy nelesní rašeliniskové vegetace Třeboňské pánve. – *Sborník Jihočeského Muzea v Českých Budějovicích, Přírodní Vědy*, 45: 45–56.
- Neuhäuslová Z. & Sofron J. (2005): Montane Fichten-(Tannen)-Buchenwälder (*Calamagrostio villosae-Fagetum*) im Böhmerwald. – *Sborník Západočeského Muzea v Plzni, Příroda*, 42: 1–19.
- Šibík J., Petrík A., Krajčiová-Šibíková I. & Dúbravcová Z. (2005): Asociácia *Dryado octopetalae-Caricetum firmae* Sillinger 1933 v Západných Karpatoch [The association *Dryado octopetalae-Caricetum firmae* Sillinger 1933 in the Western Carpathians]. – *Bulletin Slovenskej Botanickej Spoločnosti* 27: 121–129.

NOVÁ LICHENOLOGICKÁ LITERATURA XV.

New lichenological literature, XV

Zdeněk Palice

Botanický ústav AV ČR, CZ-252 43 Průhonice, e-mail: palice@ibot.cas.cz

- Aguirre-Hudson B., Farkas E. & Lőkös L. (2005): New records of *Leptorhaphis* and other ascomycete genera from the Carpathian basin. – *Herzogia* 17: 47–50.
- Ahti T. & DePriest P. (2005): Proposals to conserve eight names and reject one species name in *Cladoniaceae (Fungi)*. – *Taxon* 54: 183–188.
- Ahti T. & Hawksworth D. L. (2005): *Xanthoparmelia stenophylla*, the correct name for *X. somloënsis*, one of the most widespread usnic acid containing species of the genus. – *Lichenologist* 37: 363–366.
- Alexeeva N. (2005): Lichens from islands of the Russian part of the Gulf of Finland. – *Folia Cryptogamica Estonica* 41: 5–12.

- Altermann S. (2004): A second look at *Letharia* (Th. Fr.) Zahlbr. – Bulletin of the California Lichen Society 11: 33–36.
- Andersen H. L. & Ekman S. (2005): Disintegration of the *Micareaceae* (lichenized *Ascomycota*): a molecular phylogeny based on mitochondrial rDNA sequences. – Mycological Research 109: 21–30.
- Aragón G., Otálora M. A. G. & Martínez I. (2005): New data on the genus *Leptogium* (lichenized ascomycetes) in the Iberian Peninsula. – Nova Hedwigia 80: 199–226.
- Areskoug V. & Thor G. (2005): Distribution, status and ecology of the lichen *Cyphelium notarisii* in Sweden. – Annales Botanici Fennici 42: 317–326.
- Armstrong R. A. (2005): Growth curves of four crustose lichens. – Symbiosis 38: 47–57.
- Arup U. (2004): Three overlooked *Lecidea* species in Sweden. – Symbolae Botanicae Upsalienses 34: 39–48.
- Arup U. & Westberg M. (2005): *Candelariella medians* new to Sweden. – Graphis Scripta 17: 1–2.
- Bartók K. & Lökös L. (2004): Contributions to the lichen flora and lichen ecology in the Parâng and Lotru Mountains (southern Carpathians). – Cotribuții Botanice 39: 41–49.
- Beckett R. B., Minibayeva F. V. & Laufer Z. (2005): Extracellular reactive oxygen species production by lichens. – Lichenologist 37: 397–407.
- Belnap J. & Lange O. L. (2005): Lichens and microfungi in biological soil crusts: Community structure, physiology and ecological functions. – In: Dighton J. (ed.), The Fungal Community: Its Organization and Role in the Ecosystem Vol. 23, p. 397–712, Taylor & Francis Group, Boca Raton, Florida.
- Berger F. & Priemetzhofer F. (2005): Neue und bemerkenswerte Funde von Flechten aus Oberösterreich, Österreich. – Beiträge zur Naturkunde Oberösterreichs 14: 3–18.
- Bergamini A., Scheidegger C., Stofer S., Carvalho P., Davey S., Dietrich M., Dubs F., Farkas E., Groner U., Kärkkäinen K., Keller K., Lökös L., Lommi S., Máguas C., Mitchell R., Pinho P., Rico V. J., Aragón G., Truscott A.-M., Wolseley P. & Watt A. (2005): Performance of macrolichens and lichen genera as indicators of lichen species richness and composition. – Conservation Biology 19: 1051–1062.
- Berglund H. & Jonsson B. G. (2005): Verifying an extinction debt among lichens and fungi in northern Swedish boreal forests. – Conservation Biology 19: 338–348.
- Bjelland T. & Ekman S. (2005): Fungal diversity in rock beneath a crustose lichen as revealed by molecular markers. – Microbial Ecology 49: 598–603.
- Bjerke J. W. (2005): Synopsis of the lichen genus *Menegazzia* (*Parmeliaceae*, *Ascomycota*) in South America. – Mycotaxon 91: 423–454.
- Bjerke J. W., Elvebakk A., Domínguez E. & Dahlback A. (2005): Seasonal trends in usnic acid concentrations of Arctic, alpine and Patagonian populations of the lichen *Flavocetraria nivalis*. – Phytochemistry 66: 337–344.
- Blanco O., Crespo A. & Elix J. A. (2005): Two new species of *Xanthoparmelia* (*Ascomycota*: *Parmeliaceae*) from Spain. – Lichenologist 37: 97–100.
- Blanco O., Crespo A., Divakar P. K., Elix J. A. & Lumbsch H. T. (2005): Molecular phylogeny of parmotremoid lichens (*Ascomycota*, *Parmeliaceae*). – Mycologia 97: 150–159.
- Blinkova O. & Urbanavichus G. (2005): Ecological analysis of lichens in the Teberda State Biosphere Reserve (North-Western Caucasus, Russia). – Folia Cryptogamica Estonica 41: 23–34.
- Boustie J. & Grube M. (2005): Lichens—a promising source of bioactive secondary metabolites. – Plant Genetic Resources: characterization and utilization 3: 273–287.
- Breuss O. & John V. (2004): New and interesting records of lichens from Turkey. – Österreichische Zeitschrift für Pilzkunde 13: 281–294.
- Breuss O. & Türk R. (2004): *Involucropyrenium pusillum* (*Verrucariaceae*) – eine neue Flechtenart aus Oberösterreich. – Beiträge zur Naturkunde Oberösterreichs 13: 213–216.
- Brodo I. M. & Aptroot A. (2005): Corticolous species of *Protoparmelia* (lichenized *Ascomycotina*) in North America. – Canadian Journal of Botany 83: 1075–1081.
- Burgisser L., Habashi-Mayor C., Clerc P. & Price M. (2004): Inventaire des lichens, des mousses et des hépatiques du Bois de la Grille (commune de Vernier, canton de Genève). – Saussurea 34: 111–129.
- Carbonero E. R., Mellinger C. G., Eliasaro S., Gorin P. A. J. & Iacomini M. (2005): Chemotypes significance of lichenized fungi by structural characterization of heteropolysaccharides from the genera *Parmotrema* and *Rimelia*. – FEMS Microbiology Letters 246: 273–278.
- Carbonero E. R., Montai A. V., Mellinger C. G., Eliasaro S., Sasaki G. L., Gorin P. A. J. & Iacomini M. (2005): Glucans of lichenized fungi: significance for taxonomy of the genera *Parmotrema* and *Rimelia*. – Phytochemistry 66: 929–934.

- Castello M. & Skert N. (2005): Evaluation of lichen diversity as an indicator of environmental quality in the North Adriatic submediterranean region. – *Science of the Total Environment* 336: 201–214.
- Cezanne R. & Eichler M. (2004): Neu- und Wiederfunde von Flechten und flechtenbewohnenden Pilzen in Hessen. – *Hessische Floristische Briefe* 53: 43–52.
- Cezanne R., Eichler M. & Wirth V. (2004): Flechten-Exkursion in die Umgebung von Bad Wildungen 04./05. 10. 2002. – *Hessische Floristische Briefe* 53: 17–28.
- Chen J., Zhang M. Y., Wang L., Shimazaki H. & Tamura M. (2005): A new index for mapping lichen-dominated biological soil crusts in desert areas. – *Remote Sensing of Environment* 96: 165–175.
- Chilton R. & Orvos D. (2004): Bioaccumulation of mercury in lichen (*Hypogymnia physodes*) near a coal-fired power plant. – *Southeastern Biology* 51: 151–152.
- Czarnota P. (2004): New and some rare species of the genus *Micarea* (*Micareaceae*) in the lichen flora of Poland. – *Polish Botanical Journal* 49: 135–143.
- De los Ríos A., Sancho L. G., Grube M., Wierzchos J. & Ascaso C. (2005): Endolithic growth of two *Lecidea* lichens in granite from continental Antarctica detected by molecular and microscopy techniques. – *New Phytologist* 165: 181–190.
- De los Ríos A., Wierzchos J., Sancho L. G., Green T. G. A. & Ascaso C. (2005): Ecology of endolithic lichens colonizing granite in continental Antarctica. – *Lichenologist* 37: 383–395.
- Diederich P., Van den Broeck D., Ertz D., Heylen O., Jordaeans D. & Sérusiaux E. (2004): Report on two lichenological field meetings in the province of Luxembourg in Belgium. – *Bulletin de la Société des Naturalistes Luxembourgeois* 105: 57–64.
- Divakar P. K., Blanco O., Hawksworth D. L. & Crespo A. (2005): Molecular phylogenetic studies on the *Parmotrema reticulatum* (syn. *Rimelia reticulata*) complex, including the confirmation of *P. pseudoreticulatum*. – *Lichenologist* 37: 55–65.
- Divakar P. K., Molina M. C., Lumbsch H. T. & Crespo A. (2005): *Parmelia barrenoae*, a new lichen species related to *Parmelia sulcata* (*Parmeliaceae*) based on molecular and morphological data. – *Lichenologist* 37: 37–46.
- Dobson F. S. (2005): Lichens: An illustrated guide to the British and Irish species. 5th ed. – The Richmond Publishing Co., Slough. [480 pp.]
- Ellis C. J., Crittenden P. D., Scrimgeour C. M. & Ashcroft C. J. (2005): Translocation of ^{15}N indicates nitrogen recycling in the mat-forming lichen *Cladonia portentosa*. – *New Phytologist* 168: 423–434.
- Emmerer B. & Hafellner J. (2005): Zur aktuellen Vegetation auf Abraum- und Schlackenhalden historischer Kupferbergbaue in der Montanstufe der Niederen Tauern und der Eisenerzer Alpen (Steiermark, Österreich). – *Mitteilungen der Naturwissenschaftlichen Vereines für Steiermark* 134: 121–152.
- Ertz D., Christnach C., Wedin M. & Diederich P. (2005): A world monograph of the genus *Plectocarpon* (*Roccellaceae, Arthoniales*). – *Bibliotheca Lichenologica* 91: 1–155.
- Favero-Longo S. E., Castelli D., Salvadori O., Belluso E. & Piervittori R. (2005): Pedogenetic action of the lichens *Lecidea atrobrunnea*, *Rhizocarpon geographicum* gr. and *Sporastatia testudinea* on serpentized ultramafic rocks in an alpine environment. – *International Biodeterioration & Biodegradation* 56: 17–27.
- Fischer E. & Killmann D. (2004): Diversität der Flechtenflora in Westerwald, Lahntal und angrenzenden Gebieten. – *Zeitschrift für Naturschutz* 29: 1–145.
- Flakus A. (2005): *Gyalectea peziza* (*Gyalectaceae*, lichenized *Ascomycota*), first record in the Polish Tatry Mts. – *Biologia* 60: 373–375.
- Friedman E. I. & Sun H. J. (2005): Communities adjust their temperature optima by shifting producer-to-consumer ratio, shown in lichens as models: I. Hypothesis. – *Microbial Ecology* 49: 523–527.
- Fryday A. M. (2005): The genus *Porpidia* in northern and western Europe, with special emphasis on collections from the British Isles. – *Lichenologist* 37: 1–35.
- Gaio-Oliveira G., Dahlman L., Palmqvist K. & Mágua C. (2005): Responses of the lichen *Xanthoria parietina* (L.) Th. Fr. to varying thallus nitrogen concentrations. – *Lichenologist* 37: 171–179.
- Gaio-Oliveira G., Dahlman L., Palmqvist K., Martins-Loução M. A. & Mágua C. (2005): Nitrogen uptake in relation to excess supply and its effects on the lichens *Evernia prunastri* (L.) Ach and *Xanthoria parietina* (L.) Th. Fr. – *Planta* 220: 794–803.
- Garcia-Vallès M., Topal T. & Vendrell-Saz M. (2003): Lichenic growth as a factor in the physical deterioration or protection of Cappadocian monuments. – *Environmental Geology* 43: 776–781.
- Gauslaa Y. (2005): Lichen palatability depends on investments in herbivore defence. – *Oecologia* 143: 94–105.

- Gauslaa Y. & McEvoy M. (2005): Seasonal changes in solar radiation drive acclimation of the sun-screening compound parietin in the lichen *Xanthoria parietina*. – Basic and Applied Ecology 6: 75–82.
- Geiser L. H., Glavich D. A., Mikulin A. G., Ingersoll A. R. & Hutten M. (2004): New records of rare and unusual coastal lichens from the US Pacific Northwest. – Evansia 21: 104–110.
- Gilbert O. L. (2004): The Lichen hunters. – The Book Guild Ltd., Lewes, East Sussex. [208 pp.]
- Giordano S., Adamo P., Sorbo S. & Vingiani S. (2005): Atmospheric trace metal pollution in the Naples urban area based on results from moss and lichen bags. – Atmospheric Pollution 136: 431–442.
- Glavich D. A., Geiser L. H. & Mikulin A. G. (2005): Rare epiphytic coastal lichen habitats, modeling, and management in the Pacific Northwest. – Bryologist 108: 377–390.
- Goward T. & Spribille T. (2005): Lichenological evidence for the recognition of inland rain forests in western North America. – Journal of Biogeography 32: 1209–1219.
- Grube M. (2005): *Frigidopyrenia* – a new genus for a peculiar subarctic lichen, with notes on similar taxa. – Phyton 45: 305–318.
- Günzl B. & Fischer P. (2004): Ein Beitrag zur Verbreitung der Chemotypen ausgewählter *Cladonia*-Arten. – Aktuelle Lichenologische Mitteilungen, NF 13: 20–25.
- Guo S. & Kashiwadani H. (2004): Recent study on the phylogeny of the genus *Cladonia* (s. lat.) with the emphasis on the integrative biology. – In: Akiyama S., Miyawaki R., Kubodera T. & Higuchi M. (eds.), Proceedings of the 5th and 6th Symposia on Collection Building and Natural History Studies in Asia and the Pacific Rim, p. 207–225, National Science Museum, Tokyo.
- Hafellner J., Obermayer S. & Obermayer W. (2005): Zur Diversität der Flechten und lichenicolen Pilze im Hochschwab-Massiv (Nordalpen, Steiermark). – Mitteilungen der Naturwissenschaftlichen Vereines für Steiermark 134: 57–103.
- Hafellner J., Petutschnig W., Taurer-Zeiner C. & Mayrhofer H. (2005): Zur Flechtendiversität in den Gurktaler Alpen (Österreich: Kärnten, Steiermark und Salzburg). – Herzogia 17: 79–138.
- Hasse T. (2005): Charakterisierung der Sukzessionsstadien im *Spergulo-Corynephoretum* (Silbergrasfluren) unter besonderer Berücksichtigung der Flechten. – Tuexenia 25: 407–424.
- Hauck M. (2005): Epiphytic lichen diversity on dead and dying conifers under different levels of atmospheric pollution. – Environmental Pollution 135: 111–119.
- Hauck M. & Paul A. (2005): Manganese as a site factor for epiphytic lichens. – Lichenologist 37: 409–423.
- Hauck M. & Spribille T. (2005): The significance of precipitation and substrate chemistry for epiphytic lichen diversity in spruce-fir forests of the Salish Mountains, northwestern Montana. – Flora 200: 547–562.
- Hermansson J. & Pystina T. (2004): Calicioid lichens and fungi in the Komi Republic, Russia. – Symbolae Botanicae Upsalienses 34: 97–105.
- Holt E. A. & Severns P. M. (2005): The effects of prescribed burning on wet prairie lichen communities. – Natural Areas Journal 25: 130–136.
- Hoppert M., Flies C., Günzl B. & Schneider J. (2004): Colonization strategies of lithobiontic microorganisms on carbonate rocks. – Environmental Geology 46: 421–428.
- Hur J.-S., Harada H., Lim K.-M., Lee S. M., Vuong L. V., Oh S.-O. & Koh Y. J. (2004): Stimulation of soil microbial activity by lichen colonization in an abandoned coalmine waste dump soil. – Lichenology 3: 1–9.
- Isocrono D., Matteucci E., Faletti C. & Piervittori R. (2004): Contributo alle conoscenze licheniche nelle Alpi Pennine (Valsesia, Piemonte-Italia). – Bollettino del Museo Regionale di Scienze Naturali – Torino 21: 261–474.
- Johansson S., Søchting U., Elix J. A. & Wardlaw J. H. (2005): Chemical variation in the lichen genus *Letrovitiella* (*Ascomycota, Letrovitiaceae*). – Mycological Progress 4: 139–148.
- Jørgensen P. M. (2005): A new Atlantic species in *Fuscopannaria*, with a key to its European species. – Lichenologist 37: 221–225.
- Jürjado I., Lõhmus P., Nilson E., Randlane T., Saag A., Saag L. & Suija A. (2004): Eesti Pisisamblikud [Estonian Microlichens]. – Tartu University Press, Tartu. [583 pp.]
- Kalb K., Staiger B. & Elix J. A. (2004): A monograph of the lichen genus *Diorygma* – a first attempt. – Symbolae Botanicae Upsalienses 34: 133–181.
- Kalwij J. M., Wagner H. H. & Scheidegger C. (2005): Effects of stand-level disturbances on the spatial distribution of a lichen indicator. – Ecological Applications 15: 2015–2024.
- Kanz B., Dürhammer O. & Printzen C. (2005): Lichens and lichenicolous fungi of the Bavarian Forest. – Preslia 77: 355–403.

- Kauff F. & Büdel B. (2005): Ascoma ontogeny and apothecial anatomy in the *Gyalectaceae* (*Ostropales*, *Ascomycota*) support the re-establishment of the *Coenogoniaceae*. – *Bryologist* 108: 272–281.
- Keller C. (2005): Artificial substrata colonized by freshwater lichens. – *Lichenologist* 37: 357–362.
- Khodosovtsev A., Kuznetsova E. & Himelbrant D. (2004): Lichen genus *Caloplaca* on the Kamchatka Peninsula (Russian Far East). – *Botanica Lithuanica* 10: 195–208.
- Killmann D. (2004): Erstnachweis einer epiphyllen Flechtengesellschaft (*Fellhaneretum myrtillicolae* Spier & Aptroot) auf *Buxus sempervirens* in Deutschland. – *Decheniana* 157: 99–101.
- Kondratyuk S. Y. & Kärnefelt I. (2003): Revision of three natural groups of xanthorioid lichens (*Teloschistaceae*, *Ascomycota*). – *Ukrains'kyi Botanichnyi Zhurnal* 60: 427–437.
- Kotlova E. R. & Sinyutina N. F. (2005): Changes in the content of individual lipid classes of a lichen *Peltigera aphthosa* during dehydration and subsequent rehydration – *Russian Journal of Plant Physiology* 52: 35–42.
- Kranner I., Cram W. J., Zorn M., Wornik S., Yoshimura I., Stabentheiner E. & Pfeifhofer H. W. (2005): Antioxidants and photoprotection in a lichen as compared with its isolated symbiotic partners. – *Proceedings of the National Academy of Sciences of the United States of America* 102: 3141–3146.
- Kristmundsdóttir T., Jónsdóttir E., Ögmundsdóttir H. M. & Ingólfssdóttir K. (2005): Solubilization of poorly soluble lichen metabolites in non-toxic solvents for testing in tissue culture. – *European Journal of Pharmaceutical Sciences* 24: 539–543.
- Kukwa M. (2004): New or interesting records of lichenicolous fungi from Poland II. Species mainly from northern Poland. – *Herzogia* 17: 67–75.
- Kukwa M. (2004): Porosty z rodzaju *Lepraria* w Tatrzańskim Parku Narodowym. – *Parki Narodowe i Rezerwaty Przyrody* 23: 3–12.
- Lange O. L., Türk R. & Zimmermann D. G. (2005): Neufunde der boreal-montanen Flechte *Evernia divaricata* im trocken-warmen Main-Tauber-Gebiet und ihre Begleiter. – *Herzogia* 18: 51–62.
- Laundon J. R. (2005): The publication and typification of Sir James Edward Smith's lichens in English Botany. – *Botanical Journal of Linnean Society* 147: 483–499.
- Levia D. F. (2002): Nitrate sequestration by corticolous macrolichens during winter precipitation events. – *International Journal of Biometeorology* 46: 60–65.
- Lindblom L. & Ekman S. (2005): Molecular evidence supports the distinction between *Xanthoria parietina* and *X. aureola* (*Teloschistaceae*, lichenized *Ascomycota*). – *Mycological Research* 109: 187–199.
- Lindblom L., Ladstein M. H., Blom H. H., Ekman S. & Timdal E. (2005): *Xanthoria aureola* in Norway and a key to the species of *Xanthoria* s. str. in Scandinavia. – *Graphis Scripta* 17: 12–16.
- Litterski B. & Ahti T. (2004): World distribution of selected European *Cladonia* species. – *Symbolae Botanicae Upsalienses* 34: 205–236.
- Lücking R. & Bernecker-Lücking A. (2005): Drip-tips do not impair the development of epiphyllous rainforest lichen communities. – *Journal of Tropical Ecology* 21: 171–177.
- Lumbsch H. T., del Prado R. & Kantvilas G. (2005): *Gregorella*, a new genus to accommodate *Moelleropsis humida* and a molecular phylogeny of *Arctomiaceae*. – *Lichenologist* 37: 291–302.
- Marques A. P., Freitas M. C., Wolterbeek H. T., Steinebach O. M., Verburg T. & De Goeij J. J. M. (2005): Cell-membrane damage and element leaching in transplanted *Parmelia sulcata* lichen related to ambient SO₂, temperature, and precipitation. – *Environmental Science & Technology* 39: 2624–2630.
- Martínez I., Aragón G. & Burgaz A. R. (2004): Estudio de la flora líquenica epífita de la Sierra de Gúdar (Teruel). – *Teruel* 88: 99–120.
- Maslov A. I. & Pavlova E. A. (2005): Simple method for fractionation of *Parmelia sulcata* lichen thalli. – *Russian Journal of Plant Physiology* 52: 271–274.
- Masson D. (2005): Taxonomy, ecology and chorology of French species of *Hypotrachyna* and *Parmelinopsis* genera (lichenized *Ascomycota*, *Parmeliaceae*). – *Cryptogamie Mycologie* 26: 205–263.
- Mayrhofer H., Denchev C. M., Stoykov D. Y. & Nikolova S. O. (2005): Catalogue of the lichenized and lichenicolous fungi in Bulgaria. – *Mycologia Balcanica* 2: 3–61.
- Mayrhofer H. & Unger E. (2005): Ein Beitrag zur Diversität von lichenisierten und lichenicolen Pilzen im Gebiet des Teigitschtales (Steiermark, Österreich). – *Mitteilungen der Naturwissenschaftlichen Vereines für Steiermark* 134: 105–120.
- Mikhailova I., Trubina M., Vorobeichik E. & Scheidegger C. (2005): Influence of environmental factors on the local-scale distribution of cyanobacterial lichens: case study in the North Urals, Russia. – *Folia Cryptogamica Estonica* 41: 45–54.

- Minganti V., Capelli R., Drava G., De Pellegrini R., Brunialti G., Giordano P. & Modenesi P. (2003): Biomonitoring of Trace Metals by Different species of Lichens (*Parmelia*) in North-West Italy. – *Journal of Atmospheric Chemistry* 45: 219–229.
- Molnár K. (2004): Lichen types and list of exsiccata in the Eger Cryptogamic Herbarium (EGR). – *Folia Historico Naturalia Musei Matraensis* 28: 53–55.
- Monnet F., Bordas F., Deluchat V., Chatenet P., Botineau M. & Baudu M. (2005): Use of the aquatic lichen *Dermatocarpon luridum* as bioindicator of copper pollution: Accumulation and cellular distribution tests. – *Environmental Pollution* 138: 455–461.
- Motiejūnaitė J. (2005): Distribution of some rare and declining lichen species in lowland eastern and eastern-central Europe. – *Biologia* 60: 357–363.
- Motiejūnaitė J., Czyżewska K. & Cieśliński S. (2004): Lichens – indicators of old-growth forests in biocentres of Lithuania and North-East Poland. – *Botanica Lithuanica* 10: 59–74.
- Motiejūnaitė J. & Golubkov V. V. (2005): Cyanolichens of freshwater aquatic and subaquatic habitats in Lithuania and Belarus. – *Botanica Lithuanica* 11: 35–40.
- Motiejūnaitė J., Stončius D. & Kukwa M. (2005): Contribution to the Lithuanian flora of lichens and allied fungi. II. – *Botanica Lithuanica* 11: 41–49.
- Moutchnik E. E. & Zavarzin A. A. (2005): Lichens in the Red Data Books of Russia. – *Folia Cryptogamica Estonica* 41: 59–80.
- Mylllys L., Högnabba F., Lohtander K., Thell A., Stenroos S. & Hyvönen J. (2005): Phylogenetic relationships of *Stereocaulaceae* based on simultaneous analysis of beta-tubulin, GAPDH and SSU rDNA sequences. – *Taxon* 54: 605–618.
- Mylllys L., Lohtander K. & Stenroos S. (2003): Phylogenetic position of the asexual fungal genus *Lepraria* within *Ascomycota*. – *Cladistics* 19: 158.
- Nifontova M. G. (2005): Use of lichens and mosses for on-line testing of environmental radioactive contamination. – *Russian Journal of Plant Physiology* 52: 60–63.
- O'Brien H. E., Miadlikowska J. & Lutzoni F. (2005): Assessing host specialization in symbiotic cyanobacteria associated with four closely related species of the lichen fungus *Peltigera*. – *European Journal of Phycology* 40: 363–378.
- Öckinger E., Niklasson M. & Nilsson S. G. (2005): Is local distribution of the epiphytic lichen *Lobaria pulmonaria* limited by dispersal capacity or habitat quality? – *Biodiversity and Conservation* 14: 759–773.
- Omarsdóttir S., Freysdóttir J., Barsett H., Smestad Paulsen B. & Olafsdóttir E. S. (2005): Effects of lichen heteroglycans on proliferation and IL-10 secretion by rat spleen cells and IL-10 and TNF- α secretion by rat peritoneal macrophages in vitro. – *Phytomedicine* 12: 461–467.
- Otte V. (2003): Lichenologische Beobachtungen in der Oberlausitz. – *Abhandlungen und Berichte des Naturkundemuseums Görlitz* 75: 35–42.
- Otte V. (2004): Versuch der Identifikation der Moose und Flechten aus von Bergens „Flora Francofurtana“ (1750). – *Verhandlungen des Botanischen Vereins von Berlin und Brandenburg* 137: 277–302.
- Otte V. (2004): Zur Bedeutung der *Acer-negundo*-Anpflanzungen auf ehemaligen Rieselfeldern für die Berliner Flechtenflora. – *Verhandlungen des Botanischen Vereins von Berlin und Brandenburg* 137: 373–374.
- Otte V. (2005): Flechten der Lausche (Zittauer Gebirge). – *Berichte der Naturforschenden Gesellschaft der Oberlausitz* 13: 63–66.
- Otte V. (2005): Lichenological observations in Upper Lusatia II. – *Abhandlungen und Berichte des Naturkundemuseums Görlitz* 76: 151–156.
- Otte V., Esslinger T. L. & Litterski B. (2004): Global distribution of the European species of the lichen genus *Melanelia* Essl. – *Journal of Biogeography* 32: 1221–1241.
- Paukov A. G. & Trapeznikova S. N. (2005): Lithophilous lichens of Middle Urals. – *Folia Cryptogamica Estonica* 41: 81–88.
- Paz-Bermúdez G. & Carballal R. (2005): The portuguese specimens of the family *Parmeliaceae* (lichenized ascomycetes) in the PO herbarium (Oporto, Portugal). – *Nova Hedwigia* 81: 205–228.
- Paz-Bermúdez G., López de Silanes M. E. & Carballal R. (2005): The lichen genera *Collemopsidium* Nyl. and *Verrucaria* Schrader from the Galician seashores (NW Spain). – *Nova Hedwigia* 80: 73–87.
- Pintado A., Sancho L. G., Green T. G. A., Blanquer J. M. & Lázaro R. (2005): Functional ecology of the biological soil crust in semiarid SE Spain: sun and shade populations of *Diploschistes diacapsis* (Ach.) Lumbsch. – *Lichenologist* 37: 425–432.

- Priemetzhofer F. (2005): Silikat- und bodenbewohnende Flechten im Mittleren und Unterem Mühlviertel (Oberösterreich, Austria). – Beiträge zur Naturkunde Oberösterreichs 14: 71–146.
- Purvis O. W., Chimonides J., Jeffries T. E., Jones G. C., Read H. & Spiro B. (2005): Investigating biogeochemical signatures in the lichen *Parmelia sulcata* at Burnham Beeches, Buckinghamshire, England. – Lichenologist 37: 329–344.
- Ra H.-S. Y., Geiser L. H. & Crang R. F. E. (2005): Effects of season and low-level air pollution on physiology and element content of lichens from the U.S. Pacific Northwest. – Science of the Total Environment 343: 155–167.
- Ra H.-S. Y., Rubin L. & Crang R. F. E. (2004): Structural impacts on thallus and algal cell components of two lichen species in response to low-level air pollution in Pacific Northwest Forests. – Microscopy and Microanalysis 10: 270–279.
- Radies D. N. & Coxson D. S. (2004): Macrolichen colonization on 120–140 year old *Tsuga heterophylla* in wet temperate rainforests of central-interior British Columbia: a comparison of lichen response to even-aged versus old-growth stand structures. – Lichenologist 36: 235–247.
- Randlane T. & Saag A. (2004): Distribution patterns of some primary and secondary cetrarioid species. – Symbolae Botanicae Upsalienses 34: 359–376.
- Randlane T. & Saag A. (2005): Distribution patterns of primary and secondary species in the genus *Vulpicida*. – Folia Cryptogamica Estonica 41: 89–96.
- Rätzel S., Otte V. & Sipman H. J. M. (2005): Bemerkenswerte Flechtenfunde aus Brandenburg (incl. lichenicoler und lichenoider Pilze) IX. – Verhandlungen des Botanischen Vereins von Berlin und Brandenburg 137: 515–535.
- Rico V. J., van den Boom P. P. G. & Barrasa J. M. (2005): Morphology, chemistry and distribution of *Melanelia sorediella* (*Parmeliaceae*) and similar species in the Iberian Peninsula. – Lichenologist 37: 199–215.
- Rinino S., Bombardi V., Giordani P., Tretiach M., Crisafulli P., Monaci F. & Modenesi P. (2005): New histochemical techniques for the localization of metal ions in the lichen thallus. – Lichenologist 37: 463–466.
- Roux C. (2004): Likenkunajoj kaj supra arbarlimo en la kalkaj Alpoj de suda Francio [Peuplements lichéniques et limite supérieure des forêts dans les Alpes françaises méridionales calcaires]. – Bulletin de la Société Linnéenne de Provence 55: 71–88.
- Roux C. & Sérusiaux E. (2004): Le genre *Strigula* (lichens) en Europe et en Macaronésie. – Bibliotheca Lichenologica 90: 1–96.
- Sanders W. B. (2005): Observing microscopic phases of lichen life cycles on transparent substrata placed in situ. – Lichenologist 37: 373–382.
- Sanders W. B., Moe R. L. & Ascaso C. (2005): Ultrastructural study of the brown alga *Petroderma maculiforme* (*Phaeophyceae*) in the free-living state and in lichen symbiosis with the intertidal marine fungus *Verrucaria tavaresiae* (*Ascomycotina*). – European Journal of Phycology 40: 353–361.
- Santitoro A., Aprile G. G., Baldantoni D., Bartoli G. & Alfani A. (2004): Trace element analyses in an epiphytic lichen and its bark substrate to assess suitability for air biomonitoring. – Environmental Monitoring and Assessment 98: 59–67.
- Scherrer S., Zippler U. & Honegger R. (2005): Characterisation of the mating-type locus in the genus *Xanthoria* (lichen-forming ascomycetes, *Lecanoromycetes*). – Fungal Genetics and Biology 42: 976–988.
- Schiefelbein U. & Rätzel S. (2005): Beiträge zur Flechtenflora Mecklenburg-Vorpommern (Deutschland) und angrenzender Gebiete. – Herzogia 17: 63–77.
- Schmitt I., Martín M. P., Kautz S. & Lumbsch H. T. (2005): Diversity of non-reducing polyketide synthase genes in the *Pertusariales* (lichenized *Ascomycota*): a phylogenetic perspective. – Phytochemistry 66: 1241–1253.
- Schmitt I., Mueller G. & Lumbsch H. T. (2005): Ascoma morphology is homoplasious and phylogenetically misleading in some pyrenocarpous lichens. – Mycologia 97: 362–374.
- Schmull M. & Hauck M. (2005): *Lecidea hercynica*, a new montane epiphytic lichen from Germany. – Lichenologist 37: 485–489.
- Seaward M. R. D. & John V. (2005): A comparative study of the lichens of two cemeteries in Belgrade, Serbia. – Mycologia Balcanica 2: 65–68.
- Seymour F. A., Crittenden P. D., Dickinson M. J., Paoletti M., Montiel D., Cho L. & Dyer P. S. (2005): Breeding systems in the lichen-forming fungal genus *Cladonia*. – Fungal Genetics and Biology 42: 554–563.
- Showman R. E. & Flenniken D. G. (2004): The Macrolichens of Ohio. – Ohio Biological Survey, Columbus. [279 pp.]

- Simon D. M., Hummel C. L., Sheeley S. L. & Bhattacharya D. (2005): Heterogeneity of intron presence or absence in rDNA genes of the lichen species *Physcia aipolia* and *P. stellaris*. – Current Genetics 47: 389–399.
- Simon D. M., Moline J., Helms G., Friedl T. & Bhattacharya D. (2005): Divergent histories of rDNA group I introns in the lichen family *Physciaceae*. – Journal of Molecular Evolution 60: 434–446.
- Sipman H. J. M. & Aptroot A. (2005): Notes on *Mycomicrothelia* (*Arthopyreniaceae* s. lat.), with two new species. – Lichenologist 37: 307–311.
- Sipman H. J. M., Raus T. & Scharlau A. (2005): Some lichens have incomplete distribution ranges in the Aegean (Greece). – Folia Cryptogamica Estonica 41: 97–104.
- Skuterud L., Gwynn J. P., Gaare E., Steinnes E. & Hove K. (2005): ^{90}Sr , ^{210}Po and ^{210}Pb in lichen and reindeer in Norway. – Journal of Environmental Radioactivity 84: 441–456.
- Suija A., Nõmm M. & Boch S. (2005): New Estonian lichens and lichenicolous fungi. – Folia Cryptogamica Estonica 41: 135–136.
- Sun H. J. & Friedman E. I. (2005): Communities adjust their temperature optima by shifting producer-to-consumer ratio, shown in lichens as models: II. Experimental verification. – Microbial Ecology 49: 528–535.
- Svensson M., Johansson P. & Thor G. (2005): Lichens of wooden barns and *Pinus sylvestris* snags in Dalarna, Sweden. – Annales Botanici Fennici 42: 351–363.
- Takahashi K., Kinoshita K., Yamamoto Y., Koyama K. & Yoshimura I. (2005): Chemical constituents from lichens for pharmaceutical and industrial uses. – Folia Cryptogamica Estonica 41: 109–114.
- Taylor T. N., Hass H., Kerp H., Krings M. & Hanlin R. T. (2005): Perithecial ascomycetes from the 400 million year old Rhynie chert: an example of ancestral polymorphism. – Mycologia 97: 269–285.
- Tehler A., Dahlkild Å., Eldenäs P. & Feige G. B. (2004): The phylogeny and taxonomy of Macaronesian, European and Mediterranean *Roccella* (*Roccellaceae*, *Arthoniales*). – Symbolae Botanicae Upsalienses 34: 405–428.
- Temina M., Nevo E. & Wasser S. P. (2005): The lichen genus *Acarospora* in Israel and its vicinity. – Nova Hedwigia 80: 433–452.
- Temina M., Wasser S. P. & Nevo E. (2002): A contribution to the species diversity of lichens in Israel. – Flora Mediterranea 12: 285–298.
- Thell A., Herber B., Aptroot A., Adler M. T., Feuerer T. & Kärnefelt E. I. (2005): A preliminary phylogeographic study of *Flavopunctelia* and *Punctelia* inferred from rDNA ITS-sequences. – Folia Cryptogamica Estonica 41: 115–122.
- Thell A., Randlane T., Saag A. & Kärnefelt I. (2005): A new circumscription of the lichen genus *Nephromopsis* (*Parmeliaceae*, lichenized *Ascomycetes*). – Mycological Progress 4: 303–316.
- Thell A., Westberg M. & Kärnefelt I. (2004): Biogeography of the lichen family *Parmeliaceae* in the Nordic countries with taxonomic remarks. – Symbolae Botanicae Upsalienses 34: 429–452.
- Thor G. (2004): *Opegrapha suecica*, a new lichen from Europe. – Symbolae Botanicae Upsalienses 34: 453–454.
- Timdal E. & Zhurbenko M. P. (2004): *Psora indigirkiae*, a new species from eastern Siberia, with key to the Asian species of *Psora*. – Symbolae Botanicae Upsalienses 34: 25–29.
- Tõrra T. & Randlane T. (2005): What are diagnostic characters in *Usnea*? A survey of Baltic species. – In: Proceedings of the XVI Symposium of Mycologists and Lichenologists of Baltic States. 21–25 September 2005, p. 35–40, Cesis, Latvia.
- Tretiach M. (2004): Further additions to the Italian lichen flora. – Cryptogamie Mycologie 25: 173–183.
- Tretiach M., Crisafulli P., Pittao E., Rinino S., Roccotielo E. & Modenesi P. (2005): Isidia ontogeny and its effect on the CO_2 gas exchanges of the epiphytic lichen *Pseudevernia furfuracea* (L.) Zopf. – Lichenologist 37: 445–462.
- Türk R. (2004): Neue und seltene Flechten der Ostalpen. – Verhandlungen der Zoologisch-Botanischen Gesellschaft in Österreich 141: 63–73.
- Türk R. & Antesberger B. (2004): Epiphytische Flechten in der Achartinger Au. – Mitteilungen aus der Haus der Natur 16: 65–69.
- Türk R. & Riegler-Hager H. (2003): Eine bemerkenswerte Flechtensammlung im Kärntner Landesherbar (KL). – Wulfenia 10: 171–176.
- van den Boom P. P. G. (2005): Contribution to the flora of Portugal, lichens and lichenicolous fungi IV. – Cryptogamie Mycologie 26: 51–59.
- van den Boom P. P. G. & Brand A. M. (2005): *Lecania fructigena* Zahlbr., a coastal saxicolous lichen, new for Europe, with notes on related species. – Lichenologist 37: 277–283.

- van den Boom P. P. G. & Elix J. A. (2005): Notes on *Halecania* species, with descriptions of two new species from Asia. – *Lichenologist* 37: 237–246.
- van den Boom P. P. G. & Vězda A. (2005): *Gyalecta canariensis* sp. nov., a new lichen (*Ascomycota*) described from La Palma (Canary Islands). – *Mycotaxon* 92: 255–258.
- van Herk K. & Aptroot A. (2004): *Veldgids Korstmossen*. – Stichting Uitgeverij van de KNNV, Utrecht. [421 pp.]
- Walser J.-C., Holderegger R., Gugerli F., Hoebee S. E. & Scheidegger C. (2005): Microsatellites reveal regional population differentiation and isolation in *Lobaria pulmonaria*, an epiphytic lichen. – *Molecular Ecology* 14: 457–467.
- Wedin M., Döring H., Könberg K. & Gilenstam G. (2005): Generic delimitations in the family *Stictidaceae* (*Ostropales, Ascomycota*): the *Stictis-Conotrema* problem. – *Lichenologist* 37: 67–75.
- Wedin M. & Wiklund E. (2004): The phylogenetic relationships of *Lecanorales* suborder *Peltigerineae* revisited. – *Symbolae Botanicae Upsalienses* 34: 469–475.
- Wedin M., Wiklund E., Crewe A., Döring H., Ekman S., Nyberg Å., Schmitt I. & Lumbsch H. T. (2005): Phylogenetic relationships of *Lecanoromycetes* (*Ascomycota*) as revealed by analyses of mtSSU and nLSU rDNA sequence data. – *Mycological Research* 109: 159–172.
- Weissman L., Garty J. & Hochman A. (2005): Rehydration of the lichen *Ramalina lacera* results in production of reactive oxygen species and nitric oxide and a decrease in antioxidants. – *Applied and Environmental Microbiology* 71: 2121–2129.
- Werth S., Tømmervik H. & Elvebakk A. (2005): Epiphytic macrolichen communities along regional gradients in northern Norway. – *Journal of Vegetation Science* 16: 199–208.
- Wolterbeek H. T., Jeran Z., Verburg T., de Goeij J. J. M. & van Dam J. (2004): Chernobyl ^{137}Cs in lichens: Use of specific activity showing differences in transport routes of Cs and ^{137}Cs . – *Journal of Radioanalytical and Nuclear Chemistry* 259: 13–18.
- Yuan X. L., Xiao S. H. & Taylor T. N. (2005): Lichen-like symbiosis 600 million years ago. – *Science* 308: 1017–1020.
- Zhao Z., Ren Q. & Aptroot A. (2004): An annotated key of *Pertusaria* in China. – *Bryologist* 107: 531–541.
- Zhurbenko M. & Ahti T. (2005): Contribution to the study of the lichen genera *Cladina* and *Cladonia* in the Russian Arctic, mainly from Taimyr Peninsula and Severnaya Zemlya. – *Nova Hedwigia* 81: 79–96.
- Zhurbenko M. P., Laursen G. A. & Walker D. A. (2005): New and rare lichenicolous fungi and lichens from the North American Arctic. – *Mycotaxon* 92: 201–212.
- Zocher B. & Stocker-Wörgötter E. (2005): Secondary chemistry of lichen-forming ascomycetes: culture experiments with isolates of *Xanthoparmelia* species from Australia. – *Folia Cryptogamica Estonica* 41: 123–134.