

NOVÁ BRYOLOGICKÁ LITERATURA XIX.

New bryological literature, XIX

Jan Kučera¹, Svatava Kubelová² & Vítězslav Plášek³

¹ Jihočeská Univerzita, Přírodovědecká fakulta, Branišovská 31, CZ-370 05 České Budějovice, e-mail: kucera@prf.jcu.cz, ² Botanické oddělení, Moravské zemské muzeum, Hviezdoslavova 29a, CZ-62700 Brno, skubesova@mzm.cz; ³ Ostravská univerzita, KBE PřF, Chittussiho 10, CZ-71000 Ostrava, vitezslav.plasek@osu.cz

Výběr ze světové bryologické literatury [Selection from the world bryological literature]

- Aboal J. R., Fernández J. A., Couto J. A. & Carballeira A. (2008): Testing differences in methods of preparing moss samples. – Environmental Monitoring and Assessment 137: 371–378.
- Adamo P., Bargagli R., Giordano S., Modenesi P., Monaci F., Pittao E., Spagnuolo V. & Tretiach M. (2008): Natural and pre-treatments induced variability in the chemical composition and morphology of lichens and mosses selected for active monitoring of airborne elements. – Environmental Pollution 152: 11–19.
- Adams D. G. & Duggan P. S. (2008): Cyanobacteria-bryophyte symbioses. – Journal of Experimental Botany 59: 1047–1058.
- Afonina O. M., Ignatova E. A. & Maksimov A. I. (2006): *Stereodon fertilis* (*Pylaisiaceae, Musci*) in Russia. – Botanicheskiy Zhurnal 91: 329–335.
- Anterola A. & Shanle E. (2008): Genomic insights in moss gibberellin biosynthesis. – Bryologist 111: 218–230.
- Asakawa Y. (2007): Biologically active compounds from bryophytes. – Chenia 9: 73–104.
- Astel A., Astel K. & Biziuk M. (2008): PCA and multidimensional visualization techniques united to aid in the bioindication of elements from transplanted *Sphagnum palustre* moss exposed in the Gdańsk city area. – Environmental Science and Pollution Research 15: 41–50.
- Bączkiewicz A., Sawicki J., Buczkowska K., Polok K. & Zieliński R. (2008): Application of different DNA markers in studies on cryptic species of *Aneura pinguis* (*Jungermanniopsida, Metzgeriales*). – Cryptogamie Bryologie 29: 3–21.
- Balan C., Bilba D. & Macoveanu M. (2008): Removal of cadmium (II) from aqueous solutions by *Sphagnum* moss peat: Equilibrium study. – Environmental Engineering and Management Journal 7: 17–23.
- Baptista M. S., Vasconcelos M. T. S. D., Cabral J. P., Freitas M. C. & Pacheco A. M. G. (2008): Copper, nickel and lead in lichen and tree bark transplants over different periods of time. – Environmental Pollution 151: 408–413.
- Barandovski L., Cekova M., Frontasyeva M. V., Pavlov S. S., Stafilov T., Steinnes E. & Urumov V. (2008): Atmospheric deposition of trace element pollutants in Macedonia studied by the moss biomonitoring technique. – Environmental Monitoring and Assessment 138: 107–118.
- Basile A., Sorbo S., Aprile G., Conte B. & Cobianchi R. C. (2008): Comparison of the heavy metal bioaccumulation capacity of an epiphytic moss and an epiphytic lichen. – Environmental Pollution 151: 401–407.
- Bates J. W., Matcham H. W. & Lara F. (2007): *Dalytrichia fragilifolia* (*Bryopsida: Pottiaceae*) in Berkshire and Caernarvonshire, new to Britain. – Journal of Bryology 29: 228–234.
- Belnap J., Phillips S. L. & Smith S. D. (2007): Dynamics of cover, UV-protective pigments, and quantum yield in biological soil crust communities of an undisturbed Mojave Desert shrubland. – Flora 202: 674–686.
- Bergamini A., Müller N., Peintinger M., Schnyder N. & Stofer S. (2007): Die Wassermoosflora des Rheinfalls bei Schaffhausen im Vergleich zu den 1930er Jahren. – Botanica Helvetica 117: 202–207.
- Black P. L., Arnason J. T., Cuerrier A. (2008): Medicinal plants used by the Inuit of Qikiqtaaluk (Baffin island, Nunavut). – Botany – Botanique 86: 157–163.
- Boudreault C., Coxson D. S., Vincent E., Bergeron Y. & Marsh J. (2008): Variation in epiphytic lichen and bryophyte composition and diversity along a gradient of productivity in *Populus tremuloides* stands of northeastern British Columbia, Canada. – Ecoscience 15: 101–112.
- Breeuwer A., Heijmans M. M. P. D., Robroek B. J. M. & Berendse F. (2008): The effect of temperature on growth and competition between *Sphagnum* species. – Oecologia 156: 155–167.

- Brinda J. C., Stark L. R., Shevock J. R. & Spence J. R. (2007): An annotated checklist of the bryophytes of Nevada, with notes on collecting history in the state. – *Bryologist* 110: 673–705.
- Cailliau A. & Price M. J. (2007): Hornworts in the agricultural fields of Geneva: new findings, the soil diaspore bank and ex situ soil cultures. – *Candollea* 62: 165–172.
- Calabrese G. M. & Muñoz J. (2008): *Zygodon (Orthotrichaceae)* in the Iberian Peninsula. – *Bryologist* 111: 231–247.
- Callaghan D. A. & Ashton P. A. (2007): Bryophyte clusters and sand dune vegetation communities. – *Journal of Bryology* 29: 213–221.
- Cano M. J. & Gallego M. T. (2008): The genus *Tortula* (*Pottiaceae*, *Bryophyta*) in South America. – *Botanical Journal of the Linnean Society* 156: 173–220.
- Cassie D. M., Piercy-Normore M. D., and Belland R. J. (2008): Population Structure of *Dicranum elongatum* in Northeastern Regions of Wapusk National Park, Manitoba, Canada. – *Bryologist* 111: 302–309.
- Chambers P. A., Lacoul P., Murphy K. J. & Thomaz S. M. (2008): Global diversity of aquatic macrophytes in freshwater. – *Hydrobiologia* 595: 9–26.
- Clarke L. J., Ayre D. J. & Robinson S. A. (2008): Somatic mutation and the Antarctic ozone hole. – *Journal of Ecology* 96: 378–385.
- Cvetić T., Sabovljević A., Sabovljević M. & Grubišić D. (2007): Development of the moss *Pogonatum urnigerum* (Hedw.) P. Beauv. under *in vitro* culture conditions. – *Archives of Biological Sciences* 59: 57–61.
- Deckor E. L. & Reski R. (2007): Moss bioreactors producing improved biopharmaceuticals. – *Current Opinion In Biotechnology* 18: 393–398.
- Diéguez C., Rodríguez-López J. P. & Meléndez N. (2007): Marchantiopsid colonization mats from the Upper Aptian–Lower Albian of the Escucha Formation (Oliete Sub-Basin, Iberian Ranges, eastern Spain). – *Comptes Rendus Palevol* 6: 413–422.
- Draper I., Mazimpaka V. & Lara F. (2008): New records to the epiphytic bryophyte flora of Tunisia. – *Cryptogamie Bryologie* 29: 83–91.
- Drozd P., Plášek V., Dolný A., Kočárek P. & Jašík M. (2007): Factors or mosses – What the bryobionts prefer? – *Nowellia Bryologica* 34: 9–10.
- Ellis L. T. (2007): Bryophyte flora of Uganda. 7. *Calymperaceae* and *Octoblepharaceae*. – *Journal of Bryology* 29: 259–274.
- Equihua C. & Equihua M. (2007): Spatial distribution of *Bryopteris filicina* on tree trunks in the Selva Lacandona, Chiapas, Mexico. – *Bryologist* 110: 766–775.
- Erzberger P. & Papp B. (2007): New and noteworthy bryophyte records from Montenegro and Serbia. – *Willdenowia* 37: 339–352.
- Espinar J. L. & Clemente L. (2007): The impact of vertic soil cracks on submerged macrophyte diaspore bank depth distribution in Mediterranean temporary wetlands. – *Aquatic Botany* 87: 325–328.
- Fisher K. M. (2008): Bayesian reconstruction of ancestral expression of the LEA gene families reveals propagule-derived desiccation tolerance in resurrection plants. – *American Journal of Botany* 95: 506–515.
- Fojcik B., Stebel A., Fudali E., Plášek V., Rusińska A., Żarnowiec J., Zmrhalová M., Zubel R., Gorski P., Cykowska B. & Wilhelm M. (2007): Materiały do brioflory Ojcowskiego Parku Narodowego [Contribution to the bryoflora of Ojców National Park]. – Prądnik. Prace i Materiały Muzeum Im. Prof. W. Szafera, 17: 79–94.
- Frahm J.-P. (2008): Diversity, dispersal and biogeography of bryophytes (mosses). – *Biodiversity and Conservation* 17: 277–284.
- Fritz Ö., Gustafsson L. & Larsson K. (2008): Does forest continuity matter in conservation? – A study of epiphytic lichens and bryophytes in beech forests of southern Sweden. – *Biological Conservation* 141: 655–668.
- Fudali E. (2007): Human traces in the bryophyte flora of the summit region of Karkonosze Mts (Polish side). – *Acta Societatis Botanicorum Poloniae* 76: 345–349.
- Fujita T., Sakaguchi H., Hiwatashi Y., Wagstaff S. J., Ito M., Deguchi H., Sato T. & Hasebe M. (2008): Convergent evolution of shoots in land plants: lack of auxin polar transport in moss shoots. – *Evolution & Development* 10: 176–186.
- Fuselier L. (2008): Variation in life history characteristics between asexual and sexual populations of *Marchantia inflexa*. – *Bryologist* 111: 248–259.
- Garbary D. J., Miller A. G., Scrosati R., Kim K.-Y., & Schofield W. B. (2008): Distribution and salinity tolerance of intertidal mosses from Nova Scotian salt marshes. – *Bryologist* 111: 282–291.

- Garcia C., Sérgio C. & Jansen J. (2008): The bryophyte flora of the natural park of Serra da Estrela (Portugal): Conservation and Biogeographical approaches. – *Cryptogamie Bryologie* 29: 49–73.
- Grimm A., Zanzi R., Björnbom E. & Cukierman A. L. (2008): Comparison of different types of biomasses for copper biosorption. – *Bioresource Technology* 99: 2559–2565.
- Groeneveld E. V. G., Masse A. & Rochefort L. (2007): *Polytrichum strictum* as a nurse-plant in peatland restoration. – *Restoration Ecology* 15: 709–719.
- Grundmann M., Ansell S. W., Russell S. J., Koch M. A. & Vogel J. C. (2008): Hotspots of diversity in a clonal world - the Mediterranean moss *Pleurochaete squarrosa* in Central Europe. – *Molecular Ecology* 17: 825–838.
- Guerra J. (2007): *Pohlia* section *Cacodon* (*Mielichhoferiaceae*, *Bryophyta*) with axillary bulbils in the Iberian Peninsula. – *Anales del Jardín Botánico de Madrid* 64: 55–62.
- Hájek T. & Beckett R. P. (2008): Effect of water content components on desiccation and recovery in *Sphagnum* mosses. – *Annals of Botany* 101: 165–173.
- Hájková P., Shaw B., Hájek M., Hinterlang D. & Plášek V. (2007): The role of base saturation and altitude in habitat differentiation within *Philonotis* in springs and mires of three different European regions. – *Bryologist* 110: 776–787.
- Harmens H., Norris D. A., Koerber G. R., Buse A., Steinnes E. & Rühling Å. (2008): Temporal trends (1990–2000) in the concentration of cadmium, lead and mercury in mosses across Europe. – *Environmental Pollution* 151: 368–376.
- Harris E. S. J. (2008): Ethnobryology: traditional uses and folk classification of bryophytes. – *Bryologist* 111: 169–217.
- Hautala H., Tolvanen A. & Nuortila C. (2008): Recovery of pristine boreal forest floor community after selective removal of understorey, ground and humus layers. – *Plant Ecology* 194: 273–282.
- He-Nygrén X. (2007): Multi-gene phylogeny supports single origin of jungermannioid perigynium. – *Annales Botanici Fennici* 44: 450–462.
- Heber U., Azarkovich M. & Shuvalov V. (2007): Activation of mechanisms of photoprotection by desiccation and by light: poikilohydric photoautotrophs. – *Journal of Experimental Botany* 58: 2745–2759.
- Hedderson T. A. & Zander R. H. (2007): *Ludorugbya springbokorum* (*Pottiaceae*) a new moss genus and species from the Western Cape Province of South Africa. – *Journal of Bryology* 29: 222–227.
- Hedenäs L. (2008): Molecular variation and speciation in *Antitrichia curtipendula* s.l. (*Leucodontaceae*, *Bryophyta*). – *Botanical Journal of the Linnean Society* 156: 341–354.
- Hedenäs L. & Eldenäs P. (2007): Cryptic speciation, habitat differentiation, and geography in *Hamatocaulis vernicosus* (*Calliergonaceae*, *Bryophyta*). – *Plant Systematics and Evolution* 268: 131–145.
- Hedenäs L. & Eldenäs P. (2008): Relationships in *Scorpidium* (*Calliergonaceae*, *Bryophyta*), especially between *S. cossonii* and *S. scorpioides*. – *Taxon* 57: 121–130.
- Heijmans M. M. P. D., Mauquoy D., van Geel B. & Berendse F. (2008): Long-term effects of climate change on vegetation and carbon dynamics in peat bogs. – *Journal of Vegetation Science* 19: 307–U54.
- Heino J. & Korsu K. (2008): Testing species-stone area and species-bryophyte cover relationships in riverine macroinvertebrates at small scales. – *Freshwater Biology* 53: 558–568.
- Heino J. & Toivonen H. (2008): Aquatic plant biodiversity at high latitudes: patterns of richness and rarity in Finnish freshwater macrophytes. – *Boreal Environment Research* 13: 1–14.
- Hentschel J., Feldberg K., Zündorf H.-J., Hellwig F. H., Schneider H. & Heinrichs J. (2007): The systematic position of *Pachyglossa* and *Clasmatocolea* (*Jungermanniopsida*: *Lophocoleaceae*) inferred from nrDNA ITS sequences and morphology. – *Taxon* 56: 1136–1142.
- Hentschel J., Paton J. A., Schneider H. & Heinrichs J. (2007): Acceptance of *Lioclada* Nees and *Solenostoma* Mitt., the systematic position of *Eremonotus* Pearson and notes on *Jungermannia* L. s.l. (*Jungermanniidae*) based on chloroplast DNA sequence data. – *Plant Systematics and Evolution* 268: 147–157.
- Hentschel J., Zhu R.-L., Long D. G., Davison P. G., Schneider H., Gradstein S. R. & Heinrichs J. (2007): A phylogeny of *Porella* (*Porellaceae*, *Jungermanniopsida*) based on nuclear and chloroplast DNA sequences. – *Molecular Phylogenetics and Evolution* 45: 693–705.
- Hernández-Maqueza R., Quandt D., Werner O. & Muñoz J. (2008): Phylogeny and classification of the *Grimmiaceae/Ptychomitriaceae* complex (*Bryophyta*) inferred from cpDNA. – *Molecular Phylogenetics and Evolution* 46: 863–877.

- Hernick L., VanAller, Landing E. & Bartowski K. E. (2008): Earth's oldest liverworts – *Metzgeriothallus haronae* sp. nov. from the Middle Devonian (Givetian) of eastern New York, USA. – Review of Palaeobotany and Palynology 148: 154–162.
- Heylen O. & Hermy M. (2008): Age structure and ecological characteristics of some epiphytic liverworts (*Frullania dilatata*, *Metzgeria furcata* and *Radula complanata*). – Bryologist 111: 84–97.
- Hock Z., Szövényi P., Schneller J. J., Tóth Z. & Urmi E. (2008): Bryophyte diaspore bank: A genetic memory? Genetic structure and genetic diversity of surface populations and diaspore bank in the liverwort *Mannia fragrans* (Athyriaceae). – American Journal of Botany 95: 542–548.
- Hooijmaijers C. A. M. (2008): Desiccation tolerance in red and green gametophytes of *Jamesoniella colorata* in relation to photoprotection. – Planta 227: 1301–1310.
- Hutsemekers V., Dopagne C. & Vanderpoorten A. (2008): How far and how fast do bryophytes travel at the landscape scale?. – Diversity and Distributions 14: 483–492.
- Ingerpuu N. & Vellak K. (2007): Collections of G. C. Girgensohn (1786–1872): Lectotypes and rare species. – Journal of Bryology 29: 235–240.
- Jankowiak-Siuda K., Pacak A., Odrzykoski I., Wyatt R. & Szweykowska-Kulińska Z. (2008): Organellar inheritance in the allopolyploid moss *Plagiomnium curvatum*. – Taxon 57: 145–152.
- Jones M. R., Leith I. D., Raven J. A., Fowler D., Sutton M. A., Nemitz E., Cape J. N., Sheppard L. J. & Smith R. I. (2007): Concentration-dependent NH₃ deposition processes for moorland plant species with and without stomata. – Atmospheric Environment 41: 8980–8994.
- Kalmykova Y., Strömwall A.-M. & Steenari B.-M. (2008): Adsorption of Cd, Cu, Ni, Pb and Zn on *Sphagnum* peat from solutions with low metal concentrations. – Journal of Hazardous Materials 152: 885–891.
- Kapila S., Kumar S. S. & Kaur R. (2007): Effect of zinc, copper and iron on regeneration of *Hyophila involuta* (Hook.) Jaeg. and *Funaria hygrometrica* Hedw. – Hikobia 15: 99–103.
- Kempter H. & Frenzel B. (2008): Titanium in ombrotrophic *Sphagnum* mosses from various peat bogs of Germany and Belgium. – Science of the Total Environment 392: 324–334.
- Khatun H. & Hadiuzzaman S. (2007): Additions to the pleurocarpous mosses of Bangladesh: Family – *Plagiotheciaceae*. – Bangladesh Journal of Plant Taxonomy 14: 79–82.
- Kicsi A., Bilba D. & Macoveanu M. (2007): Removal of zinc (II) from aqueous solutions by Romanian spragnum peat moss. – Environmental Engineering and Management Journal 6: 205–209.
- Klanderud K. (2008): Species-specific responses of an alpine plant community under simulated environmental change. – Journal of Vegetation Science 19: 363–U109.
- Köchy M. & Bråkenhielm S. (2008): Separation of effects of moderate N deposition from natural change in ground vegetation of forests and bogs. – Forest Ecology and Management 255: 1654–1663.
- Kosior G., Samecka-Cymerman A., Chmielewski A., Wierzchnicki R., Derda M. & Kempers A. J. (2008): Native and transplanted *Pleurozium schreberi* (Brid.) Mitt. as a bioindicator of N deposition in a heavily industrialized area of Upper Silesia (S Poland). – Atmospheric Environment 42: 1310–1318.
- Kottke I., Haug I., Setaro S., Suárez J. P., Weiß M., Preußing M., Nebel M. & Oberwinkler F. (2008): Guilds of mycorrhizal fungi and their relation to trees, ericads, orchids and liverworts in a neotropical mountain rain forest. – Basic and Applied Ecology 9: 13–23.
- Koz B., Cevik U., Ozdemir T., Duran C., Kaya S., Gundogdu A. & Celik N. (2008): Analysis of mosses along Sarp-Samsun highway in Turkey. – Journal of Hazardous Materials 153: 646–654.
- Krab E. J., Cornelissen J. H. C., Lang S. I. & van Logtestijn R. S. P. (2008): Amino acid uptake among wide-ranging moss species may contribute to their strong position in higher-latitude ecosystems. – Plant and Soil 304: 199–208.
- Kürschner H. (2008): A key to the acrocarpous mosses (*Bryophytina* p. p., excl. *Pottiaceae*) of the Near and Middle East. Towards a bryophyte flora of the Near and Middle East, 7. – Nova Hedwigia 86: 43–103.
- Kürschner H., Stech M., Sim-Sim M., Fontinha S. & Frey W. (2007): Life form and life strategy analyses of the epiphytic bryophyte communities of Madeira's laurel and ericaceous forests. – Botanische Jahrbücher für Systematik, Pflanzengeschichte und Pflanzengeographie 127: 151–164.
- Lagerström A., Nilsson M. -C., Zackrisson O. & Wardle D. A. (2007): Ecosystem input of nitrogen through biological fixation in feather mosses during ecosystem retrogression. – Functional Ecology 21: 1027–1033.
- Lalanne A., Bardat J., Lalanne-Amara F., Gautrot T. & Ponge J.-F. (2008): Opposite responses of vascular plant and moss communities to changes in humus form, as expressed by the Humus Index. – Journal of Vegetation Science 19: 645–652.

- Lappalainen N. M., Huttunen S. & Suokanerva H. (2008): Acclimation of a pleurocarpous moss *Pleurozium schreberi* (Britt.) Mitt. to enhanced ultraviolet radiation *in situ*. – Global Change Biology 14: 321–333.
- Ligrone R., Carafa A., Duckett J. G., Renzaglia K. S. & Ruel K. (2008): Immunocytochemical detection of lignin-related epitopes in cell walls in bryophytes and the charalean alga *Nitella*. – Plant Systematics and Evolution 270: 257–272.
- Ligrone R., Carafa A., Lumini E., Bianciotto V., Bonfante P. & Duckett J. G. (2007): Glomeromycotan associations in liverworts: A molecular cellular and taxonomic analysis. – American Journal of Botany 94: 1756–1777.
- Lüttge U., Meirelles S. T. & de Mattos E. A. (2008): Strong quenching of chlorophyll fluorescence in the desiccated state in three poikilohydric and homoioclorophyllous moss species indicates photo-oxidative protection on highly light-exposed rocks of a tropical inselberg. – Journal of Plant Physiology 165: 172–181.
- Mälson K., Backéus I. & Rydin H. (2008): Long-term effects of drainage and initial effects of hydrological restoration on rich fen vegetation. – Applied Vegetation Science 11: 99–106.
- Man R., Kayahara G. J., Rice J. A. & MacDonald G. B. (2008): Eleven-year responses of a boreal mixedwood stand to partial harvesting: Light, vegetation, and regeneration dynamics. – Forest Ecology and Management 255: 697–706.
- Merdanić H., Smailhodžić H., Ferhatović D. & Osmanović Z. (2007): Indicating water quality by moss. – Journal of Environmental Protection and Ecology 8: 520–525.
- Nabe H., Funabiki R., Kashino Y., Koike H. & Satoh K. (2007): Responses to desiccation stress in bryophytes and an important role of dithiothreitol-insensitive non-photochemical quenching against photoinhibition in dehydrated states. – Plant and Cell Physiology 48: 1548–1557.
- Newton A. E. & Tangney R. S. (eds.) (2007): Pleurocarpous mosses: systematics and evolution. – Systematics Association Special Volume 71, CRC Press. [434 pp.]
- Nimptsch J. & Pflugmacher S. (2008): Decomposing leaf litter: The effect of allochthonous degradation products on the antioxidant fitness and photosynthesis of *Vesicularia dubyana*. – Ecotoxicology and Environmental Safety 69: 541–545.
- Nöske N. M., Hilt N., Werner F. A., Brehm G., Fiedler K., Sipman H. J. M. & Gradstein S. R. (2008): Disturbance effects on diversity of epiphytes and moths in a montane forest in Ecuador. – Basic and Applied Ecology 9: 4–12.
- Ohyama K. & Takeura M. (2008): Molecular evolution of mitochondrial introns in the liverwort *Marchantia polymorpha*. – Proceedings of the Japan Academy Series B – Physical and Biological Sciences 84: 17–23.
- Økland R. H., Rydgren K. & Økland T. (2008): Species richness in boreal swamp forests of SE Norway: The role of surface microtopography. – Journal of Vegetation Science 19: 67–74.
- Pakeman R. J., Reid C. L., Lennon J. J. & Kent M. (2008): Possible interactions between environmental factors in determining species optima. – Journal of Vegetation Science 19: 201–U4.
- Panov V. V. (2006): Some features of the *Sphagnum* moss cover development in bogs. – Botanicheskiy Zhurnal 91: 393–401.
- Papp B., Ódor P. & Szurdoki E. (2007): Bryophyte biodiversity monitoring system in Hungary, eastern central Europe. – Chenia 9: 143–158.
- Peck J. & Muir P. (2008): Biomass inventory and regrowth rate of harvestable tree and shrub moss in the Oregon coast range. – Western Journal of Applied Forestry 23: 34–39.
- Peck J., Hoganson H., Muir P., Ek A. & Frelich L. (2008): Using inventory projections to evaluate management options for the nontimber forest product of epiphytic moss. – Forest Science 54: 185–194.
- Peck J. E. & Frelich L. E. (2008): Moss harvest truncates the successional development of epiphytic bryophytes in the Pacific Northwest. – Ecological Applications 18: 146–158.
- Pekka L., Halmeenpää H., Ecke F., Vuori K.-M., Mokrotovarova O., Öhlander B. & Ingri J. (2008): Assessing pollution in the Kola River, northwestern Russia, using metal concentrations in water and bryophytes. – Boreal Environment Research 13: 15–30.
- Pesch R., Schröder W. & Schmidt G. (2007): Nitrogen accumulation in forests. Exposure monitoring by mosses. – The Scientific World 7: 151–158.
- Phuyal M., Artz R. R. E., Sheppard L., Leith I. D. & Johnson D. (2008): Long-term nitrogen deposition increases phosphorus limitation of bryophytes in an ombrotrophic bog. – Plant Ecology 196: 111–121.
- Plášek V. (2007): Gatunki z rodziny *Orthotrichaceae* (*Bryophyta*) i problemy ich ochrony. – In: Kasza H. & Klama H., ed., Zapobieganie zanieczyszczeniu, przekształcaniu i degradacji środowiska XIV, p. 137–

- 140, Wydział Nauk o Materiałach i Środowisku, Instytut Ochrony i Inżynierii Środowiska, Wydaw. Akademii Techniczno-Humanistycznej, Bielsko-Biała.
- Plášek V., Drozd P. & Wierzcholska S. (2007): The chorology, ecology & population biology of the genus *Orthotrichum*, *Ulota* & *Zygodon* in the Central Europe. Part I. Methods & first results. – *Nowellia Bryologica* 33: 39–47.
- Plášek V. & Filipová K. (2007): *Bryophyta a Tracheophyta* Přírodní památky Věřňovice. – In: Lis J.A. & Mazur M. (2007), Przyrodnicze wartości Polsko-Czeskiego pogranicza jako wspólne dziedzictwo Unii Europejskiej, p. 99–111, Centrum Studiów nad Bioróżnorodnością, Uniwersytet Opolski, Opole.
- Plášek V., Franková H., Nociarová D. & Podstawková D. (2007): Rozšíření a ekologické nároky epifytních mechů ovlněně vertikální a horizontální struktury Moravské brány. – In: Lis J.A. & Mazur M. (2007), Przyrodnicze wartości Polsko-Czeskiego pogranicza jako wspólne dziedzictwo Unii Europejskiej, p. 221, Centrum Studiów nad Bioróżnorodnością, Uniwersytet Opolski, Opole.
- Plášek V. & Wierzcholska S. (2007): Bryophytes collected during the Bryological Meeting in Vierves-sur-Viroin. – *Nowellia Bryologica* 34: 84–85.
- Plášek V. & Wierzcholska S. (2007): *Orthotrichum*, *Ulota* & *Zygodon* in the Central Europe – their chorology, ecology and conservation problems. – *Nowellia Bryologica* 34: 11–12.
- Pressel S., Ligrone R., Duckett J. G. & Davis E. C. (2008): A novel ascomycetous endophytic association in the rhizoids of the leafy liverwort family *Schistochilaceae* (*Jungermanniidae*, *Hepaticopsida*). – *American Journal of Botany* 95: 531–541.
- Pressel S., Matcham H. W. & Duckett J. G. (2007): Studies of protonemal morphogenesis in mosses. XI. *Bryum* and allied genera: a plethora of propagules. – *Journal of Bryology* 29: 241–258.
- Price J. S., Whittington P. N., Elrick D. E., Strack M., Brunet N. & Faux E. (2008): A method to determine unsaturated hydraulic conductivity in living and undecomposed *Sphagnum* moss. – *Soil Science Society of America Journal* 72: 487–491.
- Proctor M. C. F., Oliver M. J., Wood A. J., Alpert P., Stark L. R., Cleavitt N. L. & Mishler B. D. (2007): Desiccation-tolerance in bryophytes: a review. – *Bryologist* 110: 595–621.
- Pursell R. A. (2007): *Fissidentaceae*. – In: Flora Neotropica 101, Organization for Flora Neotropica by The New York botanical Garden: New York. [i–iii, 1–279]
- Pyatygina S. S., Opritov V. A. & Vodeneev V. A. (2008): Signaling role of action potential in higher plants. – *Russian Journal of Plant Physiology* 55: 285–291.
- Rensing S. A., Lang D., Zimmer A. D., Terry A., Salamov A., Shapiro H., Nishiyama T., Perroud P.-F., Lindquist E. A., Kamisugi Y., Tanahashi T., Sakakibara K., Fujita T., Oishi K., Shin-I T., Kuroki Y., Toyoda A., Suzuki Y., Hashimoto S.-i., Yamaguchi K., Sugano S., Kohara Y., Fujiyama A., Anterola A., Aoki S., Ashton N., Barbazuk W. B., Barker E., Bennetzen J. L., Blankenship R., Cho S. H., Dutcher S. K., Estelle M., Fawcett J. A., Gundlach H., Hanada K., Heyl A., Hicks K. A., Hughes J., Lohr M., Mayer K., Melkozernov A., Murata T., Nelson D. R., Pils B., Prigge M., Reiss B., Renner T., Rombauts S., Rushton P. J., Sanderfoot A., Schween G., Shiu S.-H., Stueber K., Theodoulou F. L., Tu H., Van de Peer Y., Verrier P. J., Waters E., Wood A., Yang L., Cove D., Cuming A. C., Hasebe M., Lucas S., Mishler B. D., Reski R., Grigoriev I. V., Quatrano R. S. & Boore J. L. (2008): The *Physcomitrella* genome reveals evolutionary insights into the conquest of land by plants. – *Science* 319: 64–69.
- Rikkinen J. & Virtanen V. (2008): Genetic diversity in cyanobacterial symbionts of thalloid bryophytes. – *Journal of Experimental Botany* 59: 1013–1021.
- Ringwall A., Snäll T., Ekström M. & Ståhl G. (2007): Unrestricted guided transect sampling for surveying sparse species. – *Canadian Journal of Forest Research – Revue Canadienne de Recherche Forestière* 37: 2575–2586.
- Robroek B. J. M., Limpens J., Breeuwer A., van Ruijven J. & Schouten M. G. C. (2007): Precipitation determines the persistence of hollow *Sphagnum* species on hummocks. – *Wetlands* 27: 979–986.
- Ros R. M. & Cano M. J. (2008): Identity of North African endemic bryophytes, 2. – *Cryptogamie Bryologie* 29: 135–141.
- Ros R. M., Mazimpaka V., Abou-Salama U., Aleffi M., Blockeel T. L., Brugués M., Cano M. J., Cros R. M., Dia M. G., Dirkse G. M., El Saadawi W., Erdağ A., Ganeva A., González-Mancebo J. M., Herrnstadt I., Khalil K., Kürschner H., Lanfranco E., Losada-Lima A., Refai M. S., Rodríguez-Núñez S., Sabovljević M., Sérgio C., Shabbara H., Sim-Sim M. & Söderström L. (2007): Hepatics and Anthocerotes of the Mediterranean, an annotated checklist. – *Cryptogamie Bryologie* 28: 351–437.
- Ros R. M., Muñoz J., Werner O. & Rams S. (2008): New typifications and synonyms in *Tortula* sect. *Pottia* (*Pottiaceae*, *Musci*). – *Taxon* 57: 279–288.
- Rozbrojová Z. & Hájek M. (2008): Changes in nutrient limitation of spring fen vegetation along environmental gradients in the West Carpathians. – *Journal of Vegetation Science* 19: 613–620.

- Rozzi R., Armesto J. J., Goffinet B., Buck W., Massardo F., Silander J., Arroyo M. T. K., Russell S., Anderson C. B., Cavieres L. A. & Callicott J. B. (2008): Changing lenses to assess biodiversity: patterns of species richness in sub-Antarctic plants and implications for global conservation. – *Frontiers In Ecology and the Environment* 6: 131–137.
- Sabovljević M., Vukojević V., Sabovljević A., Mihajlović N., Dražić G. & Vučinić Ž. 2007): Determination of heavy metal deposition in the county of Obrenovac (Serbia) using mosses as bioindicators. III. Copper (Cu), iron (Fe) and mercury (Hg). – *Archives of Biological Sciences* 59: 351–361.
- Salemaa M., Mäkipää R. & Oksanen J. (2008): Differences in the growth response of three bryophyte species to nitrogen. – *Environmental Pollution* 152: 82–91.
- Samecka-Cymerman A., Kolon K. & Kempers A. J. (2008): A preliminary investigation in using *Pohlia nutans* and *Larix decidua* as biomonitor of air pollution by the coke industry in Walbrzych (SW Poland). – *Polish Journal of Environmental Studies* 17: 121–128.
- Savaroglu F. & Potoglu Erkara I. (2008): Observations of spore morphology of some *Pottiaceae* Schimp. species (*Bryophyta*) in Turkey. – *Plant Systematics and Evolution* 271: 93–99.
- Saxena D. K., Srivastava K. & Singh S. (2008): Retrospective metal data of the last 100 years deduced by moss, *Barbula* sp from Mussoorie city, Garhwal Hills, India. – *Current Science* 94: 901–904.
- Schröder W., Pesch R., Englert C., Harmens H., Suchara I., Zechmeister H. G., Thöni L., Mařkovská B., Jeran Z., Grodzinska K. & Alber R. (2008): Metal accumulation in mosses across national boundaries: Uncovering and ranking causes of spatial variation. – *Environmental Pollution* 151: 377–388.
- Selkirk P. M., Whinam J. P., Downing A. J. & Skotnicki M. L. (2008): Mosses of sub-Antarctic Heard Island: an updated list and discussion of their distribution. – *Polar Record* 44: 155–164.
- Shakya K., Chettri M. K. & Sawidis T. (2008): Impact of heavy metals (copper, zinc, and lead) on the chlorophyll content of some mosses. – *Archives of Environmental Contamination and Toxicology* 54: 412–421.
- Shannon J. M. & Morgan J. W. (2007): Floristic variation in *Sphagnum*-dominated peatland communities of the Central Highlands, Victoria. – *Cunninghamia* 10: 59–76.
- Shaw A. J., Cao T., Wang L.-s., Flatberg K. I., Flatberg B., Shaw B., Zhou P., Boles S., & Terracciano S. (2008): Genetic variation in three Chinese peat mosses (*Sphagnum*) based on microsatellite markers, with primer information and analysis of ascertainment bias. – *Bryologist* 111: 271–281.
- Shaw A. J., Holz I., Cox C. J. & Goffinet B. (2008): Phylogeny, character evolution, and biogeography of the gondwanic moss family *Hypopterygiaceae* (*Bryophyta*). – *Systematic Botany* 33: 21–30.
- Shields J. A., Webster C. R. & Glime J. A. (2007): Bryophyte community response to silvicultural opening size in a managed northern hardwood forest. – *Forest Ecology and Management* 252: 222–229.
- Shimamura M., Yamaguchi T. & Deguchi H. (2008): Airborne sperm of *Conocephalum conicum* (*Conocephalaceae*). – *Journal of Plant Research* 121: 69–71.
- Skrzypek G., Kałużyński A., Wojtuń B. & Jędrysek M.-O. (2007): The carbon stable isotopic composition of mosses: A record of temperature variation. – *Organic Geochemistry* 38: 1770–1781.
- Söderström L., Urmi E. & Váňa J. (2007): The distribution of *Hepaticae* and *Anthocerotae* in Europe and Macaronesia - Update 1-427. – *Cryptogamie Bryologie* 28: 299–350.
- Sotiaux A., Steperaere H. & Vanderpoorten A. (2007): Bryophyte checklist and European red list of the Brussels-Capital Region, Flanders and Wallonia (Belgium). – *Belgian Journal of Botany* 140: 174–196.
- Staples G. W. & Imada C. T. (2006) [2007]. Checklist of Hawaiian anthocerotes and hepatics. – *Tropical Bryology* 28: 15–47.
- Startsev N., Lieffers V. J. & Landhäusser S. M. (2008): Effects of leaf litter on the growth of boreal feather mosses: Implication for forest floor development. – *Journal of Vegetation Science* 19: 253–260.
- Stech M. & Frey W. (2008): A morpho-molecular classification of the mosses (*Bryophyta*). – *Nova Hedwigia* 86: 1–21.
- Stech M., Pfeiffer T. & Frey W. (2008): Phylogenetic relationships within *Dendroligotrichum* and *Polytrichadelphus* (*Polytrichaceae*, *Polytrichopsida*) in the Palaeoaustral region. Studies in austral temperate rainforest bryophytes 32. – *Australian Systematic Botany* 21: 39–49.
- Stenøien H. K. (2008): Slow molecular evolution in 18S rDNA, *rbcL* and *nad5* genes of mosses compared with higher plants. – *Journal of Evolutionary Biology* 21: 566–571.
- Stotler R. E. & Crandall-Stotler B. J. (2008): Correct author citations for some upper rank names of liverworts (*Marchantiophyta*). – *Taxon* 57: 289–292.
- Studlar S. M. & Peck J. E. (2007): Commercial moss harvest in the Appalachian Mountains of West Virginia: targeted species and incidental take. – *Bryologist* 110: 752–765.

- Suchara I. & Sucharová J. (2008): Mercury distribution around the Spolana chlor-alkali plant (central Bohemia, Czech Republic) after a catastrophic flood, as revealed by bioindicators. – Environmental Pollution 151: 352–361.
- Suzuki T. & Iwatsuki Z. (2007): A new approach to the infrageneric classification of the genus *Fissidens* (Fissidentaceae, Bryopsida). – Hikobia 15: 67–85.
- Taylor P. J., Eppley S. M. & Jesson L. K. (2007): Sporophytic inbreeding depression in mosses occurs in a species with separate sexes but not in a species with combined sexes. – American Journal of Botany 94: 1853–1859.
- Tomlinson S., Matthes U., Richardson P. J. & Larson D. W. (2008): The ecological equivalence of quarry floors to alvars. – Applied Vegetation Science 11: 73–82.
- Troitsky A. V., Ignatov M. S., Bobrova V. K. & Milyutina I. A. (2007): Contribution of genosystematics to current concepts of phylogeny and classification of bryophytes. – Biochemistry (Moscow) 72: 1368–1376.
- Tsegmed Ts. (2006): Checklist and distributions of the mosses in Mongolia with addition. – Botanikiin Khureelengiin Erdem Shinhilgeenii Buteel 16: 18–40.
- Tusiime F. Mugizi, Byarujali S. M. & Bates J. W. (2007): Diversity and distribution of bryophytes in three forest types of Bwindi Impenetrable National Park, Uganda. – African Journal of Ecology 45, Suppl. 3: 79–87.
- Vanderpoorten A., Devos N., Goffinet B., Hardy O. J. & Shaw A. J. (2008): The barriers to oceanic island radiation in bryophytes: insights from the phylogeography of the moss *Grimmia montana*. – Journal of Biogeography 35: 654–663.
- Vázquez M. D., Fernández J. A., Real C., Villares R., Aboal J. R. & Carballeira A. (2007): Design of an aquatic biomonitoring network for an environmental specimen bank. – Science of the Total Environment 388: 357–371.
- Vilnet A. A., Milyutina I. A., Konstantinova N. A., Ignatov M. S. & Troitsky A. V. (2007): Phylogeny of the genus *Lophozia* (Dumort.) Dumort. s. str. inferred from nuclear and chloroplast sequences ITS1–2 and TRNL–F. – Russian Journal of Genetics 43: 1306–1313.
- Virkutyte J., Vadakojyte S., Sinkevicius S. & Sillanpaa M. (2008): Heavy metal distribution and chemical partitioning in Lake Saimaa (SE Finland) sediments and moss *Pleurozium schreberi*. – Chemistry and Ecology 24: 119–132.
- Wallace J. & McJannet D. (2008): Modelling interception in coastal and montane rainforests in northern Queensland, Australia. – Journal of Hydrology 348: 480–495.
- Wanek W. & Pörtl K. P. (2008): Short-term ^{15}N uptake kinetics and nitrogen nutrition of bryophytes in a lowland rainforest, Costa Rica. – Functional Plant Biology 35: 51–62.
- Weber W. A. & Wittmann R. C. (2007): Bryophytes of Colorado: mosses, liverworts, and hornworts. – Pilgrims Process, Santa Fe, New Mexico, USA. [231 pp.]
- Wickett N. J. & Goffinet B. (2008): Origin and relationships of the myco-heterotrophic liverwort *Cryptothallus mirabilis* Malmb. (*Metzgeriales, Marchantiophyta*). – Botanical Journal of the Linnean Society 156: 1–12.
- Wickett N. J., Zhang Y., Hansen S. K., Roper J. M., Kuehl J. V., Plock S. A., Wolf P. G., dePamphilis C. W., Boore J. L. & Goffinet B. (2008): Functional gene losses occur with minimal size reduction in the plastid genome of the parasitic liverwort *Aneura mirabilis*. – Molecular Biology and Evolution 25: 393–401.
- Xu S.-j., Chen W.-h., Chen Y.-w., He M. & Wang Y. (2007): Effects of thermostress on the plasma membrane permeability of desert moss *Tortula desertorum* examined by in-situ micro-FTIR analysis. – Spectroscopy and Spectral Analysis 27: 2417–2421.
- Yang S.-D., Wang R., Li G. & Egashira K. (2008): Selection of a vegetable water-retaining material and its effects on the growth and quality of cucumber under drought stress part 1: Selection of a vegetable water-retaining material. – Journal of the Faculty of Agriculture Kyushu University 53: 133–136.
- Zander R. H. (2008): Statistical evaluation of the clade “*Rhabdoweisiaceae*”. – Bryologist 111: 292–301.
- Zechmeister H. G., Hohenwallner D., Hanus-Illnar A., Hagendorfer H., Roder I. & Riss A. (2008): Temporal patterns of metal deposition at various scales in Austria during the last two decades. – Atmospheric Environment 42: 1301–1309.
- Zhang Y. & Liang-Dong G. (2007): Arbuscular mycorrhizal structure and fungi associated with mosses. – Mycorrhiza 17: 319–325.
- Zotz G. & Kahler H. (2007): A moss “canopy” – small-scale differences in microclimate and physiological traits in *Tortula ruralis*. – Flora 202: 661–666.

Bryologické publikace z České a Slovenské republiky [Bryological publications issued in the Czech Republic and Slovakia]

1. Články a knižní publikace [Papers and books]

- Čech L., Dvořáčková K. & Juřička J. (eds.) (2006): Seznam zjištěných taxonů mechorostů. – In: Čech L., Dvořáčková K. & Juřička J. (eds.), Výsledky floristického kurzu České botanické společnosti v Jihlavě (2.–8. července 2005) [Results of the floristic summer school of the Czech Botanical Society in Jihlava (2–8 July 2005)], Zprávy České botanické společnosti 41, Suppl. 2006/1, p. 69.
- Duda J. & Duda J. (2005): Doplněk k poznání játrovek pohoří Roháče na Slovensku [Complement to the knowledge of liverwort flora of Roháče Mts, Slovakia]. – Časopis Slezského Zemského Muzea, Ser. A, 54: 161–166.
- Duda J. & Duda J. (2005): Mechiorosty na kopcích na severních svazích Kněhyně a Tanečnice v Moravskoslezských Beskydech [Bryophytes of hills in northern slopes of Kněhyně Mt. and Tanečnice Mt. in Moravskoslezské Beskydy Mts. (Czech Republic)]. – Časopis Slezského Zemského Muzea, Ser. A, 54: 85–89.
- Duda J. & Duda J. (2007): Mechiorosty hory Malý Manín (Slovensko) [Bryophytes of Malý Manín hill (Slovakia)]. – Časopis Slezského Zemského Muzea, Ser. A, 56: 175–176.
- Duda J. & Duda J. (2007): Mechiorosty hory Velký Manín (Slovensko) [Bryophytes of Velký Manín hill (Slovakia)]. – Časopis Slezského Zemského Muzea, Ser. A, 56: 172–174.
- Hájek M. & Hájková P. (2007): Hlavní typy rašeliníšť ve střední Evropě z botanického hlediska [Major mire types in Central Europe from the botanical point of view]. – Zprávy České botanické společnosti 42, Mater. 22: 19–28.
- Hlaváček R., Nesvadbová J., Pecháčková S. & Franklová H. (2008): Botanický průzkum lokality uvažované pro výstavbu radaru u Míšova v Brdech. – Calluna 13/1: 5–7.
- Kochjarová J., Šoltés R. & Hrvnák R. (2007): Mrižkovec pobrežný (*Cinclidotus aquaticus*) na Slovensku: súčasnosť a prognózy [*Cinclidotus aquaticus* in Slovakia: present state and prognoses]. – Bryonora 40: 1–6.
- Košnar J., Paulič R. & Matějková I. (2007): Májová bryologicko-fanerogamologická exkurze do okolí Žichovic. – Calluna 12/1: 16–19.
- Kubešová S. & Novotný I. (2008): Mechiorosty. – In: Skořepa H. (ed.), Přírodní poměry Boskovicka. Vlastivěda Boskovicka, Sv. 1, p. 79–83, Muzeum Boskovicka, Boskovice.
- Kubešová S. (2007): Mechiorosty DP lomu Mokrá a EVL Sivický les. – In: Sborník odborného semináře Monitoring stavu životního prostředí v lomových prostorech, p. 45–46, Brno.
- Kučera J. (ed.) (2007): Zajímavé bryofloristické nálezy X [Interesting bryofloristic records, X]. – Bryonora 40: 41–43.
- Kučera J., Kubešová S., Hájek M. & Plášek V. (2007): Nová bryologická literatura XVIII. [New bryological literature, XVIII]. – Bryonora 40: 53–65.
- Kučera J., Váňa J. & Wojtuń B. (2007): Mechiorosty. – In: Kolektiv autorů, Krkonoše: Příroda – historie – život, p. 201–204, Nakl. Miloš Uhlíř – Baset, Praha.
- Mikulášková E., Berka T., Holá E., Košnar J., Kubešová S., Marková I., Mudrová R. & Musil Z. (2007): Mechiorosty zaznamenané v průběhu 20. podzimního setkání Bryologicko-lichenologické sekce v NP Šumava [Bryophytes recorded during the 20th Autumn Meeting of the Bryological and Lichenological Section in the Šumava National Park (Bohemian Forest)]. – Bryonora 40: 14–27.
- Pecháčková S. & Mudrová R. (2008): Exkurze kolem Kamenného rybníka – tentokrát za mechiorosty. – Calluna 13/1: 1–2.
- Plášek V. & Marková I. (2007): *Orthotrichum pulchellum* (*Orthotrichaceae, Musci*), new to the Czech Republic. – Acta Musei Moraviae, Sci. Biol., 92: 223–228.
- Plášek V. (2007): Mechiorosty Přírodní památky Koryto řeky Ostravice. – Práce a studie Muzea Beskyd, Přírodní vědy, 19: 185–192.
- Plášek V. (2007): Potvrzení výskytu mechu *Orthotrichum alpestre* Hornsch. ex Bruch. & Schimp. v České republice a nově zjištěná historická lokalita tohoto druhu na Slovensku [Confirmation of the occurrence of moss *Orthotrichum alpestre* Hornsch. ex Bruch & Schimp. in the Czech Republic and a newly discovered historical locality in Slovakia]. – Bryonora 40: 27–30.
- Plášek V. (2007): Recent findings of epiphytic mosses from *Orthotrichaceae* family in Poloniny National Park (Slovakia). – Časopis Slezského Zemského Muzea, Ser. A, 56: 277–283.
- Plášek V. (2007): Mechiorosty přírodní rezervace Velký Polom (Moravskoslezské Beskydy) [Bryophytes of Natural Reserve „Velký Polom“ in Moravskoslezské Beskydy Mts.]. – Časopis Slezského Zemského Muzea, ser. A, 56: 167–171.

- Štechová T. & Štech M. (2007): Ohrožené mechorosty rašelinišť České republiky. [Endangered bryophytes of mires in the Czech Republic] – Zprávy České botanické společnosti 42, Mater. 22: 113–117.
- Wierzcholska S., Dajdok Z. & Wuczyński A. (2007): Recent occurrence of moss *Archidium altenifolium* (*Bryophyta, Archidiaceae*) in Lower Silesia (SW Poland). – Časopis Slezského Zemského Muzea, Ser. A, 56: 285–288.

2. Recenze [Reviews]

- Hájek M. (2007): Recenze. – Bryonora 40: 51–52. [Klement J. & Valachovič M. (eds) (2007): Rastlinné spoločenstvá Slovenska 4. Vysokohorská vegetácia. – Veda, Bratislava.]
- Váňa J. (2007): Recenze. – Bryonora 40: 51. [Bednarek-Ochyra H. (2006): A taxonomic monograph of the moss genus *Codriophorus* P. Beauv. (*Grimmiaceae*). – W Szafer Institute of Botany, Polish Academy of Science, Kraków.]

3. Rukopisné práce [Manuscripts]

- Kubešová S., Novotný I. & Sutorý K. (2007): Inventarizační průzkum cévnatých rostlin a mechorostů PR Olšina Sklenné, PR Řeka, PR Stíří důl, PP Peperek, PP Rozštípená skála, PP Tisůvka, PP Brožova skála, Světka, Hudecká skála. – Ms., 77 pp. [depon in: Botanické oddělení, Moravské zemské muzeum, Brno]
- Novotný I. & Dědečková M. (2006): Návrh metodiky pro mech *Hilpertia velenovskyi* (Schiffn.) R.H.Zander [*Bryophyta, Dicranaceae*]. – Ms., 24 pp. [depon in: Botanické oddělení, Moravské zemské muzeum, Brno]
- Novotný I. & Dědečková M. (2006): Návrh metodiky pro mech Děrozub pochybný *Trematodon ambiguus* (Hedw.) Hornsch. [*Bryophyta, Dicranaceae*]. – Ms., 18 pp. [depon in: Botanické oddělení, Moravské zemské muzeum, Brno]
- Novotný I. & Kubešová S. (2005): Inventarizační průzkum bryoflóry NPR Děvín – Kotel – Soutěska. – Ms., 30 pp. [depon. in: Správa CHKO Pálava, Mikulov]
- Novotný I. & Kubešová S. (2005): Inventarizační průzkum bryoflóry NPR Tabulová, Růžový vrch a Kočičí kámen. – Ms., 26 pp. [depon in: Správa CHKO Pálava, Mikulov]
- Novotný I. & Kubešová S. (2005): Inventarizační průzkum mechiorostů NPR Ransko. – Ms., 28 pp. [depon. in: Správa CHKO Žďárské vrchy, Žďár n. S.]
- Zmrhalová M. (2004): Inventarizační průzkum NPR Králický Sněžník. Obor: Bryologie. – Ms., 25 pp. + append. [depon. in: AOPK, Olomouc, Praha]
- Zmrhalová M. (2005): Inventarizační průzkum údolí Bílé Opavy (část NPR Praděd). Obor: bryologie. – Ms., 19 pp. [depon. in: Správa CHKO Jeseníky, Jeseník]

4. Nebryologické práce s údaji o výskytu na území ČR a SR [Non-bryological papers with bryophyte records from the territory of the Czech Republic and Slovakia]

- Bufková I., Prach K. & Bastl M. (2005): Relationships between vegetation and environment within the montane floodplain of the Upper Vltava River (Šumava National Park, Czech Republic). – Silva Gabreta, Suppl. 2: 5–76.
- Hájková P. & Hájek M. (2005): Diversity of *Calthion* wet meadows in the western part of flysch Carpathians: regional classification based on national formal definitions. – Thaiszia 15: 85–116.
- Hofhanzlová E. & Ekrt L. (2006): Floristický a vegetační inventarizační průzkum Národní přírodní rezervace Zhejral [Floristic and vegetation research at the Zhejral reserve]. – Acta Rerum Naturalium 2: 19–37.
- Hofhanzlová E., Ekrt L. & Štechová T. (2005): Floristický a vegetační průzkum rašeliniště Na Klátově [Floristic and vegetation research in the „Na Klátově“ peat bog]. – Acta Rerum Naturalium 1: 45–52.
- Hradílek Z. & Duchoslav M. (2007): Flóra a vegetace Národní přírodní rezervace Žebračka u Přerova [Flora and vegetation of the Žebračka National Nature Reserve near the town of Přerov (Central Moravia, Czech Republic)]. – Časopis Slezského Zemského Muzea, Ser. A, 56: 193–226.
- Strnad L. & Ekrt L. (2007): Nález nové lokality kyvoru lékařského (*Asplenium ceterach*, *Aspleniaceae*, *Pteridophyta*) a přehled jeho rozšíření v České republice. – Zprávy České Botanické Společnosti 42: 221–229.