

NOVÁ BRYOLOGICKÁ LITERATURA XXI.

New bryological literature, XXI

Jan Kučera¹, Svatava Kubešová², Michal Hájek³ & Vítězslav Plášek⁴

¹ Jihoceská Univerzita, Přírodovědecká fakulta, Branišovská 31, CZ-370 05 České Budějovice, e-mail: kucera@prf.jcu.cz; ² Botanické oddělení, Moravské zemské muzeum, Hvězdoslavova 29a, CZ-62700 Brno, skubesova@mzm.cz; ³ Masarykova univerzita, Ústav botaniky a zoologie, Kotlářská 2, CZ-61137 Brno, hajek@sci.muni.cz; ⁴ Ostravská univerzita, KBE PřF, Chittussiho 10, CZ-71000 Ostrava, vitezslav.plasek@osu.cz

Výběr ze světové bryologické literatury [Selection from the world bryological literature]

- Aleffi M. (2008): Biologia ed ecologia delle briofite. – Antonio Delfino Editore, Roma. [xii+434 pp.]
- Aničić M., Tasić M., Frontasyeva M. V., Tomašević M., Rajšić S., Mijić Z. & Popović A. (2009): Active moss biomonitoring of trace elements with *Sphagnum girgensohnii* moss bags in relation to atmospheric bulk deposition in Belgrade, Serbia. – Environmental Pollution 157: 673–679.
- Aničić M., Tasić M., Frontasyeva M. V., Tomašević M., Rajšić S., Strelkova L. P., Popović A. & Steinnes E. (2009): Active biomonitoring with wet and dry moss: a case study in an urban area. – Environmental Chemistry Letters 7: 55–60.
- Anterola A., Shanle E., Mansouri K., Schuette S. & Renzaglia K. (2009): Gibberellin precursor is involved in spore germination in the moss *Physcomitrella patens*. – Planta 229: 1003–1007.
- Ariyanti N. S., Bos M. M., Kartawinata K., Tjitosoedirdjo S. S., Guhardja E. & Gradstein S. R. (2008): Bryophytes on tree trunks in natural forests, selectively logged forests and cacao agroforests in Central Sulawesi, Indonesia. – Biological Conservation 141: 2516–2527.
- Arróniz -Crespo M., Leake J. R., Horton P. & Phoenix G. K. (2008): Bryophyte physiological responses to, and recovery from, long-term nitrogen deposition and phosphorus fertilisation in acidic grassland. – New Phytologist 180: 864–874.
- Asakawa Y. (2008): Liverworts – potential source of medicinal compounds. – Current Pharmaceutical Design 14: 3067–3088.
- Asakawa Y., Ludwiczuk A., Nagashima F., Toyota M., Hashimoto T., Tori M., Fukuyama Y. & Harinantenaina L. (2009): Bryophytes: bio- and chemical diversity, bioactivity and chemosystematics. – Heterocycles 77: 99–150.
- Atwood J. J. (2009): A taxonomic revision of *Schlotheimia* subg. *Stegotheca* (*Orthotrichaceae*). – Bryologist 112: 290–300.
- Bacaro G., Rocchini D., Bonini I., Marignani M., Maccherini S. & Chiarucci A. (2008): The role of regional and local scale predictors for plant species richness in Mediterranean forests. – Plant Biosystems 142: 630–642.
- Bach L. H., Frostegård Å. & Ohlson M. (2009): Site identity and moss species as determinants of soil microbial community structure in Norway spruce forests across three vegetation zones. – Plant and Soil 318: 81–91.
- Banner A. & LePage P. (2008): Long-term recovery of vegetation communities after harvesting in the coastal temperate rainforests of northern British Columbia. – Canadian Journal of Forest Research – Revue Canadienne De Recherche Forestière 38: 3098–3111.
- Bartholomew-Began S. E. (2009): Juvenile gametophyte development in the *Blasiales*. 3. Sporeling ontogeny of *Cavicularia densa*. – Bryologist 112: 354–358.
- Basile A., Sorbo S., Bassi P., Napolitano E., Cogoni A. E. & Cobianchi R. C. (2008): Effects of heavy metals on protonemal development and ultrastructure in populations of the moss *Funaria hygrometrica* Hedw. (*Bryophyta*) from a mine and an unpolluted site. – Fresenius Environmental Bulletin 17: 1956–1963.
- Bates J. W., Wibbelmann M. H., Proctor M. C.F. (2009): Salinity responses of halophytic and non-halophytic bryophytes determined by chlorophyll fluorometry. – Journal of Bryology 31:
- Beever J. E. & Fife A. J. (2008): *Hypnobartlettia fontana* is an environmental form of *Cratoneuropsis relaxa* (*Bryophyte: Amblystegiaceae*). – New Zealand Journal of Botany 46: 341–345.
- Benavides J. C. & Sastre-De Jesús I. (2009): Digitized images provide more accuracy and efficiency to estimate bryophyte cover. – Bryologist 112: 12–18.

- Bengtsson B. O. & Cronberg N. (2009): The effective size of bryophyte populations. – *Journal of Theoretical Biology* 258: 121–126.
- Benscoter B. W. & Vitt D. H. (2008): Spatial patterns and temporal trajectories of the bog ground layer along a post-fire chronosequence. – *Ecosystems* 11: 1054–1064.
- Bergamini A., Peintinger M., Fakheran S., Moradi H., Schmid B. & Joshi J. (2009): Loss of habitat specialists despite conservation management in fen remnants 1995–2006. – *Perspectives In Plant Ecology, Evolution and Systematics* 11: 65–79.
- Bignal K. L., Ashmore M. R. & Headley A. D. (2008): Effects of air pollution from road transport on growth and physiology of six transplanted bryophyte species. – *Environmental Pollution* 156: 332–340.
- Boiko M. F. (2008): Cheklist mokhopodibnykh Ukrayiny [A Checklist of the Bryobionta of Ukraine]. – Ajlant, Kherson. [232 pp.]
- Botting R. S., Campbell J., Fredeen A. L. (2008): Contrasting arboreal and terrestrial macrolichen and bryophyte communities in old-growth sub-boreal spruce forests of central British Columbia. – *Bryologist* 111: 607–619.
- Bragazza L. (2008): A climatic threshold triggers the die-off of peat mosses during an extreme heat wave. – *Global Change Biology* 14: 2688–2695.
- Brugués M., Puche F., Cezón K. *Tortella bambbergeri* (*Pottiaceae*) in the Iberian Peninsula, with an updated key to Iberian *Tortella*. – *Bryologist* 112: 164–168.
- Callaghan D. A. (2008): Scale dependency and area of occupancy: *Tortula freibergii* in north-west England. – *Journal of Bryology* 30: 279–282.
- Campioli M., Samson R., Michelsen A., Jonasson S., Baxter R. & Lemeur R. (2009): Nonvascular contribution to ecosystem NPP in a subarctic heath during early and late growing season. – *Plant Ecology* 202: 41–53.
- Cano M. J. (2008): Taxonomic revision of *Hennediella* Paris (*Pottiaceae, Bryophyta*). – *Bryophytorum Bibliotheca* 64: 1–142.
- Cao T., Wang M., An L., Yu Y., Lou Y., Guo S., Zuo B., Liu Y., Wu J., Cao Y. & Zhu Z. (2009): Air quality for metals and sulfur in Shanghai, China, determined with moss bags. – *Environmental Pollution* 157: 1270–1278.
- Casas C., Brugués M., Cros R. M., Ruiz E. & Barrón A. (2009): Checklist of Mosses of the Spanish Central Pyrenees. – *Cryptogamie Bryologie* 30: 33–65.
- Cesa M., Azzalini G., De Toffol V., Fontanive M., Fumagalli F., Nimis P. L. & Riva G. (2009): Moss bags as indicators of trace element contamination in Pre-alpine streams. – *Plant Biosystems* 143: 173–180.
- Cevik U. & Celik N. (2009): Ecological half-life of ^{137}Cs in mosses and lichens in the Ordu province, Turkey by Cevik and Celik. – *Journal of Environmental Radioactivity* 100: 23–28.
- Clarke L. J., Ayre D. J. & Robinson S. A. (2009): Genetic structure of East Antarctic populations of the moss *Ceratodon purpureus*. – *Antarctic Science* 21: 51–58.
- Cleavitt N. L., Williams S. A. & Slack N. G. (2009): Relationship of bryophyte occurrence to rock type in upstate New York and coastal Maine. – *Northeastern Naturalist* 16: 67–84.
- Coskun M., Steinnes E., Coskun M. & Cayir A. (2009): Comparison of epigeic moss (*Hypnum cupressiforme*) and lichen (*Cladonia rangiformis*) as biomonitor species of atmospheric metal deposition. – *Bulletin of Environmental Contamination and Toxicology* 82: 1–5.
- Crandall-Stotler B., Stotler R. E. & Long D. (2009): Phylogeny and classification of the *Marchantiophyta*. – *Edinburgh Journal of Botany* 66: 155–198.
- Crawford M., Jesson L. K. & Garnock-Jones P. J. (2009): Correlated evolution of sexual system and life-history traits in mosses. – *Evolution* 63: 1129–1142.
- da Costa D. P. (2008): *Metzgeriaceae* (*Hepaticae*). – *Flora Neotropica Monograph* 102: 1–169.
- Dazy M., Masfaraud J.-F. & Ferard J.-F. (2009): Induction of oxidative stress biomarkers associated with heavy metal stress in *Fontinalis antipyretica* Hedw. – *Chemosphere* 75: 297–302.
- Delgadillo C. M. (2009): Floristic corridors for moss distribution across the Neovolcanic Belt of Mexico. IV. The Toluca and Chalco corridors. – *Journal of Bryology* 31: 30–40.
- Demars B. O. L. & Edwards A. C. (2008): Tissue nutrient concentrations in aquatic macrophytes: comparison across biophysical zones, surface water habitats and plant life forms. – *Chemistry and Ecology* 24: 413–422.
- Devos N. & Vanderpoorten A. (2009): Range disjunctions, speciation, and morphological transformation rates in the liverwort genus *Leptoscyphus*. – *Evolution* 63: 779–792.
- Dickson J. H., Hofbauer W., Porley R., Schmidl A., Kofler W. & Oeggl K. (2009): Six mosses from the Tyrolean Iceman's alimentary tract and their significance for his ethnobotany and the events of his last days. – *Vegetation History and Archaeobotany* 18: 13–22.

- Djamali M., Kuerschner H., Akhani H., de Beaulieu J.-L., Amini A., Andrieu-Ponel V., Ponel P. & Stevens L. (2008): Palaeoecological significance of the spores of the liverwort *Riella* (*Riellaceae*) in a late Pleistocene long pollen record from the hypersaline Lake Urmia, NW Iran. – Review of Palaeobotany and Palynology 152: 66–73.
- dos Santos N. D. & da Costa D. P. (2008): The importance of Private Natural Heritage Reserves for conservation of Atlantic rain forest bryoflora: a study at El Nagual, Mage, Rio de Janeiro State, Brazil. – Acta Botanica Brasilica 22: 359–372.
- Draper I. & Hedenäs L. (2008): *Sciuro-hypnum tromsoense* (Kaurin & Arnell) Draper & Hedenäs, a distinct species from the European mountains. – Journal of Bryology 30: 271–278.
- Draper I. & Hedenäs L. (2009): *Sciuro-hypnum dovrense* (Limpr.) Draper et Hedenäs comb. nov., a distinct Eurasian alpine species. – Cryptogamie Bryologie 30: 289–299.
- Dulger B., Hacioglu N. & Uyar G. (2009): Evaluation of antimicrobial activity of some mosses from Turkey. – Asian Journal of Chemistry 21: 4093–4096.
- Dynesius M., Åström M. & Nilsson C. (2008): Microclimatic buffering by logging residues and forest edges reduces clear-cutting impacts on forest bryophytes. – Applied Vegetation Science 11: 345–354.
- Dynesius M., Hylander K. & Nilsson C. (2009): High resilience of bryophyte assemblages in streamside compared to upland forests. – Ecology 90: 1042–1054.
- Eppinga M. B., Rietkerk M., Wassen M. J. & De Ruiter P. C. (2009): Linking habitat modification to catastrophic shifts and vegetation patterns in bogs. – Plant Ecology 200: 53–68.
- Erdağ A. & Kürschner H. (2009): *Cinclidotus vardaranus* Erdağ & Kurschner (*Bryopsida*, *Pottiaceae*) sp. nov. from Eastern Turkey, with some remarks on the speciation centre of the genus. – Nova Hedwigia 88: 183–188.
- Erzberger P. & Schröder W. (2008): The genus *Schistidium* (*Grimmiaceae*, *Musci*) in Hungary. – Studia Botanica Hungarica 39: 27–88.
- Ewald J. (2008): Plant species richness in mountain forests of the Bavarian Alps. – Plant Biosystems 142: 594–603.
- Fenton N. J. & Bergeron Y. (2008): Does time or habitat make old-growth forests species rich? Bryophyte richness in boreal *Picea mariana* forests. – Biological Conservation 141: 1389–1399.
- Fischer E., Killmann D. & Buchbender V. (2008): Zum Status von *Notothylas orbicularis* und *Anthoceros neesii* (*Anthocerotopsida*) im Westerwald/Rheinland-Pfalz. – Decheniana 161: 33–39.
- Frahm J.-P. (2009): Zur Unterscheidung von *Thamnobryum subserratum* (Hook. ex Harv.) Nog. & Z. Iwats. und *Thamnobryum neckeroides* (Hook.) E. Lawton. – Archive for Bryology 39: 1–7.
- Frahm J.-P. (2009): A preliminary study of the infraspecific taxa of *Hypnum cupressiforme* in Europe. – Archive for Bryology 40: 1–10.
- Frahm J.-P. (2009): *Marchantia paleacea* erstmalig in Deutschland gefunden. – Archive for Bryology 47: 1–3.
- Frahm J.-P. & Lüth M. (2008): The Bryophyte Flora of the Maltese Islands. – Archive for Bryology 29: 1–10.
- Frahm J.-P., Lüth M. & Van Melick H. (2008): Kommentierte Artenliste der Moose von Sardinien. – Archive for Bryology 31: 1–13.
- Fritz K. M., Glime J. M., Hribljan J. & Greenwood J. L. (2009): Can bryophytes be used to characterize hydrologic permanence in forested headwater streams? – Ecological Indicators 9: 681–692.
- Fritz O. (2009): Vertical distribution of epiphytic bryophytes and lichens emphasizes the importance of old beeches in conservation. – Biodiversity and Conservation 18: 289–304.
- Fritz O., Niklasson M. & Churski M. (2009): Tree age is a key factor for the conservation of epiphytic lichens and bryophytes in beech forests. – Applied Vegetation Science 12: 93–106.
- Fudali E., Szczepański M., Rusińska A., Rosadziński S. & Wolski G. (2009): The current distribution in Poland of some European neophytic bryophytes with supposed invasive tendencies. – Acta Societatis Botanicorum Poloniae 78: 73–80.
- García-Avila D., De Luna E., Newton A. E. (2009): Phylogenetic relationships of the *Thuidiaceae* and the non-monophyly of the *Thuidiaceae* and the *Leskeaceae* based on *rbcL*, *rps4* and the *rps4-trnS* intergenic spacer. – Bryologist 112: 80–93.
- Gensel P. G. (2008): The earliest land plants. – Annual Review of Ecology Evolution and Systematics 39: 459–477.
- Goffinet B. & Shaw A. J. (eds.) (2009) [2008]: Bryophyte Biology, second edition. – Cambridge University Press, Cambridge. [xiv + 565 pp.]
- Gorski P. (2009): The effects of hikers' paths on the distribution of liverworts in the Tatra Mountains (Western Carpathians). – Cryptogamie Bryologie 30: 229–242.

- Graf M. D. & Rochefort L. (2008): Techniques for restoring fen vegetation on cut-away peatlands in North America. – *Applied Vegetation Science* 11: 521–528.
- Gradstein S. R. (2008): Epiphytes and deforestation in the tropics. – *Abhandlungen aus dem Westfälischen Museum für Naturkunde* 70: 417–424.
- Granath G., Strengbom J., Breeuwer A., Heijmans M. M. P. D., Berendse F. & Rydin H. (2009): Photosynthetic performance in *Sphagnum* transplanted along a latitudinal nitrogen deposition gradient. – *Oecologia* 159: 705–715.
- Guerra J., Jiménez-Martínez J. F., Cano M. J. & Brugués M. (2008): Molecular and morphological evidence on the status of *Bryum mildeanum* Jur. and records revision in the Iberian Peninsula. – *Journal of Bryology* 30: 297–302.
- Haig D. (2008): Homologous versus antithetic alternation of generations and the origin of sporophytes. – *Botanical Review* 74: 395–418.
- Hájek T. & Adamec L. (2009): Mineral nutrient economy in competing species of *Sphagnum* mosses. – *Ecological Research* 24: 291–302.
- Hájek T., Tuittila E.-S., Ilomets M. & Laiho R. (2009): Light responses of mire mosses - a key to survival after water-level drawdown?. – *Oikos* 118: 240–250.
- Hájková P., Hájek M. & Kintrová K. (2009): How can we effectively restore species richness and natural composition of a *Molinia*-invaded fen?. – *Journal of Applied Ecology* 46: 417–425.
- Hallingbäck, T., Carlsson P., Johansson P., Kristensson G. & Sögård P. (eds.) (2008): Göteborgstraktens Mossor. – Föreningen Mossornas Vänner, Uppsala.[128 pp. + CD]
- Harpel J. A. (2009): Wilfred Borden Schofield (1927–2008). – *Bryologist* 112: 257–267.
- He Si (ed.) (2008): Moss Flora of China, English Version. Volume 7, *Amblystegiaceae* to *Plagiotheciaceae*. – Science Press & Missouri Botanical Garden, Beijing, New York & St. Louis. [vii + 258 pp.]
- Hedenäs L. (2009): Relationships among arctic and non-arctic haplotypes of the moss species *Scorpidium cossonii* and *Scorpidium scorpioides* (*Calliergonaceae*). – *Plant Systematics and Evolution* 277: 217–231.
- Heras P. P., Infante S. M. & Izaguirre L. M. (2009): On the use of mosses in the building of a XVth century ship in Northern Spain. – *Cryptogamie Bryologie* 30: 177–184.
- Hock Z., Szövényi P., Schneller J. J., Urmi E. & Tóth Z. (2009): Population genetic consequences of the reproductive system in the liverwort *Mannia fragrans*. – *Plant Ecology* 202: 123–134.
- Hodgetts N. G. (2008): A morphological revision of the genus *Herbertus* S. Gray (*Herbaceae*, *Marchantiophyta*) in Africa, including the East African islands. – *Journal of Bryology* 30: 239–263.
- Hollingsworth M. L., Clark A. A., Forrest L. L., Richardson J., Pennington R. T., Long D. G., Cowan R., Chase M. W., Gaudeul M. & Hollingsworth P. M. (2009): Selecting barcoding loci for plants: evaluation of seven candidate loci with species-level sampling in three divergent groups of land plants. – *Molecular Ecology Resources* 9: 439–457.
- Homen N. & Gabriel R. (2008) Briófitos raros dos Açores/Azorean rare bryophytes. – Princípia Editora, Lda., Parede. [95 pp]
- Hughes P. D. M., Lomas-Clarke S. H., Schulz J. & Barber K. E. (2008): Decline and localized extinction of a major raised bog species across the British Isles: evidence for associated land-use intensification. – *Holocene* 18: 1033–1043.
- Hylander K. & Nemomissa S. (2008): Home garden coffee as a repository of epiphyte biodiversity in Ethiopia. – *Frontiers In Ecology and the Environment* 6: 524–528.
- Hylander K. & Nemomissa S. (2009): Complementary roles of home gardens and exotic tree plantations as alternative habitats for plants of the Ethiopian montane rainforest. – *Conservation Biology* 23: 400–409.
- Hylander K. (2009): No increase in colonization rate of boreal bryophytes close to propagule sources. – *Ecology* 90: 160–169.
- Iserman M. (2007): Diversity of bryophytes in an urban area of NW Germany. – *Lindbergia* 33: 75–82.
- Jeschke M., Kiehl K., Pfadenhauer J. & Gigon A. (2008): Long-term effects of former management on the diversity of vascular plants, mosses and lichens in a calcareous grassland. – *Botanica Helvetica* 118: 95–109.
- Karlin E. F., Boles S. B., Ricca M., Temsch E. M., Greilhuber J. & Shaw A. J. (2009): Three-genome mosses: complex double allopolyploid origins for triploid gametophytes in *Sphagnum*. – *Molecular Ecology* 18: 1439–1454.
- Kauserud H., Mathiesen C. & Ohlson M. (2008): High diversity of fungi associated with living parts of boreal forest bryophytes. – *Botany – Botanique* 86: 1326–1333.
- Klanderud K. & Totland O. (2008): Diversity-stability relationships of an alpine plant community under simulated environmental change. – *Arctic Antarctic and Alpine Research* 40: 679–684.

- Koch K., Frahm J.-P. & Pollawatn R. (2009): The cuticle of the *Buxbaumia viridis* sporophyte. – Flora 204: 34–39.
- Köckinger H., Suanjak M., Schriebl A. & Schröck C. (2008): Die Moos Kärntens. Sonderreihe Natur Kärnten, Band 4. – Verlag der Naturwissenschaftlichen Vereins für Kärnten, Klagenfurt. [319 pp.]
- Kotelko R., Doering M. & Piercy-Normore M. D. (2008): Species diversity and genetic variation of terrestrial lichens and bryophytes in a boreal jack pine forest of central Canada. – Bryologist 111: 594–606.
- Krzaczkowski L., Wright M. & Gairin J. E. (2008): Les Bryophytes, source potentielle de médicaments de demain [Bryophytes, a potent source of drugs for tomorrow's medicine]? – Médecine Sciences: M/S 24: 947–953.
- Lai M.-J., Zhu R.-L. & Chantanaorrapint S. (2008): Liverworts and hornworts of Thailand: an updated checklist and bryofloristic accounts. – Annales Botanici Fennici 45: 321–341.
- Lang D., Zimmer A. D., Rensing S. A. & Reski R. (2008): Exploring plant biodiversity: the *Physcomitrella* genome and beyond. – Trends In Plant Science 13: 542–549.
- Lang S. I., Cornelissen J. H. C., Hölzer A., ter Braak C. J. F., Ahrens M., Callaghan T. V. & Aerts R. (2009): Determinants of cryptogam composition and diversity in *Sphagnum*-dominated peatlands: the importance of temporal, spatial and functional scales. – Journal of Ecology 97: 299–310.
- Lara F., Garilleti R., Medina R. & Mazimpaka V. (2009): A new key to the genus *Orthotrichum* Hedw. in Europe and the Mediterranean Region. – Cryptogamie Bryologie 30: 129–142.
- Larraín J., Herrera F., Budke J. M. & Goffinet B. (2009): Phylogenetic affinities and conservation status of the Chilean endemic *Costesia spongiosa* (*Gigaspermaceae*). – Bryologist 112: 278–286.
- Limpens J., Robroek B. J. M., Heijmans M. M. P. D. & Tomassen H. B. M. (2008): Mixing ratio and species affect the use of substrate-derived CO₂ by *Sphagnum*. – Journal of Vegetation Science 19: 841–848.
- Liu X.-Y., Xiao H.-Y., Liu C.-Q., Li Y.-Y. & Xiao H.-W. (2008): Atmospheric transport of urban-derived NH_x: Evidence from nitrogen concentration and δ¹⁵N in epilithic mosses at Guiyang, SW China. – Environmental Pollution 156: 715–722.
- Lo Giudice R. & Bonanno G. (2009): Bioaccumulation of heavy metals in mosses from Etna Volcano and Iblei Mountains (Eastern Sicily, Italy). – Cryptogamie Bryologie 30: 143–155.
- Löbel S., Snäll T. & Rydin H. (2009): Mating system, reproduction mode and diaspore size affect metacommunity diversity. – Journal of Ecology 97: 176–185.
- Loisel J., Garneau M. & Helie J.-F. (2009): Modern *Sphagnum* δ¹³C signatures follow a surface moisture gradient in two boreal peat bogs, James Bay lowlands, Québec. – Journal of Quaternary Science 24: 209–214.
- Lüth M. (2008): Bildatlas der Moose Deutschlands, Vol. 5:*Andreaeaceae – Timmiaceae*. – Bryologische Arbeitsgemeinschaft Deutschlands, Bonn. [128 color plates]
- Lüth M. (2009): Bildatlas der Moose Deutschlands, Vol. 6:*Amblystegiaceae – Thuidiaceae*. – Bryologische Arbeitsgemeinschaft Deutschlands, Bonn. [106 color plates]
- Mälson K. & Rydin H. (2009): Competitive hierarchy, but no competitive exclusions in experiments with rich fen bryophytes. – Journal of Bryology 31: 41–45.
- Marino P., Raguso R. & Goffinet B. (2009): The ecology and evolution of fly dispersed dung mosses (Family *Splachnaceae*): Manipulating insect behaviour through odour and visual cues. – Symbiosis 47: 61–76.
- Marović G., Franić Z., Senčar J., Bituh T. & Vugrinec O. (2008): Mosses and some mushroom species as bioindicators of radiocaesium contamination and risk assessment. – Collegium Antropologicum 32, Suppl. 2: 109–114.
- Marsteller R. (2008): Die Moosgesellschaften des Verbandes *Fissidention taxifolii* Marst. 2006. 129. Beitrag zur Moosvegetation Thüringens. – Herzogia 21: 199–216.
- Martínez-Abaigar J., Otero S., Tomás R. & Núñez-Olivera E. (2009): Effects of enhanced ultraviolet radiation on six aquatic bryophytes. – Cryptogamie Bryologie 30: 157–175.
- McManus H. A. & Qiu Y.-L. (2008): Life cycles in major lineages of photosynthetic eukaryotes, with a special reference to the origin of land plants. – Fieldiana: Botany, New Series 47: 17–33.
- Meier E. & Paal J. (2009): Cryptogams in Estonian alvar forests: species composition and their substrata in stands of different age and management intensity. – Annales Botanici Fennici 46: 1–20.
- Merced-Alejandro A. & Sastre-De Jesús I. (2009): A developmental sequence for paraphyses in *Neckeropsis* (*Neckeraceae*). – Bryologist 112: 342–353.
- Mewari N. & Kumar P. (2008): Antimicrobial activity of extracts of *Marchantia polymorpha*. – Pharmaceutical Biology 46: 819–822.

- Migaszewski Z. M., Galuszka A., Crock J. G., Lamothe P. J. & Dołęgowska S. (2009): Interspecies and interregional comparisons of the chemistry of PAHs and trace elements in mosses *Hylocomium splendens* (Hedw.) B.S.G. and *Pleurozium schreberi* (Brid.) Mitt. from Poland and Alaska. – Atmospheric Environment 43: 1464–1473.
- Moschen R., Kühl N., Rehberger I. & Lücke A. (2009): Stable carbon and oxygen isotopes in sub-fossil *Sphagnum*: Assessment of their applicability for palaeoclimatology. – Chemical Geology 259: 262–272.
- Müller F. (2008): Neu- und Wiederfunde von Moosen für Sachsen-Anhalt aus dem Nationalpark Harz einschließlich der Lectotypifizierung von *Scapania paludicola*. – Herzogia 21: 251–255.
- Muñoz J., Hespanhol H., Cezón K. & Séneca A. (2009): *Grimmia horrida* (Grimmiaceae), a new species from the Iberian Peninsula. – Bryologist 112: 325–328.
- Novák M., Erel Y., Zemanová L., Bottrell S. H. & Adamová M. (2008): A comparison of lead pollution record in *Sphagnum* peat with known historical Pb emission rates in the British Isles and the Czech Republic. – Atmospheric Environment 42: 8997–9006.
- Núñez-Olivera E., Otero S., Tomás R. & Martínez-Abaigar J. (2009): Seasonal variations in UV-absorbing compounds and physiological characteristics in the aquatic liverwort *Jungermannia exsertifolia* subsp. *cordifolia* over a 3-year period. – Physiologia Plantarum 136: 73–85.
- Ochyra R., Bednarek-Ochyra H. & Smith R. I. L. (2008): New and rare moss species from the Antarctic. – Nova Hedwigia 87: 457–477.
- Ohyama K., Takemura M., Oda K., Fukuzawa H., Kohchi T., Nakayama S., Ishizaki K., Fujisawa M. & Yamato K. (2009): Gene content, organization and molecular evolution of plant organellar genomes and sex chromosomes – Insights from the case of the liverwort *Marchantia polymorpha*. – Proceedings of the Japan Academy, Series B, Physical and Biological Sciences 85: 108–124.
- Olsson S., Rumsey F., Grundmann M., Russell S., Enroth J. & Quandt D. (2009): The origin of the British and Macaronesian endemic *Thamnobryum* species (Neckeraceae). – Journal of Bryology 31: 1–10.
- Otero S., Cezón K., Martínez-Abaigar J. & Núñez-Olivera E. (2008): Ultraviolet-absorbing capacity of aquatic bryophytes from Tierra del Fuego (Argentina). – Journal of Bryology 30: 290–296.
- Pankratov T. A. & Dedysh S. N. (2009): Cellulolytic streptomycetes from *Sphagnum* peat bogs and factors controlling their activity. – Microbiology 78: 227–233.
- Papp B., Erzberger P. & Dragičević S. (2008): Contributions to the bryophyte flora of the Orjen mts, Montenegro. – Studia Botanica Hungarica 39: 101–112.
- Patino J., Gonzalez-Mancebo J. M. & Fernandez-Palacios J. M. (2009): Epiphytic bryophytes in Canarian subtropical montane cloud forests: the importance of the time since disturbance and host identity. – Canadian Journal of Forest Research – Revue Canadienne De Recherche Forestière 39: 48–63.
- Pearman D. A., Preston C. D., Rothero G. P. & Walker K. J. (2008): The Flora of Rum: An Atlantic Island Reserve. – Published by the authors, Truro. [479 pp.]
- Péli E. R., Ötvös E., Juhász A. & Benkő Z. (2008): Effects of elevated air CO₂ concentrations on heavy metal contents of terricolous mosses and lichens: a preliminary study of bioindication under doubled air CO₂ concentration. – Cereal Research Communications 36, Suppl. 5: 2011–2014.
- Pellerin S., Mercure M., Desaulniers A. S. & Lavoie C. (2009): Changes in plant communities over three decades on two disturbed bogs in southeastern Quebec. – Applied Vegetation Science 12: 107–118.
- Pena M. J., Darvill A. G., Eberhard S., York W. S. & O'Neill M. A. (2008): Moss and liverwort xyloglucans contain galacturonic acid and are structurally distinct from the xyloglucans synthesized by hornworts and vascular plants. – Glycobiology 18: 891–904.
- Pereira S. M. P. & Porto K. C. (2009): Effect of fragmentation on the community structure of epixylic bryophytes in Atlantic Forest remnants in the Northeast of Brazil. – Biodiversity and Conservation 18: 317–337.
- Perhans K., Leif A., Jonsson F., Nordin U., Söderström B. & Gustafsson L. (2009): Retention patches as potential refugia for bryophytes and lichens in managed forest landscapes. – Biological Conservation 142: 1125–1133.
- Piccolo G. R. A., da Silva P. & Vital D. M. (2008): *Hyphomycetes* (conidial fungi) associated with decomposing bryophytes. – Acta Botanica Brasilica 22: 599–606.
- Plášek V., Sawicki J., Trávníčková V. & Pasečná M. (2009): *Orthotrichum moravicum* (Orthotrichaceae), a new moss species from the Czech Republic. – Bryologist 112: 329–336.
- Porley R. (2008): Arable bryophytes: a field guide to the mosses, liverworts and hornworts of cultivated land in Britain and Ireland. – WILDGuides Ltd., Old Basing. [140 pp.]
- Pouliquen H., Delepee R., Thorin C., Haury J., Larhantec-Verdier M., Morvan M.- & Le Bris H. (2009): Comparison of water, sediment, and plants for the monitoring of antibiotics: a case study on a river dedicated to fish farming. – Environmental Toxicology and Chemistry 28: 496–502.

- Pressel S., Duckett J. G., Ligrone R. & Proctor M. C. F. (2009): Effects of de- and rehydration in desiccation-tolerant liverworts: a cytological and physiological study. – International Journal of Plant Sciences 170: 182–199.
- Printarakul N., Wongkuna K., Kornochalert S. & Santanachote K. (2009): Database of bryophytes and their ecological parameters in the CMU Herbarium. – Maejo International Journal of Science and Technology 3: 99–129.
- Rajandu E., Kikas K. & Paal J. (2009): Bryophytes and decaying wood in *Hepatica* site-type boreo-nemoral *Pinus sylvestris* forests in Southern Estonia. – Forest Ecology and Management 257: 994–1003.
- Ricca M., Beecher F. W., Boles S. B., Temsch E., Greilhuber J., Karlin E. F. & Shaw A. J. (2008): Cytotype variation and allopolyploidy in north american species of the *Sphagnum subsecundum* complex (*Sphagnaceae*). – American Journal of Botany 95: 1606–1620.
- Rice S. K., Aclander L. & Hanson D. T. (2008): Do bryophyte shoot systems function like vascular plant leaves or canopies? functional trait relationships in *Sphagnum* mosses (*Sphagnaceae*). – American Journal of Botany 95: 1366–1374.
- Robroek B. J. M., Schouten M. G. C., Limpens J., Berendse F. & Poorter H. (2009): Interactive effects of water table and precipitation on net CO₂ assimilation of three co-occurring *Sphagnum* mosses differing in distribution above the water table. – Global Change Biology 15: 680–691.
- Rudolphi J. (2009): Ant-mediated dispersal of asexual moss propagules. – Bryologist 112: 73–79.
- Rzepka A. & Krupa J. (2008): Ca²⁺ reduces the effect of hypoxia in mosses *Mnium undulatum* and *Polytrichum commune*. – Biologia Plantarum 52: 740–742.
- Sabovljević M. & Frahm J.-P. (2008): Genetic structure of the rare and endangered moss *Campylopus oerstedianus* (*Dicranaceae*) in Europe. – Biologia 63: 1073–1077.
- Sabovljević M. & Frahm J.-P. (2009): Genetic diversity and phylogeography of the rare riparian moss *Dichelyma capillaceum* (With.) Myr. inferred from *trnL-F* plastid DNA sequences. – Archives of Biological Sciences 61: 85–91.
- Sabovljević M., Natcheva R., Dihorù G., Tsakiri E., Dragićević S., Erdağ A. & Papp B. (2008): Check-list of the mosses of SE Europe. – Phytologia Balcanica 14: 207–244.
- Sagar T. & Wilson P. (2009): Niches of common bryophytes in a semi-arid landscape. – Bryologist 112: 30–41.
- Samecka-Cymerman A., Stankiewicz A., Kolon K. & Kempers A. J. (2009): Bioindication of trace metals in *Brachythecium rutabulum* around a copper smelter in legnica (Southwest Poland): use of a new form of data presentation in the form of a self-organizing feature map. – Archives of Environmental Contamination and Toxicology 56: 717–722.
- Sawicki J., Nadolski L., Hościk M. & Zieliński R. (2008): Identification of *Conocephalum conicum* and *C. salebrosum* (*Hepaticophyta*) on the base of DNA markers. – Časopis Slezského Zemského Muzea, Ser. A, 57: 259–264.
- Sawicki J., Plášek V. & Szczecińska M. (2009): Molecular evidence does not support the current division of *Orthotrichum* subgenus *Gymnoporus*. – Plant Systematics and Evolution 279: 125–137.
- Schouten M. A., Verweij P. A., Barendregt A., Kleukers R. M. J. C., Kalkman V. J. & de Ruiter P. C. (2009): Determinants of species richness patterns in the Netherlands across multiple taxonomic groups. – Biodiversity and Conservation 18: 203–217.
- Sérgio C. & Garcia C. (2009): Noteworthy range extensions of two *Aneura* (*Jungmanniopsida*, *Metzgeriales*) species new for the Iberian Peninsula: *Aneura maxima* (Schiffn.) Steph. and *A. pseudopinguis* (Herzog) Pócs. – Cryptogamie Bryologie 30: 207–215.
- Shaw B., Terracciano S. & Shaw A. J. (2009): A genetic analysis of two recently described peat moss species, *Sphagnum atlanticum* and *S. bergianum* (*Sphagnaceae*). – Systematic Botany 34: 6–12.
- Shi Y.-q., Zhu C.-j., Yuan H.-q., Li B.-q., Gao J., Qu X.-j., Sun B., Cheng Y.-n., Li S., Li X. & Lou H.-X. (2009): Marchantin C, a novel microtubule inhibitor from liverwort with anti-tumor activity both in vivo and in vitro. – Cancer Letters 276: 160–170.
- Siddiqui K. U., Islam M. A., Ahmed Z. U., Begum Z. N. T., Hassan M. A., Khondker M., Rahman M. M., Kabir S. M. H., Ahmad M., Ahmed A. T. A., Rahman A. K. A. & Haque E. U. (eds.) (2007): Bryophytes, Pteridophytes, Gymnosperms. – Encyclopedia of Flora and Fauna of Bangladesh 5: 1–391.
- Singh A. P. & Nath V. (2007): Hepaticae of Khasi and Jaintia Hills: Eastern Himalayas. – Bishen Singh Mahendra Pal Singh, Dehra Dun. [ix + 382 pp.]
- Skrzypek G., Akagi T., Drzewicki W. & Jędrysek M.-O. (2008): Stable isotope studies of moss sulfur and sulfate from bog surface waters. – Geochemical Journal 42: 481–492.
- Söderström L., Hagborg A., von Konrat M. & Renner M. (2008): Early land plants today: liverwort checklist of checklists. – Fieldiana: Botany, New Series 47: 105–130.

- Sonneitner M., Dullinger S., Wanek W. & Zechmeister H. (2009): Microclimatic patterns correlate with the distribution of epiphyllous bryophytes in a tropical lowland rain forest in Costa Rica. – *Journal of Tropical Ecology* 25: 321–330.
- Spagnuolo V., Terracciano S. & Giordano S. (2009): Trace element content and molecular biodiversity in the epiphytic moss *Leptodon smithii*: Two independent tracers of human disturbance. – *Chemosphere* 74: 1158–1164.
- Spagnuolo V., Terracciano S., Cobianchi R. C. & Giordano S. (2008): Taxonomy of the *Hypnum cupressiforme* complex in Italy based on ITS and *trnL* sequences and ISSR markers. – *Journal of Bryology* 30: 283–289.
- Spitale D., Petraglia A. & Tomaselli M. (2009): Structural equation modelling detects unexpected differences between bryophyte and vascular plant richness along multiple environmental gradients. – *Journal of Biogeography* 36: 745–755.
- Sporn S. G., Bos M. M., Hoffstätter-Müncheberg M., Kessler M. & Gradstein S. R. (2009): Microclimate determines community composition but not richness of epiphytic understory bryophytes of rainforest and cacao agroforests in Indonesia. – *Functional Plant Biology* 36: 171–179.
- Stalheim T., Ballance S., Christensen B. E. & Granum P. E. (2009): Sphagnan – a pectin-like polymer isolated from *Sphagnum* moss can inhibit the growth of some typical food spoilage and food poisoning bacteria by lowering the pH. – *Journal of Applied Microbiology* 106: 967–976.
- Stark Lloyd R., McLetchie D. Nicholas & Roberts S. P. (2009): Gender differences and a new adult eukaryotic record for upper thermal tolerance in the desert moss *Syntrichia caninervis*. – *Journal of Thermal Biology* 34: 131–137.
- Stebel A. & Ochyra R. (eds.) (2008): Bryophytes of the Polish Carpathians. – Sorus, Poznań. [339 pp.]
- Stech M., Sim-Sim M., Esquivel M. G., Fontinha S., Tangney R., Lobo C., Gabriel R. & Quandt D. (2008): Explaining the ‘anomalous’ distribution of *Echinodium* (*Bryopsida: Echinodiaceae*): Independent evolution in Macaronesia and Australasia. – *Organisms Diversity & Evolution* 8: 282–292.
- Strømgård J. & Nordin A. (2008): Commercial forest fertilization causes long-term residual effects in ground vegetation of boreal forests. – *Forest Ecology and Management* 256: 2175–2181.
- Sun S.-Q., Wang D.-Y., He M. & Zhang C. (2009): Monitoring of atmospheric heavy metal deposition in Chongqing, China – based on moss bag technique. – *Environmental Monitoring and Assessment* 148: 1–9.
- Szövényi P., Terracciano S., Ricca M., Giordano S. & Shaw A. J. (2008): Recent divergence, intercontinental dispersal and shared polymorphism are shaping the genetic structure of amphi-Atlantic peatmoss populations. – *Molecular Ecology* 17: 5364–5377.
- Thebaud G. & Petel G. (2008): A contribution to the revision of ombrotrophic and ombrominerotrophic mediterranean mires communities. – *Phytocoenologia* 38: 287–304.
- Thompson D. K. & Waddington J. M. (2008): *Sphagnum* under pressure: towards an ecohydrological approach to examining *Sphagnum* productivity. – *Ecohydrology* 1: 299–308.
- Tipping E., Vincent C. D., Lawlor A. J. & Lofts S. (2008): Metal accumulation by stream bryophytes, related to chemical speciation. – *Environmental Pollution* 156: 936–943.
- Turetsky M. R., Crow S. E., Evans R. J., Vitt D. H. & Wieder R. K. (2008): Trade-offs in resource allocation among moss species control decomposition in boreal peatlands. – *Journal of Ecology* 96: 1297–1305.
- Turnbull J. D. & Robinson S. A. (2009): Accumulation of DNA damage in Antarctic mosses: correlations with ultraviolet-B radiation, temperature and turf water content vary among species. – *Global Change Biology* 15: 319–329.
- Turnbull J. D., Leslie S. J. & Robinson S. A. (2009): Desiccation protects two Antarctic mosses from ultraviolet-B induced DNA damage. – *Functional Plant Biology* 36: 214–221.
- Uribe M. J. (2008): Monograph of *Frullania* subgenus *Meteoriopsis* (*Frullaniaceae, Marchantiophyta*). – *Caldasia* 30: 49–94.
- Uyar G., Avcil E., Ören M., Karaca F. & Öncel M. S. (2009): Determination of Heavy Metal Pollution in Zonguldak (Turkey) by Moss Analysis (*Hypnum cupressiforme*). – *Environmental Engineering Science* 26: 183–194.
- Uyar G., Ören M., Yildirim Y. & Öncel M. S. (2008): Biomonitoring of metal deposition in the vicinity of Eregli steel plant in Turkey. – *Environmental Forensics* 9: 350–363.
- Wang X. Q., Yang P. F., Liu Z., Liu W. Z., Hu Y., Chen H., Kuang T. Y., Pei Z. M., Shen S. H. & He Y. K. (2009): Exploring the mechanism of *Physcomitrella patens* desiccation tolerance through a proteomic strategy. – *Plant Physiology* 149: 1739–1750.

- Werner F. A. & Gradstein S. R. (2009): Diversity of dry forest epiphytes along a gradient of human disturbance in the tropical Andes. – *Journal of Vegetation Science* 20: 59–68.
- Werner O., Patino J., González-Mancebo M. J., de Almeida Gabriel R. M. & Ros R. M. (2009): The taxonomic status and the geographical relationships of the Macaronesian endemic moss *Fissidens luisieri* (*Fissidentaceae*) based on DNA sequence data. – *Bryologist* 112: 315–324.
- Wiedermann M. M., Gunnarsson U., Ericson L. & Nordin A. (2009): Ecophysiological adjustment of two *Sphagnum* species in response to anthropogenic nitrogen deposition. – *New Phytologist* 181: 208–217.
- Wiedermann M. M., Gunnarsson U., Nilsson M. B., Nordin A. & Ericson L. (2009): Can small-scale experiments predict ecosystem responses? An example from peatlands. – *Oikos* 118: 449–456.
- Wilson D., Stock W. D. & Hedderston T. (2009): Historical nitrogen content of bryophyte tissue as an indicator of increased nitrogen deposition in the Cape Metropolitan Area, South Africa. – *Environmental Pollution* 157: 938–945.
- Wu P.-c., Crosby M. R. & He S. (eds.) (2008): Moss Flora of China, English Version. Volume 8, *Sematophyllaceae to Polytrichaceae*. – Science Press & Missouri Botanical Garden, Beijing, New York & St. Louis. [385 pp.]
- Wu Y., Chen Y., Yi Y. & Shen Z. (2009): Responses to copper by the moss *Plagiomnium cuspidatum*: Hydrogen peroxide accumulation and the antioxidant defense system. – *Chemosphere* 74: 1260–1265.
- Xie C.-F., Lou H.-X. (2009): Secondary metabolites in bryophytes: An Ecological Aspect. – *Chemistry & Biodiversity* 6: 303–312.
- Xu S.-jun, Liu C.-j., Jiang P.-a., Cai W.-m. & Wang Y. (2009): The effects of drying following heat shock exposure of the desert moss *Syntrichia caninervis*. – *Science of the Total Environment* 407: 2411–2419.
- Yam R. S. W., O'Hare M., Dunbar M. J., Baattrup-Pedersen A. & Larsen S. E. (2009): River management: use of macrophyte community structure in UK rivers as an example for ecological assessment for hydromorphological impacts. – In: Recent Advances in Water Resources, Hydraulics & Hydrology, Proceedings of the 4th IASME/WSEAS International Conference on Water Resources, Hydraulics and Hydrology, p. 147–151, WSEAS Press.
- Yano O. (2008): Catálogo dos antóceros e hepáticas brasileiros: literatura original, basônimo, localidade-tipo e distribuição geográfica. – *Boletim do Instituto de Botânica* (São Paulo) 19: 1–110.
- Yayintas O. T. & Yapıcı B. M. (2009): In vitro antimicrobial activity of *Brachythecium campestre* and *Eurhynchium pulchellum* extracts. – *Asian Journal of Chemistry* 21: 2193–2197.
- Zechmeister H. G., Richter A., Smidt S., Hohenwallner D., Roder I., Maringer S. & Wanek W. (2008): Total nitrogen content and $\delta^{15}\text{N}$ signatures in moss tissue: indicative value for nitrogen deposition patterns and source allocation on a nationwide scale. – *Environmental Science & Technology* 42: 8661–8667.

Bryologické publikace z České a Slovenské republiky [Bryological publications issued in the Czech Republic and Slovakia]

1. Články a knižní publikace [Papers and books]

- Dolný A. & Plášek V. (2009): *Sphagnum* mosses as a microhabitat for dragonflies bogs (about dragonfly-affinity for *Sphagnum* ecological-group). – In: Bryja J., Řehák Z. & Zukal J. (eds.), Zoologické dny – Brno 2009, Sborník abstraktů, p. 56.
- Drozd P., Ceh Š., Jašík M., Krupař M. & Plášek V. (2009): Kompozice bezobratlých v mechovém patru horských a podhorských lesních ekosystémů. – In: Bryja J., Řehák Z. & Zukal J. (eds.), Zoologické dny – Brno 2009, Sborník abstraktů, p. 52–53.
- Duda J. & Duda J. (2008): Mechiorsty okolí soutoku Bílé a Černé Ostravice v Beskydách. – *Časopis Slezského Zemského Muzea*, Ser. A, 57: 185–187.
- Gąbka M. & Lamentowicz M. (2008): Vegetation-Environment Relationships in Peatlands Dominated by *Sphagnum fallax* in Western Poland. – *Folia Geobotanica* 43: 413–429.
- Gloser J. (2008): Antarktické vegetační oázy. 3. Mechiorsty. – *Živa* 2008/3: 117–120.
- Hradílek Z. & Novotný I. (2008): Mechiorsty [Bryophytes]. – In: Jongepierová I. (ed.), Louky Bílých Karpat [Grasslands of the White Carpathian Mountains], p. 65–70, ZO ČSOP Bílé Karpaty, Veselí nad Moravou.
- Hradílek Z. (2008): *Funaria pulchella* – nový druh pro Českou republiku [*Funaria pulchella*, a new moss species to the Czech Republic]. – *Bryonora* 42: 6–10.
- Kočárek P., Grucmanová Š., Filipcová Z., Bradová L. & Plášek V. (2009): Mosses in the diet of groundhopper *Tetrix tenuicornis* (Orthoptera: Tetrigidae). – In: Bryja J., Řehák Z. & Zukal J. (eds.), Zoologické dny – Brno 2009, Sborník abstraktů, p. 103.

- Kubinská A. (2007): Bryoflóra Zoborských vrchov. – In: Řehořek V. (ed.), Lišajníky, machorasty a cievnaté rastliny Zoborských vrchov, p. 52–61, Slovenská poľnohospodárska univerzita v Nitre, Nitra.
- Kučera J. (ed.) (2008): Zajímavé bryofloristické nálezy XII. [Interesting bryofloristic records, XII]. – Bryonora 42: 38–41.
- Kučera J., Kubešová S., Hájek M. & Plášek V. (2008): Nová bryologická literatura XX. [New bryological literature, XX]. – Bryonora 42: 49–57.
- Kučera T., Peksa O. & Košnar J. (2006): K problematice původu acidofilních borů na Třeboňsku [About the origin of acidophilous Scots Pine forests in the Třeboňsko Basin (South Bohemia)]. – In: Kučera T. & Navrátilová J. (eds.), Biotopy a jejich vegetační interpretace v ČR, p. 91–106, Česká botanická společnost, Praha.
- Mikulášková E. (2008): Diverzita a ekologie šumavských mechovostů. – Živa 2008/6: 249–252.
- Novotný I. (2008): Distribution of *Ditrichum zonatum* (Brid.) Kindb. in the Czech Republic, Slovakia and adjacent areas. – Bryonora 42: 1–5.
- Plášek V. & Cimalová Š. (2008): Zajímavé botanické nálezy z regionu severní Moravy a Slezska II. – Časopis Slezského Zemského Muzea, Ser. A, 57: 271–275.
- Plášek V. & Poklembová P. (2008): Rozšíření a ekologie epifytických mechů z čeledi *Orthotrichaceae* v údolí řeky Bystřice. – Časopis Slezského Zemského Muzea, Ser. A, 57: 155–163.
- Soldán Z. (2009): Species composition and diversity of bryophytes in biological soil crusts on natural substrata. – Novitates Botanica Universitatis Carolinae 19/2008: 25–26.
- Soldán Z. (2009): Species composition and diversity of bryophytes on anthropogenic substrata. – Novitates Botanica Universitatis Carolinae 19/2008: 45–50.
- Štechová T. & Manukjanová A. (2008): Broumarské slatiny – nová lokalita mechu *Hamatocaulis vernicosus* ve východních Čechách. – Orchis 27: 2–5.
- Štechová T., Hájek M., Hájková P. & Navrátilová J. (2008): Comparison of habitat requirements of the mosses *Hamatocaulis vernicosus*, *Scorpidium cossonii* and *Warnstorffia exannulata* in different parts of temperate Europe. – Preslia 80: 399–410.
- Štechová T., Holá E., Štech M. & Mikulášková E. (2007): Recentně známé lokality mechu *Hamatocaulis vernicosus* (Mitt.) Hedenäs v západních Čechách a na Šumavě. – Erica 14: 5–12.

2. Rukopisné práce [Manuscripts]

- Holá E. (2008): Ekologická studie epixylických druhů *Lophozia ascendens* a *Anastrophyllum hellerianum* (*Lophoziacaeae*). – Ms., 39 pp. [MSc. thesis, University of South Bohemia, České Budějovice]
- Novotný I. & Dědečková M. (2008): Návrh metodiky monitoringu pro mech útlovláska bledá *Ditrichum pallidum* (Hedw.) Hampe [Bryophyta, Dicranales]. – Ms., 18 pp. [depon. in: Botanické oddělení, Moravské zemské muzeum, Brno & AOPK, Praha]
- Papáčková D. (2009): Biologie vzácných druhů mechovostů. – Ms., 25 pp. [Bc. thesis, University of South Bohemia, České Budějovice]
- Štechová T. (2008): Výsledky intenzivního monitoringu druhu *Hamatocaulis vernicosus* (*Calliergonaceae*) v roce 2008. – Ms., 89 pp. + append. [depon. in: AOPK ČR, Praha].
- Váňa J. (2007): Mechovosti rašeliníšť v okolí Jizerky (Mechovosti vybraných rašeliníšť Jizerských hor III). – Ms., 34 pp. [depon. in: Správa CHKO Jizerské hory, Liberec].
- Váňa J. (2007): Mechovosti zkoumaných území v Trojské kotlině – stav v roce 2007. – Ms., 5 pp. [depon. in: ŽS-IOŽP-MIS Praha].

3. Nebryologické práce s údaji o výskytu na území ČR a SR [Non-bryological papers with bryophyte records from the territory of CR and SR]

- Bastl M., Burian M., Kučera J., Prach K., Rektorová L. & Štech M. (2008): Central European pine bogs changing along an altitudinal gradient. – Preslia 80: 349–363.
- Bufková I. & Rydlo J. (2008): Vodní makrofyta a mokřadní vegetace odstavených říčních ramen horní Vltavy (Hornovltavský luh, NP Šumava) [Water macrophytes and wetland vegetation in old cut meanders of the Upper Vltava River (Šumava National Park)]. – Silva Gabreta 14: 93–134.
- Ekrlová E., Ekrt L., Košnar J., Zapomělová E. & Čejková A. (2008): Míčovka kulkonosná (*Pilularia globulifera*) znova objevena v České republice [*Pilularia globulifera* rediscovered in the Czech Republic]. – Zprávy České botanické společnosti 43: 193–208.
- Hradílek Z. (2008): Rybníky u Předmostí. Botanický inventarizační průzkum. Závěrečná zpráva. – Ms., 33 pp. (Depon. in: Magistrát města Přerova, odbor životního prostředí).
- Hradílek Z. & Kincl L. (2008): Flóra a vegetace Národní přírodní rezervace Hůrka u Hranic. – Časopis Slezského Zemského Muzea, Ser. A, 57: 237–258.

- Laburďová J. (2008): Vegetace lučních pramenišť západočeské zřídelní oblasti. – Ms., 57 pp. [Bc. Thesis, in Czech, depon. in: University of South Bohemia, České Budějovice].
- Zelený D. (2008): Lesní vegetace v údolí Vltavy severně od Zlaté Koruny (okres Český Krumlov) [Forest vegetation in the Vltava river valley north of Zlatá Koruna (district of Český Krumlov)]. – Zprávy České botanické společnosti 43: 111–169.

NOVÁ LICHENOLOGICKÁ LITERATURA XVIII.

New lichenological literature, XVIII

Zdeněk Palice

Botanický ústav AV ČR, CZ-252 43 Průhonice, e-mail: palice@ibot.cas.cz

- Adamo P., Bargagli R., Giordano S., Modenesi P., Monaci F., Pittao E., Spagnuolo V. & Tretiach M. (2008): Natural and pre-treatments induced variability in the chemical composition and morphology of lichens and mosses selected for active monitoring of airborne elements. – Environmental Pollution 152: 11–19.
- Amtoft A., Lutzoni F. & Miadlikowska J. (2008): *Dermatocarpon (Verrucariaceae)* in the Ozark Highlands, North America. – Bryologist 111: 1–40.
- Aptroot A., Lücking R., Sipman H. J. M., Umaña L. & Chaves J. L. (2008): Pyrenocarpous lichens with bitunicate asci: A first assessment of the lichen biodiversity inventory in Costa Rica. – Bibliotheca Lichenologica 97: 1–162.
- Aptroot A. & Schumm F. (2008): Key to *Ramalina* species known from Atlantic islands, with two new species from the Azores. – Sauteria 15: 21–57.
- Aptroot A. & Stapper N. J. (2008): Flechten im Nationalpark Eifel und in den angrenzenden Ardennen – ein Exkursionsbericht. – Aktuelle Lichenologische Mitteilungen, N.F., 15: 14–43.
- Aragón G., Belinchón R. & Izquierdo P. (2008): Valoración de la diversidad de líquenes epífitos en bosques de quercíneas mediante un nuevo índice liquénico (IDLE). Aplicación a la Red Natura 2000. – Botanica Complutensis, 32: 37–48.
- Armaleo D., Zhang Y. & Cheung S. (2008): Light might regulate divergently depside and depsidone accumulation in the lichen *Parmotrema hypotropum* by affecting thallus temperature and water potential. – Mycologia 100: 565–576.
- Armstrong R. A. (2008): Lichen competition: two-dimensional warfare in slow motion. – Microbiologist 9: 27–31.
- Asplund J. & Gauslaa Y. (2008): Mollusc grazing limits growth and early development of the old forest lichen *Lobaria pulmonaria* in broadleaved deciduous forests. – Oecologia, 155: 93–99.
- Bajgier-Kowalska M. (2008): Lichenometric dating of landslide episodes in the Western part of the Polish Flysch Carpathians. – Catena 72: 224–234.
- Baptista M. S., Vasconcelos M. T. S. D., Cabral J. P., Freitas M. C. & Pacheco A. M. G. (2008): Copper, nickel and lead in lichen and tree bark transplants over different periods of time. – Environmental Pollution 151: 408–413.
- Baruffo L., Picotto M. & Tretiach M. (2008): Intrathalline variation of chlorophyll a fluorescence emission in the epiphytic lichen *Flavoparmelia caperata*. – Bryologist 111: 455–462.
- Basile A., Sorbo S., Aprile G., Conte B. & Castaldo Cobianchi R. (2008): Comparison of the heavy metal bioaccumulation capacity of an epiphytic moss and an epiphytic lichen. – Environmental Pollution 151: 401–407.
- Beiggi S. & Piercy-Normore M. D. (2007): Evolution of ITS ribosomal RNA secondary structures in fungal and algal symbionts of selected species of *Cladonia* sect. *Cladonia* (*Cladoniaceae, Ascomycota*). – Journal of Molecular Evolution 64: 528–542.
- Belinchón R., Martínez I., Escudero A., Aragón G. & Valladares F. (2007): Edge effects on epiphytic communities in a Mediterranean *Quercus pyrenaica* forest. – Journal of Vegetation Science 18: 81–90.
- Benedict J. B. (2008): Experiments on lichen growth, III. The shape of the age-size curve. – Arctic, Antarctic, and Alpine Research 40: 15–26.