

NOVÁ BRYOLOGICKÁ LITERATURA XXV.

New bryological literature, XXV

Jan Kučera¹, Svatava Kubešová² & Vítězslav Plášek³

¹Jihočeská Univerzita, Přírodovědecká fakulta, Branišovská 31, CZ-370 05 České Budějovice, e-mail: kucera@prf.jcu.cz; ²Botanické oddělení, Moravské zemské muzeum, Hvězdovská 29a, CZ-62700 Brno, skubesova@mzm.cz; ³Ostravská univerzita, KBE PřF, Chittussiho 10, CZ-71000 Ostrava, vitezslav.plasek@osu.cz

Výběr ze světové bryologické literatury [Selection from the world bryological literature]

- Adamo P., Giordano S., Sforza A. & Bargagli R. (2011): Implementation of airborne trace element monitoring with devitalised transplants of *Hypnum cupressiforme* Hedw.: assessment of temporal trends and element contribution by vehicular traffic in Naples city. – Environmental Pollution 159: 1620–1628.
- Akiyama H. (2010): Taxonomic revision of the genus *Trismegistia* (*Pylaisiadelphaceae*, *Musci*). – Humans and Nature 21: 1–77.
- Allan Green T. G., Sancho Leopoldo G. & Pintado A. (2011): Ecophysiology of desiccation/rehydration cycles in mosses and lichens. – In: Lüttge U., Beck E. & Bartels D. (eds), Plant Desiccation Tolerance, Ecological Studies Vol. 215, Springer Verlag, p. 89–120.
- Arróniz-Crespo M., Gwynn-Jones D., Callaghan T. V., Núñez-Olivera E., Martínez-Abaigar J., Horton P. & Phoenix G. K. (2011): Impacts of long-term enhanced UV-B radiation on bryophytes in two sub-Arctic heathland sites of contrasting water availability. – Annals of Botany 108: 557–565.
- Asada T., Warner B. & Aravena R. (2005): Effects of the early stage of decomposition on change in carbon and nitrogen isotopes in *Sphagnum* litter. – Journal of Plant Interactions 1: 229–237.
- Asakawa Y. (2011): Bryophytes: chemical diversity, synthesis and biotechnology. A review. – Flavour and Fragrance Journal 26: 318–320.
- Atala C. (2011): Water transport and gas exchange in the non-vascular plant *Dendroligotrichum dendroides* (Brid. ex Hedw.) Broth. (Polytrichaceae, Bryophyta). – Gayana Botanica 68: 89–92.
- Bachtrög D., Kirkpatrick M., Mank J. E., McDaniel S. F., Pires J. C., Rice W. R. & Valenzuela N. (2011): Are all sex chromosomes created equal? – Trends in Genetics 27: 350–357.
- Bakalin V. (2011): Notes on *Lophozia* VI. Taxonomy and distribution of *Lophozia* and *Schistochilopsis* (Lophoziacae) in North America north of Mexico. – Bryologist 114: 298–315.
- Bakalin V. A., Chernyagina O. A. & Kirichenko V. E. (2011): Characteristics of hepatic floras of thermal habitats in Kamchatka. – Contemporary Problems of Ecology 4: 28–34.
- Bartels D. & Hussain S. S. (2011): Resurrection plants: physiology and molecular biology. – In: Lüttge U., Beck E. & Bartels D. (eds), Plant Desiccation Tolerance, Ecological Studies Vol. 215, Springer Verlag, p. 339–364.
- Bartels D., Lüttge U. & Beck E. (2011): Synopsis: drying without dying. – In: Lüttge U., Beck E. & Bartels D. (eds), Plant Desiccation Tolerance, Ecological Studies Vol. 215, Springer Verlag, p. 367–373.
- Baumann M. (2011): Zum Vorkommen von *Tayloria tenuis* im Erzgebirge [On the occurrence of *Tayloria tenuis* in the Ore Mountains (Saxony)]. – Herzogia 24: 103–119.
- Belén Estébanez P., Draper I. D. de Atauri & Medina Bujalance R. (2011): Briófitos: una aproximación a las plantas terrestres más sencillas [Bryophytes: an approximation to the simplest land plants]. – In: Montesinos J. L. V. (ed.), Biodiversidad, Aproximación a la diversidad botánica y zoológica de España, Memorias de la Real Sociedad Española de Historia Natural, 2^a ép., 9, Madrid, p. 19–73.
- Benassi M., Stark L. R., Brinda J. C., McLetchie D. N., Bonine M. & Mishler B. D. (2011): Plant size, sex expression and sexual reproduction along an elevation gradient in a desert moss. – Bryologist 114: 277–288.
- Benavides J. C. & Sastre-De Jesús I. (2011): Diversity and rarity of epiphyllous bryophytes in a superhumid tropical lowland forest of Choco-Colombia. – Cryptogamie Bryologie 32: 119–133.
- Bergmeier E., Blockeel T., Böhling N., Fournaraki C., Gotsiou P., Jahn R., Lansdown R. & Turland N. (2011): An inventory of the vascular plants and bryophytes of Gavdopoula island (S Aegean, Greece) and its phytogeographical significance. – Willdenowia 41: 179–190.
- Bernard E. L. & Schäfer-Verwimp A. (2011): Checklist of the bryophytes of the Guadeloupe archipelago and Martinique (French West Indies). – Cryptogamie Bryologie 32: 233–272.

- Bidartondo M. I., Read D. J., Trappe J. M., Merckx V., Ligrone R. & Duckett J. G. (2011): The dawn of symbiosis between plants and fungi. – *Biology Letters* 7: 574–577.
- Bönsel A. & Sonneck A.-G. (2011): Effects of a hydrological protection zone on the restoration of a raised bog: a case study from Northeast-Germany 1997–2008. – *Wetlands Ecology and Management* 19: 183–194.
- Boquete M. T., Fernández J. A., Aboal J. R. & Carballeira A. (2011): Analysis of temporal variability in the concentrations of some elements in the terrestrial moss *Pseudoscleropodium purum*. – *Environmental and Experimental Botany* 72: 210–216.
- Boquete M. T., Fernández J. A., Aboal J. R. & Carballeira A. (2011): Are terrestrial mosses good biomonitorsof atmospheric deposition of Mn? – *Atmospheric Environment* 45: 2704–2710.
- Brown R. C. & Lemmon B. E. (2011): Spores before sporophytes: hypothesizing the origin of sporogenesis at the algal-plant transition. – *New Phytologist* 190: 875–881.
- Bu Z.-J., Rydin H. & Chen X. (2011): Direct and interaction-mediated effects of environmental changes on peatland bryophytes. – *Oecologia* 166: 555–563.
- Budke J. M., Goffinet B. & Jones C. S. (2011): A hundred-year-old question: is the moss calyptra covered by a cuticle? A case study of *Funaria hygrometrica*. – *Annals of Botany* 107: 1279–1286.
- Burgess K. S., Fazekas A. J., Kesanakurti P. R., Graham S. W., Husband B. C., Newmaster S. G., Percy D. M., Hajibabaei M. & Barrett S. C. H. (2011): Discriminating plant species in a local temperate flora using the *rbcL* plus *matK* DNA barcode. – *Methods in Ecology and Evolution* 2: 333–340.
- Callaghan D. & Bowyer H. (2011): Chemical characteristics of the soil occupied by *Cephaloziella massalongi* and *C. nicholsonii* in Cornwall (UK). – *Journal of Bryology* 33: 173–176.
- Canivet L. & Denayer F.-O. (2011): Staining of protonemal cells of *Ceratodon purpureus* (Hedw.) Brid. for better highlighting micronuclei in bryophytes. – *Cryptogamie Bryologie* 32: 211–220.
- Cano M. J., Guerra J., Jiménez J. A., Gallego M. T. & Orgaz D. J. (2010): An updated bryophytes checklist of the region of Murcia (Southeastern Spain). – *Anales de Biología* 32: 101–131.
- Chang Y. & Graham S. W. (2011): Bryophyte-specific primers for retrieving plastid genes suitable for phylogenetic inference. – *American Journal of Botany* 98: e109–e113.
- Chang Y. & Graham S. W. (2011): Inferring the higher-order phylogeny of mosses (*Bryophyta*) and relatives using a large, multigene plastid data set. – *American Journal of Botany* 98: 839–849.
- Chater C., Kamisugi Y., Movahedi M., Fleming A., Cuming A. C., Gray J. E. & Beerling D. J. (2011): Regulatory mechanism controlling stomatal behavior conserved across 400 million years of land plant evolution. – *Current Biology* 21: 1025–1029.
- Chuah-Petiot M. S. (2011): A checklist of *Hepaticae* and *Anthocerotae* of Malaysia. – *Polish Botanical Journal* 56: 1–44.
- Costa D. P., Pôrto K. C., Luizi-Ponzo A. P., Ilkiu-Borges A. L., Bastos C. J. P., Câmara P. E. A. S., Peralta D. F., Bôas-Bastos S. B. V., Imbabayah C. A. A., Henriques D. K., Gomes H. C. S., Rocha L. M., Santos N. D., Siviero T. S., Vaz-Imbabayah T. F. & Churchill S. P. (2011): Synopsis of the Brazilian moss flora: checklist, distribution and conservation. – *Nova Hedwigia* 93: 277–334.
- Crowley K. F. & Bedford B. L. (2011): Mosses influence phosphorus cycling in rich fens by driving redox conditions in shallow soils. – *Oecologia* 167: 253–264.
- Cushman J. C. & Oliver M. J. (2011): Understanding vegetative desiccation tolerance using integrated functional genomics approaches within a comparative evolutionary framework. – In: Lüttge U., Beck E. & Bartels D. (eds), *Plant Desiccation Tolerance*, Ecological Studies Vol. 215, Springer Verlag, p. 307–338.
- Dandotiya D., Govindappa H., Suman S. & Uniyal P. L. (2011): Checklist of the bryophytes of India. – *Archive for Bryology* 88: 1–126.
- Day M. J. & Currah R. S. (2011): *In vitro* degradation of the moss *Hylocomium splendens* by three pleosporalean fungi. – *Canadian Journal of Microbiology* 57: 382–391.
- Day M. J. & Currah R. S. (2011): Role of selected dark septate endophyte species and other hyphomycetes as saprobes on moss gametophytes. – *Botany – Botanique* 89: 349–359.
- de Carvalho R. C., Branquinho C. & da Silva J. M. (2011): Physiological consequences of desiccation in the aquatic bryophyte *Fontinalis antipyretica*. – *Planta* 234: 195–205.
- De Schrijver A., De Frenne P., Ampoorter E., Van Nevel L., Demey A., Wuyts K. & Verheyen K. (2011): Cumulative nitrogen input drives species loss in terrestrial ecosystems. – *Global Ecology and Biogeography* 20: 803–816.
- Devos N., Renner M. A. M., Gradstein R., Shaw A. J., Laenen B. & Vanderpoorten A. (2011): Evolution of sexual systems, dispersal strategies and habitat selection in the liverwort genus *Radula*. – *New Phytologist* 192: 225–236.

- Dołęgowska S. & Migaszewski Z. M. (2011): PAH concentrations in the moss species *Hylocomium splendens* (Hedw.) B.S.G. and *Pleurozium schreberi* (Brid.) Mitt. from the Kielce area (south-central Poland). – Ecotoxicology and Environmental Safety 74: 1636–1644.
- Eckstein R. L., Pereira E., Milbau A. & Graae B. J. (2011): Predicted changes in vegetation structure affect the susceptibility to invasion of bryophyte-dominated subarctic heath. – Annals of Botany 108: 177–183.
- Evasdotter L. (2011): Rikkärrsrestaurering med grävmaskin – kan man gräva sig till biologisk mångfald? [Successful restoration of a rich fen]. – Svensk Botanisk Tidskrift 105: 94–98.
- Farrant J. M. & Moore J. P. (2011): Programming desiccation-tolerance: from plants to seeds to resurrection plants. – Current Opinion in Plant Biology 14: 340–345.
- Frenzke L., Wanke S., Isnard S., Stoll A., Neinhuis C. & Rowe N. P. (2011): Stem biomechanics of the giant moss *Dendroligotrichum dendroides* s.l. and its significance for growth form diversity in mosses. – Journal of Bryology 33: 229–236.
- Fudali E. (2010): Mosses of Kocioł Łomniczki glacial cirque (Karkonosze Mts) in relation to ecological and phytocoenotical diversity of habitats. – Roczniki Akademii Rolniczej w Poznaniu 389 (Botanika – Steciana 14): 11–17.
- Fudali E. (2010): Mosses of the Kocioł Mały Staw glacial cirque (Karkonosze Mts) and their supposed response to tourism impact in the 20th century. – Acta Botanica Silesiaca 5: 111–130.
- Fudali E. (2010): Zmiany we florze mchów Kotła Łomniczki (Karkonosze) w latach 1869–2008. – Przyroda Sudetów 13: 17–26.
- Gahtori D., Chaturvedi P. & Singh S. (2011): Using bryophytes as a tool to cure European foulbrood disease of honey bee: an eco-friendly approach. – Current Science 101: 420–423.
- Gehrig-Downie C., Obregón A., Bendix J. & Gradstein S. R. (2011): Epiphyte biomass and canopy microclimate in the tropical lowland cloud forest of French Guiana. – Biotropica 43: 591–596.
- Godfrey M. (2011): Some chromosome counts for the genus *Cephaloziella*. – Field Bryology 105: 10–11.
- Goffinet B., Budke J. M. & Newman L. C. (2011): *Micromitriaceae*: A new family of highly reduced mosses. – Taxon 60: 1241–1244.
- Gornall J. L., Woodin S. J., Jónsdóttir I. S. & van der Wal R. (2011): Balancing positive and negative plant interactions: how mosses structure vascular plant communities. – Oecologia 166: 769–782.
- Górski P. & Váňa J. (2011): *Gymnomitrion adustum* – a liverwort new to Slovakia and Poland found in the Tatra Mts (Western Carpathians). – Cryptogamie Bryologie 32: 279–284.
- Greven H. (2011): Ecology and distribution of *Grimmia crinita* Brid. – Field Bryology 104: 18–21.
- Gudmundsdottir R., Olafsson J. S., Palsson S., Gislason G. M. & Moss B. (2011): How will increased temperature and nutrient enrichment affect primary producers in sub-Arctic streams? – Freshwater Biology 56: 2045–2058.
- Guerra J., Jiménez-Martínez J. F., Cano M. J. & Jiménez-Fernández J. A. (2011): A contribution to the phylogenetic study of *Mielichhoferiaceae-Mniaceae* (Bryophyta) based on molecular sequence data. – Nova Hedwigia 93: 47–56.
- Gundale M. J., Deluca T. H. & Nordin A. (2011): Bryophytes attenuate anthropogenic nitrogen inputs in boreal forests. – Global Change Biology 17: 2743–2753.
- Guo S.-L., Zhen Y.-Y., Lou Y.-X. & Shen L. (2011): Genetic background of morphological variation in *Macromitrium cavaleriei* Card. & Ther. based on RAPD. – Plant Science Journal 29: 312–318.
- Hájek M., Roleček J., Cottenie K., Kintrová K., Horská M., Pouličková A., Hájková P., Franková M. & Dítě D. (2011): Environmental and spatial controls of biotic assemblages in a discrete semi-terrestrial habitat: comparison of organisms with different dispersal abilities sampled in the same plots. – Journal of Biogeography 38: 1683–1693.
- Hájková P., Hájek M., Rybníček K., Jiroušek M., Tichý L., Králová S. & Mikulášková E. (2011): Long-term vegetation changes in bogs exposed to high atmospheric deposition, aerial liming and climate fluctuation. – Journal of Vegetation Science 22: 891–904.
- Harmens H., Norris D. A., Cooper D. M., Mills G., Steinnes E., Kubin E., Thöni L., Aboal J. R., Alber R., Carballera A., Coşkun M., De Temmerman L., Frolova M., González-Miqueo L., Jeran Z., Leblond S., Liiv S., Maňkovská B., Pesch R., Poikolainen J., Rühling A., Santamaría J. M., Simonéié P., Schröder W., Suchara I., Yurukova L. & Zechmeister H. G. (2011): Nitrogen concentrations in mosses indicate the spatial distribution of atmospheric nitrogen deposition in Europe. – Environmental Pollution 159: 2852–2860.
- He Q. & Zhu R.-L. (2010): Spore output in 24 Asian bryophytes. – Acta Bryolichenologica Asiatica 3: 125–129.

- He Q. & Zhu R.-L. (2011): Spore output in selected species of *Lejeuneaceae* (*Marchantiophyta*) from China. – *Cryptogamie Bryologie* 32: 107–112.
- Heber U. & Lüttge U. (2011): Lichens and bryophytes: light stress and photoinhibition in desiccation/rehydration cycles – mechanisms of photoprotection. – In: Lüttge U., Beck E. & Bartels D. (eds), *Plant Desiccation Tolerance*, Ecological Studies Vol. 215, Springer Verlag, p. 121–137.
- Hedenäs L. (2011): Relationships among *Cratoneuron curvicaule*, *C. filicinum* var. *filicinum*, and *C. filicinum* var. *atrovirens* (Bryophyta: Amblystegiaceae). – *Journal of Bryology* 33: 99–104.
- Hedenäs L. & Bisang I. (2011): The overlooked dwarf males in mosses – unique among green land plants. – *Perspectives in Plant Ecology, Evolution and Systematics* 13: 121–135.
- Heinrichs J., Kreier H.-P., Feldberg K., Schmidt A. R., Zhu R.-L., Shaw B., Shaw A. J. & Wisseman V. (2011): Formalizing morphologically cryptic biological entities: new insights from DNA taxonomy, hybridization, and biogeography in the leafy liverwort *Porella platyphylla* (Jungermanniopsida, Porellales). – *American Journal of Botany* 98: 1252–1262.
- Heinrichs J., Reiner-Drehwald M. E., Feldberg K., Grimaldi D. A., Nascimbene P. C., von Konrat M. & Schmidt A. R. (2011): *Kaolakia borealis* nov. gen. et sp. (Porellales, Jungermanniopsida): A leafy liverwort from the Cretaceous of Alaska. – *Review of Palaeobotany and Palynology* 165: 235–240.
- Hentschel J. (2011): First confirmed record of *Brachythecium erythrorrhizon* (Brachytheciaceae, Musci) in Germany. – *Herzogia* 24: 159–162.
- Holá E., Kučera J. & Těšitel J. (2011): Comparison of gemma production among three *Lophozia* species during the growing season. – *Flora* 206: 763–768.
- Hugonnot V., Boudier P., Mulot P.-E. & Vergne T. (2011): *Cephalozia macrostachya* Kaal. indicateur de l'état de conservation d'une tourbière en Sologne (Cher, France); son utilité dans la problématique de gestion [*Cephalozia macrostachya* Kaal. indicator of the conservation status of a bog in Sologne (Cher, France); its usefulness in the management issue]. – *Acta Botanica Gallica* 158: 329–345.
- Iurian A. R., Hofmann W., Lettner H., Tuerk R. & Cosma C. (2011): Long term study of Cs-137 concentrations in lichens and mosses. – *Romanian Journal of Physics* 56: 983–992.
- Iwai Y., Murakami K., Gomi Y., Hashimoto T., Asakawa Y., Okuno Y., Ishikawa T., Hatakeyama D., Echigo N. & Kuzuhara T. (2011): Anti-influenza activity of marchantins, macrocyclic bisbibenzyls contained in liverworts. – *Plos One* 6: e19825.
- Jabłońska E., Pawlikowski P., Jarzombkowski F., Chormański J., Okruszko T. & Kłosowski S. (2011): Importance of water level dynamics for vegetation patterns in a natural percolation mire (Rospuda fen, NE Poland). – *Hydrobiologia* 674: 105–117.
- Jaroszewicz B., Sondej I., Stebel A., Pirożnikow E. & Kwiatkowska-Falińska A. J. (2011): Biodiversity of bryophytes growing on the faeces of ungulates – a case study from north-eastern Poland. – *Cryptogamie Bryologie* 32: 221–231.
- Jassey V. E. J., Chiapusio G., Gilbert D., Buttler A., Toussaint M.-L. & Binet P. (2011): Experimental climate effect on seasonal variability of polyphenol/phenoloxidase interplay along a narrow fen-bog ecological gradient in *Sphagnum fallax*. – *Global Change Biology* 17: 2945–2957.
- Jiang Y.-B., Liu X.-H., Tian R.-X. & Shao X.-M. (2011): Field-sampling methods for investigating ground-bryophyte populations in forest vegetation. – *Polish Journal of Ecology* 59: 317–327.
- Jofre J., Goffinet B., Marino P., Raguso R. A., Nihei S. S., Massardo F. & Rozzi R. (2011): First evidence of insect attraction by a Southern Hemisphere *Splachnaceae*: The case of *Tayloria dubyi* Broth. in the Reserve Biosphere Cape Horn, Chile. – *Nova Hedwigia* 92: 317–326.
- Kapfer J., Grytnes J.-A., Gunnarsson U. & Birks H. J. B. (2011): Fine-scale changes in vegetation composition in a boreal mire over 50 years. – *Journal of Ecology* 99: 1179–1189.
- Kayes L. J., Puettmann K. J. & Anderson P. D. (2011): Short-term bryoid and vascular vegetation response to reforestation alternatives following wildfire in conifer plantations. – *Applied Vegetation Science* 14: 326–339.
- Köckinger H. & Kučera J. (2011): *Hymenostylium xerophilum*, sp. nov., and *H. gracillimum*, comb. nov., two neglected European mosses and their molecular affinities. – *Journal of Bryology* 33: 195–209.
- Kophimai Y., Cornejo C., Peintinger M., Bergamini A. & Scheidegger C. (2011): Characterization of nuclear microsatellite loci in the calcareous fen specialist *Scorpidium cossonii* (Calliergonaceae). – *American Journal of Botany* 98: e290–e292.
- Koponen T. J. (2010): Compilation of Musci in *Symbolae sinicae* in the light of H. Handel-Mazzetti's letters to V. F. Brotherus. – *Bryobrothera* 10: 3–78.
- Krause C., Garnica S., Bauer R. & Nebel M. (2011): Aneuraceae (Metzgeriales) and tulasnelloid fungi (Basidiomycota) – a model for early steps in fungal symbiosis. – *Fungal Biology* 115: 839–851.

- Kürschner H. & Frey W. (2011): Liverworts, Mosses and Hornworts of Southwest Asia (Marchantiophyta, Bryophyta, Anthocerotophyta). – Nova Hedwigia, Suppl. [Beiheft] 139: 1–240.
- Lafleur B., Pare D., Fenton N. J. & Bergeron Y. (2011): Growth and nutrition of black spruce seedlings in response to disruption of *Pleurozium* and *Sphagnum* moss carpets in boreal forested peatlands. – Plant and Soil 345: 141–153.
- Lakatos M. (2011): Lichens and Bryophytes: Habitats and Species. – In: Lüttge U., Beck E. & Bartels D. (eds), Plant Desiccation Tolerance, Ecological Studies Vol. 215, Springer Verlag, p. 65–87.
- Lavocat E. & Schäfer-Verwimp A. (2011): Checklist of the bryophytes of the Guadeloupe archipelago and Martinique (French West Indies). – Cryptogamie Bryologie 32: 232–272.
- Lawley M. (2011): The psychology of discovery: finding and recognizing uncommon bryophytes. – Field Bryology 104: 22–27.
- Liebner S., Zeyer J., Wagner D., Schubert C., Pfeiffer E.-M. & Knoblauch C. (2011): Methane oxidation associated with submerged brown mosses reduces methane emissions from Siberian polygonal tundra. – Journal of Ecology 99: 914–922.
- Liere K., Weihe A. & Boerner T. (2011): The transcription machineries of plant mitochondria and chloroplasts: Composition, function, and regulation. – Journal of Plant Physiology 168: 1345–1360.
- Ligrone R. & Duckett J. G. (2011): Morphology versus molecules in moss phylogeny: new insights (or controversies) from placental and vascular anatomy in *Oedipodium griffithianum*. – Plant Systematics and Evolution 296: 275–282.
- Limpens J., Granath G., Gunnarsson U., Aerts R., Bayley S., Bragazza L., Bubier J., Buttler A., van den Berg L. J. L., Francez A.-J., Gerdol R., Grosvernier P., Heijmans M. M. P. D., Hoosbeek M. R., Hotes S., Ilomets M., Leith I., Mitchell E. A. D., Moore T., Nilsson M. B., Nordbakken J.-F., Rochefort L., Rydin H., Sheppard L. J., Thormann M., Wiedermann M. M., Williams B. L. & Xu B. (2011): Climatic modifiers of the response to nitrogen deposition in peat-forming *Sphagnum* mosses: a meta-analysis. – New Phytologist 191: 496–507.
- Liu X.-Y., Xiao H.-Y. & Liu C.-Q. (2011): Physiological and isotopic signals in epilithic mosses for indicating anthropogenic sulfur on the urban-rural scale. – Ecological Indicators 11: 1245–1250.
- Liu Y., Xue J.-Y., Wang B., Li L. & Qiu Y.-L. (2011): The mitochondrial genomes of the early land plants *Treubia lacunosa* and *Anomodon rugelii*: Dynamic and Conservative Evolution. – PLoS ONE: e25836.
- Lloret F. & González-Mancebo J. M. (2011): Altitudinal distribution patterns of bryophytes in the Canary Islands and vulnerability to climate change. – Flora 206: 769–781.
- Lüttge U., Beck E. & Bartels D. (eds) (2011): Plant Desiccation Tolerance. – Ecological Studies, Vol. 215, Springer, Heidelberg. [386 pp.]
- Ma W.-Z. & Liu W.-Y., Song L. (2011): Species composition and distribution of bryophytes on different substrate types in middle mountain moist evergreen broad-leaved forest in Ailao Mountains. – Plant Diversity and Resources 33: 443–450.
- Maciel-Silva A. S. & dos Santos N. D. (2011): Detecting herbivory in two mosses from an Atlantic Forest, Brazil. – Journal of Bryology 33: 140–147.
- Maier E. (2010): The genus *Grimmia* Hedw. (Grimmiaceae, Bryophyta) – a morphological-anatomical study. – Boissiera, Vol. 63, Conservatoire Botanique de Genève. [377 pp.]
- Maier E. (2011): Keys to *Grimmia* in Europe. – Field Bryology 105: 12–31.
- Manninen S., Woods C., Leith I. D. & Sheppard L. J. (2011): Physiological and morphological effects of long-term ammonium or nitrate deposition on the green and red (shade and open grown) *Sphagnum capillifolium*. – Environmental and Experimental Botany 72: 140–148.
- Marschall M. (2010): Photosynthetic responses, carbohydrate composition and invertase activity in fructan accumulating bryophytes (*Porella platyphylla* and *Sphagnum flexuosum*) under different environmental conditions (carbohydrate treatments, dark starvation, low temperature, desiccation). – Acta Biologica Hungarica 61: 120–129.
- Marschall M. & Borbely P. (2011): Photosynthetic responses of the desiccation intolerant *Sphagnum angustifolium* in relation to increasing its desiccation tolerance by exogenous ABA. – Acta Biologica Szegediensis 55: 119–121.
- Martinčič A. (2011): Seznam jetrenjakov (Marchantiophyta) in rogovnjakov (Anthocerotophyta) Slovenije [Annotated checklist of Slovenian liverworts (Marchantiophyta) and hornworts (Anthocerotophyta)]. – Scopia 72: 1–38.
- Masuzaki H., Tsubota H., Shimamura M., Yamaguchi T. & Deguchi H. (2010): A taxonomic revision of the genus *Apometzgeria* (Metzgeriaceae, Marchantiophyta). – Hikobia 15: 427–452.
- Merget B. & Wolf M.. (2010): A molecular phylogeny of Hypnales (Bryophyta) inferred from ITS2 sequence-structure data. – BMC Research Notes 3: 320.

- Migaszewski Z. M., Lamothe P. J., Crock J. G., Gałuszka A. & Dołęgowska S. (2011): The role of sample preparation in interpretation of trace element concentration variability in moss bioindication studies. – *Environmental Chemistry Letters* 9: 323–329.
- Michel P., Burritt D. J. & Lee W. G. (2011): Bryophytes display allelopathic interactions with tree species in native forest ecosystems. – *Oikos* 120: 1272–1280.
- Ming R., Bendahmane A. & Renner S. S. (2011): Sex chromosomes in land plants. – *Annual Review of Plant Biology* 62: 485–514.
- Morris J. L., Richardson J. B. & Edwards D. (2011): Lower Devonian plant and spore assemblages from Lower Old Red Sandstone strata of Tredomen Quarry, South Wales. – *Review of Palaeobotany and Palynology* 165: 183–208.
- Mörsky S. K., Haapala J. K., Rinnan R., Saarnio S., Silvola J., Martikainen P. J. & Holopainen T. (2011): Minor effects of long-term ozone exposure on boreal peatland species *Eriophorum vaginatum* and *Sphagnum papillosum*. – *Environmental and Experimental Botany* 72: 455–463.
- Muller S. D., Daoud-Bouattour A., Belouahem-Abed D., Ben Haj Jilani I., Saad-Limam S. B., Benslama M., Ferchichi-Ben Jamaa H. & Ghrabi-Gammar Z. (2010): Peat mosses (*Sphagnum*) and related plant communities of North Africa. I. The Numidian-Kroumirian Range (Algeria-Tunisia). – *Flora Mediterranea* 20: 159–178.
- Muller S. D., Rhazi L., Saber Er-R., Rifai N., Daoud-Bouattour A., Bottollier-Curte M., Saad-Limam S. B. & Ghrabi-Gammar Z. (2011): Peat mosses (*Sphagnum*) and related plant communities of North Africa. II. The Tingitanean-Rifan range (northern Morocco). – *Nova Hedwigia* 93: 335–352.
- Nan W., Tan B. C. & Davison G. W. H. (2011): Bryophytes as a major dietary component for the Blood Pheasant, *Ithaginis cruentus*. – *Journal of Bryology* 33: 258–260.
- Nomura T. & Hasezawa S. (2011): Regulation of gemma formation in the copper moss *Scopelophila cataractae* by environmental copper concentrations. – *Journal of Plant Research* 124: 631–638.
- Oguri E., Yamaguchi T., Shimamura M., Tsubota H. & Deguchi H. (2010): Phylogenetic relationships among *Leucobryum juniperoides* and related species (Leucobryaceae, Musci) inferred from ITS sequences. – *Hikobia* 15: 453–462.
- Ohtsubo M., Konishi K., Madama H., Higashi T. & Kanayama M. (2011): Retention and biodegradation of oil by peat moss. – *Science Bulletin of the Faculty of Agriculture Kyushu University* 66: 11–20.
- Ochyra R. (2011): A valid description of *Orthotrichum microcarpum* (Orthotrichaceae). – *Polish Botanical Journal* 56: 105–107.
- Ochyra R. & Bednarek-Ochyra H. (2011): Lectotypification of *Hypnum boreale* (Bryopsida), an early name for a bryalean species and its taxonomic status. – *Nova Hedwigia* 93: 525–536.
- Ochyra R., Stebel A. & Bednarek-Ochyra H. (2011): *Grimmia teretinervis* (Grimmiaceae) and *Didymodon validus* (Pottiaceae), two moss species new to Poland. – In: Zemanek B. (ed.), Geobotanist and Taxonomist: A volume dedicated to Professor Adam Zająć on the 70th anniversary of his birth, p. 47–67, Institute of Botany, Jagiellonian University, Cracow, Poland.
- Oishi Y. (2011): Protective management of trees against debarking by deer negatively impacts bryophyte diversity. – *Biodiversity and Conservation* 20: 2527–2536.
- Orgaz D. J., Cano M. J. & Guerra J. (2009): New data on distribution, morphology and habitat of *Brachythecium mildeanum* (Schimp.) Schimp. (Brachytheciaceae, Bryophyta) in the Iberian Peninsula. – *Anales de Biología* 31: 37–41.
- Pajunen A. M., Oksanen J. & Virtanen R. (2011): Impact of shrub canopies on understorey vegetation in western Eurasian tundra. – *Journal of Vegetation Science* 22: 837–846.
- Pan S. & Zhang Z.-H. (2011): Bryophyte communities from abandoned mercury mine in Eastern Guizhou Province. – *Bulletin of Botanical Research* 31: 241–248.
- Papp B., Natcheva R. & Ganeva A. (2011): The bryophyte flora of Northern Mt Strandzha. – *Phytologia Balcanica* 17: 21–32.
- Pejin B., Vujišić L., Sabovljević M., Sabovljević A., Tešević V. & Vajs V. (2011): Fatty acids of some moss species from Germany. – *Asian Journal of Chemistry* 23: 5187–5188.
- Pejin B., Vujišić L., Sabovljević M., Tešević V. & Vajs V. (2011): An insight into fatty acid composition of *Calliergonella cuspidata*. – *Asian Journal of Chemistry* 23: 5161–5162.
- Pérez-Llamazares A., Aboal J. R., Carballeira A. & Fernández J. A. (2011): Cellular location of K, Na, Cd and Zn in the moss *Pseudoscleropodium purum* in an extensive survey. – *Science of the Total Environment* 409: 1198–1204.
- Perroud P.-F., Cove D. J., Quatrano R. S. & McDaniel S. F. (2011): An experimental method to facilitate the identification of hybrid sporophytes in the moss *Physcomitrella patens* using fluorescent tagged lines. – *New Phytologist* 191: 301–306.

- Pisarenko O. Yu., Lapshina E. D. & Mułdiyarov E. Ya. (2011): Cenotic positions and ecological amplitudes of mosses in the vegetation of the Great Vasyugan Swamp. – Contemporary Problems of Ecology 4: 283–295.
- Pokorný L., Oliván G. & Shaw A. J. (2011): Phylogeographic patterns in two Southern Hemisphere species of *Calyptrochaeta* (Doltoniaceae, Bryophyta). – Systematic Botany 36: 542–553.
- Popić J. M., Salbu B., Strand T. & Skipperud L. (2011): Assessment of radionuclide and metal contamination in a thorium rich area in Norway. – Journal of Environmental Monitoring 13: 1730–1738.
- Pouliot R., Rochefort L., Karofeld E. & Mercier C. (2011): Initiation of *Sphagnum* moss hummocks in bogs and the presence of vascular plants: Is there a link? – Acta Oecologica – International Journal of Ecology 37: 346–354.
- Privitera M., Aleffi M., Bertani G., Campisi P., Carratello A., Codogno M., Cogoni A., Miserere L., Perego S., Puglisi M., Sguazzin F., Tacchi R. & Zimbone A. (2010): Survey of the bryophyte diversity of the Carnic Alps, including some interesting records for the Italian bryoflora. – Flora Mediterranea 20: 193–209.
- Reynolds L. A. & McLetchie D. N. (2011): Short distances between extreme microhabitats do not result in ecotypes in *Syntrichia caninervis*. – Journal of Bryology 33: 148–153.
- Ricardi-Branco F., Feria R. S., Jasper A. & Guerra-Sommer M. (2011): *Hepaticites iporangae* n. sp., Rio Bonito Formation, early Permian (Sakmarian), Paraná Basin, Brazil, Western Gondwana. – Journal of Paleontology 85: 360–368.
- Ricca M., Szövényi P., Temsch E. M., Johnson M. G. & Shaw A. J. (2011): Interploidal hybridization and mating patterns in the *Sphagnum subsecundum* complex. – Molecular Ecology 20: 3202–3218.
- Rice S. K., Neal N., Mango J. & Black K. (2011): Relationships among shoot tissue, canopy and photosynthetic characteristics in the feathermoss *Pleurozium schreberi*. – Bryologist 114: 367–378.
- Rico R. R. G. (2011): Methods for cell clearing and rehydration in thalloid liverworts. – Tropical Bryology 33: 12–18.
- Risk A. C., Richardson C. & Davison P. (2011): Epiphyllous bryophytes in the Appalachian Plateau of Kentucky and Tennessee, U.S.A. – Bryologist 114: 289–297.
- Roberge J.-M., Bengtsson S. B. K., Wulff S. & Snall T. (2011): Edge creation and tree dieback influence the patch-tracking metapopulation dynamics of a red-listed epiphytic bryophyte. – Journal of Applied Ecology 48: 650–658.
- Rowntree J. K., Pressel S., Ramsay M. M., Sabovljević A. & Sabovljević M. (2011): *In vitro* conservation of European bryophytes. – In Vitro Cellular & Developmental Biology – Plant 47: 55–64.
- Rudolphi J., Caruso A., von Cräutlein M., Laaka-Lindberg S., Ryömä R. & Berglund H. (2011): Relative importance of thinned and clear-cut stands for bryophyte diversity on stumps. – Forest Ecology and Management 261: 1911–1918.
- Saatkamp A., Aleffi M., de Biaggi M., Delhaye S., Dentant C., Désamoré A., Frahm J.-P., Hébrard J.-P., Laenen B., Leccia M.-F., Lüth M., Miserere L., Offerhaus B., Philippe M. & Tacchi M. (2011): Bryophytes du Parc National du Mercantour : Méthodes et résultats de la campagne d'inventaires 2010 ATBI+M [Bryophytes of Mercantour National Park: methods and results of the 2010 ATBL+M inventory campaign]. – Biocosme Mésogén 28: 33–50.
- Saatkamp A., Guyon M. & Philippe M. (2011): Moss caulking of boats in upper French Rhône and Saône (Eastern France) from the 3rd to the 20th century and the use of *Neckera crispa* Hedwig. – Vegetation History and Archaeobotany 20: 293–304.
- Sabovljević A., Soković M., Glamočlija J., Ćirić A., Vujičić M., Pejin B. & Sabovljević M. (2011): Bioactivities of extracts from some axenically farmed and naturally grown bryophytes. – Journal of Medicinal Plants Research 5: 565–571.
- Sabovljević M., Cvetković J., Živković S., Vujičić M. & Sabovljević A. (2011): Genetic structure of the rare moss species *Rhodobryum ontariense* in Vojvodina (Serbia) as inferred by isozymes. – Archives of Biological Sciences 63: 151–155.
- Savaroğlu F., İşçen C. F., Vatan A. P. Ö., Kabadere S., İlhan S. & Uyar R. (2011): Determination of antimicrobial and antiproliferative activities of the aquatic moss *Fontinalis antipyretica* Hedw. – Turkish Journal of Biology 35: 361–369.
- Sérgio C. & Garcia C. (2011): Bryophyte flora of São Tomé e Príncipe Archipelago (West Africa): annotated catalogue. – Cryptogamie Bryologie 32: 145–196.
- Shanker A., Sharma V. & Daniell H. (2011): Phylogenomic evidence of bryophytes' monophyly using complete and incomplete data sets from chloroplast proteomes. – Journal of Plant Biochemistry and Biotechnology 20: 288–292.

- Shen L., Guo S.-L., Yang W., Cao T. & Glime J. M. (2011): Physiological responses of *Hypnum fertile* Sendtn. (Musci: Hypnaceae) to short-term extreme temperature stress. – Bulletin of Botanical Research 31: 40–48.
- Schmalholz M. & Hylander K. (2011): Microtopography creates small-scale refugia for boreal forest floor bryophytes during clear-cut logging. – Ecography 34: 637–648.
- Sierocka I., Rojek A., Bielewicz D., Karłowski W., Jarmolowski A. & Szwejkowska-Kulinska Z. (2011): Novel genes specifically expressed during the development of the male thalli and antheridia in the dioecious liverwort *Pellia endiviifolia*. – Gene 485: 53–62.
- Sleath J. (2011): The dual process theory as applied to bryological identification. – Field Bryology 105: 32–36.
- Soudzilovskaia N. A., Graae B. J., Douma J. C., Grau O., Milbau A., Shevtsova A., Wolters L. & Cornelissen J. H. C. (2011): How do bryophytes govern generative recruitment of vascular plants? – New Phytologist 190: 1019–1031.
- Spagnuolo V., Zampella M., Giordano S. & Adamo P. (2011): Cytological stress and element uptake in moss and lichen exposed in bags in urban area. – Ecotoxicology and Environmental Safety 74: 1434–1443.
- Staniaszek-Kik M. (2010): Liverworts on decaying wood and tree-fall disturbances in forest communities in the Karkonosze (the Sudetes, SW Poland). – Acta Botanica Silesiaca 5: 131–156.
- Steenbock C. M., Stockey R. A., Beard G. & Tomescu A. M. F. (2011): A new family of leafy liverworts from the middle eocene of Vancouver Island, British Columbia, Canada. – American Journal of Botany 98: 998–1006.
- Stevens C. J., Duprè C., Dorland E., Gaudnik C., Gowing D. J. G., Bleeker A., Diekmann M., Alard D., Bobbink R., Fowler D., Corcket E., Mountford J. O., Vandvik V., Arrestad P. A., Muller S. & Dise N. B. (2011): The impact of nitrogen deposition on acid grasslands in the Atlantic region of Europe. – Environmental Pollution 159: 2243–2250.
- Stewart K. J., Lamb .E. G., Coxson D. S. & Siciliano S. D. (2011): Bryophyte-cyanobacterial associations as a key factor in N₂-fixation across the Canadian Arctic. – Plant and Soil 344: 335–346.
- Street L. E., Subke J. A., Sommerkorn M., Heinemeyer A. & Williams M. (2011): Turnover of recently assimilated carbon in arctic bryophytes. – Oecologia 167: 325–337.
- Sukkharak P., Ludwiczuk A., Asakawa Y. & Gradstein R. (2011): Studies on the genus *Thysananthus* (Marchantiophyta, Lejeuneaceae) 3. Terpenoid chemistry and chemotaxonomy of selected species of *Thysananthus* and *Dendrolejeunea fruticosa*. – Cryptogamie Bryologie 32: 199–209.
- Sun S.-Q., He M., Wang G.-X. & Cao T. (2011): Heavy metal-induced physiological alterations and oxidative stress in the moss *Brachythecium piligerum* Chad. – Environmental Toxicology 26: 453–458.
- Sun S.-Q., Wang G.-X., He M. & Cao T. (2011): Effects of Pb and Ni stress on oxidative stress parameters in three moss species. – Ecotoxicology and Environmental Safety 74: 1630–1635.
- Takezawa D., Komatsu K. & Sakata Y. (2011): ABA in bryophytes: how a universal growth regulator in life became a plant hormone? – Journal of Plant Research 124: 437–453.
- Thouvenot L., Gradstein S. R., Hagborg A., Söderström L. & Bardat J. (2011): Checklist of the liverworts and hornworts of New Caledonia. – Cryptogamie Bryologie 32: 287–390.
- Tuba Z. & Lichtenthaler H. K. (2011): Ecophysiology of homoiochlorophyllous and poikilochlorophyllous desiccation-tolerant plants and vegetations. – In: Lütge U., Beck E. & Bartels D. (eds), Plant Desiccation Tolerance, Ecological Studies Vol. 215, Springer Verlag, p. 157–183.
- Tuba Z., Slack N. G. & Stark L. R. (eds) (2011): Bryophyte Ecology and Climate Change. – Cambridge University Press, Cambridge, U.K. [528 pp.]
- Uğur A., Özden B., Saç M. M. & Yener G. (2011): Biomonitoring of ²¹⁰Po and ²¹⁰Pb using lichens and mosses around coal-fired power plants in Western Turkey. – Journal of Environmental Radioactivity 102: 535–542.
- van der Valk R. (2011): On estimating the duration of phenological stages in bryophytes. – Lindbergia 34: 44–50.
- Váňa J. & Long D. G. (2011): Cephaloziaceae (Marchantiophyta) of the Sino-Himalayan region. – Nova Hedwigia 93: 153–164.
- Vera E. I. (2011): *Livingstonites gabriellae* gen. et sp. nov., permineralized moss (Bryophyta: Bryopsida) from the Aptian Cerro Negro formation of Livingston Island (South Shetland Islands, Antarctica). – Ameghiniana 48: 122–128.
- Vilnet A. A., Konstantinova N. A. & Troitsky A. V. (2011): Taxonomical rearrangements of Solenostomataceae (Marchantiophyta) with description of a new family Endogemmataceae based on *trnL-F* cpDNA analysis. – Folia Cryptogamica Estonica 48: 125–133.

- Waddington J. M., Lucchese M. C. & Duval T. P. (2011): *Sphagnum* moss moisture retention following the re-vegetation of degraded peatlands. – *Ecohydrology* 4: 359–366.
- Wang H., Suo F.-Y., Wang J.-H., Ruxiangul W. & Mamtimin S. (2011): Comparative anatomy of stems and leaves of six species (*Encalypta* Hedw.) from Xinjiang. – *Bulletin of Botanical Research* 31: 524–530.
- Wang J., Lai M. J. & Zhu R. L. (2011): Liverworts and hornworts of Taiwan: an updated checklist and floristic accounts. – *Annales Botanici Fennici* 48: 369–395.
- Wang X.-R., Li M., Zhao J.-C. & Li L. (2011): Spore morphology of 26 species of family Pottiaceae and its systematic significance. – *Bulletin of Botanical Research* 31: 139–146.
- Wang Y.-Y., Wang Y.-F., Wang D.-H., Li M. & Zuo Q. (2011): Effect of habitat fragmentation on bryophytes diversity in the Thousand-Island lake region. – *Plant Science Journal* 29: 435–445.
- Wheeler J. A., Hermanutz L. & Marino P. M. (2011): Feathermoss seedbeds facilitate black spruce seedling recruitment in the forest-tundra ecotone (Labrador, Canada). – *Oikos* 120: 1263–1271.
- Wickett N. J., Forrest L. L., Budke J. M., Shaw B. & Goffinet B. (2011): Frequent pseudogenization and loss of the plastid-encoded sulfate-transport gene *cysA* throughout the evolution of liverworts. – *American Journal of Botany* 98: 1263–1275.
- Wilkie D. & La Farge C. (2011): Bryophytes as heavy metal biomonitor in the Canadian High Arctic. – *Arctic, Antarctic and Alpine Research* 43: 289–300.
- Xiao H.-Y., Tang C.-G., Zhu R.-G., Wang Y.-L., Xiao H.-W. & Liu C.-Q. (2011): Tracing sources of coal combustion using stable sulfur isotope ratios in epilithic mosses and coals from China. – *Journal of Environmental Monitoring* 13: 2243–2249.
- Xie F. & Zhang Z.-H. (2011): Accumulation characteristic of heavy metals between gametophytes and sporophytes of moss *Funaria hygrometrica*. – *Bulletin of Botanical Research* 31: 117–120.
- Yahr R., Coppins B. J. & Ellis C. J. (2011): Preserved epiphytes as an archaeological resource in pre-industrial vernacular buildings. – *Journal of Archaeological Science* 38: 1191–1198.
- Zhilidov A. V., Robarts R. D., Pavlov D. F., Kamari J., Gurtovaya T. Yu., Merilainen J. J. & Pospelov I. N. (2011): Long-term changes of heavy metal and sulphur concentrations in ecosystems of the Taymyr Peninsula (Russian Federation) North of the Norilsk Industrial Complex. – *Environmental Monitoring and Assessment* 181: 539–553.
- Zona D., Oechel W. C., Richards J. H., Hastings S., Kopetz I., Ikawa H. & Oberbauer S. (2011): Light-stress avoidance mechanisms in a *Sphagnum*-dominated wet coastal Arctic tundra ecosystem in Alaska. – *Ecology* 92: 633–644.
- Zotina T. A., Kalacheva G. S. & Bolsunovsky A. Ya (2011): Biochemical fractionation and cellular distribution of americium and plutonium in the biomass of freshwater macrophytes. – *Journal of Radioanalytical and Nuclear Chemistry* 290: 447–451.
- Zuo Q., Higuchi M., Wang Y. F., Arikawa T. & Hirayama Y. (2011): The status of *Struckia* Mull.Hal. (Plagiotheciaceae, Bryopsida) inferred from multiple nuclear and chloroplast loci. – *Journal of Bryology* 33: 221–228.
- Zvereva E. L. & Kozlov M. V. (2011): Impacts of industrial polluters on bryophytes: a meta-analysis of observational studies. – *Water, Air & Soil Pollution* 218: 573–586.

Bryologické publikace z České a Slovenské republiky [Bryological publications issued in the Czech Republic and Slovakia]

1. Články a knižní publikace [Papers and books]

- Cykowska B. & Vončina G. (2011): Recent occurrence of moss *Buxbaumia viridis* (Bryophyta, Buxbaumiaceae) in the Kłodzko region (Central and Eastern Sudetes, SW Poland). – *Časopis Slezského Zemského Muzea*, v. přír., ser. A, 60: 85–89.
- Hradilek Z. & Halda J. P. (2010): Mechiorosty a lišejníky Národní přírodní rezervace Hůrka u Hranic. – *Acta Musei Richnoviensis*, Sect. Natur., 17(2): 29–56.
- Hradilek Z. & Koval Š. (2011): Skrytá krása mechiorostů. – *Naše příroda* 2: 58–65.
- Jónsdóttir I. S. (2011): Diversity of plant life histories in the Arctic. – *Preslia* 83: 281–300.
- Kučera J. (2011): Nová bryologická literatura XXIV [New bryological literature, XXII]. – *Bryonora* 47: 62–80.
- Kučera J. (ed., 2011): Zajímavé bryofloristické nálezy XVII [Interesting bryofloristic records, XVII]. – *Bryonora* 47: 57–59.

- Milewicz M. & Sawicki J. (2011): Molecular identification of sex in dioecious moss *Nyholmiella obtusifolia* (Orthotrichaceae) on the basis of ISSR markers. – Časopis Slezského Zemského Muzea, v. přír., ser. A, 60: 1–6.
- Novotný I., Kubešová S., Doskočilová A., Hradílek Z., Koval Š., Marková I., Musil Z., Plášek V., Uhřeková Šmelková D., Vicherová E. & Zmrhalová M. (2011): Mechorosty zaznamenané v průběhu 17. jarního bryologicko-lichenologického setkání v Chřibech [Bryophytes recorded during the 17th Spring Meeting of the Bryological and Lichenological Section of the ČBS in Chřiby (SE Moravia)]. – Bryonora 47: 1–8.
- Sawicki J., Plášek V. & Szczecińska M. (2011): Recent data on occurrence the sporophytes in populations of dioecious moss, *Nyholmiella obtusifolia* in North-Eastern Poland. – Časopis Slezského Zemského Muzea, v. přír., ser. A, 60: 7–10.
- Štechová T., Manukjanová A. & Čejková A. (2011): Bryoflóra tří rašelinných luk v Orlických horách [Bryophyte flora of three fen meadows in the Orlické hory Mts]. – Bryonora 47: 52–56.
- Váňa J. (2010): Mechorosty západní části Klánovického lesa [Bryophytes of the western part of the Klánovický les forest]. – Natura Pragensis 20: 104–106.
- Váňa J. (2011): Opravy a doplnky k článku „Přínos botaniků 19. století narozených a působících na českém území k nomenklatuře a taxonomii mechanrostů. I. Philipp [Filip] Maximilian Opiz“ [Corrections and additions to the article „Contribution of botanists of 19th century born and working in the area of present Czech Republic to the nomenclature and taxonomy of bryophytes. I. Philipp [Filip] Maximilian Opiz“] Bryonora 47: 27–29.
- Váňa J. (2011): Přínos botaniků 19. století narozených a působících na českém území k nomenklatuře a taxonomii mechanrostů. II. August Karl [Carl] Joseph Corda [Contribution of botanists of 19th century born and working in the area of present Czech Republic to the nomenclature and taxonomy of bryophytes. II. August Karl [Carl] Joseph Corda]. – Bryonora 47: 13–26.
- Vrtalová K., Mátl M., Plaček J., Novotný I. & Zmrhalová M. (2011): Bryologický a lichenologický průzkum v přírodním parku Velký Kosíř na Prostějovsku [Bryological and lichenological investigation in the Nature park of Velký Kosíř close to the town of Prostějov]. – Bryonora 47: 35–40.
- Zmrhalová M. & Koval Š. (2010): Nové poznatky o výskytu mechů šikouška zeleného (*Buxbaumia viridis*) v CHKO Jeseníky. – Campanula. Sborník referátů, konference ke 40. výročí CHKO Jeseníky (1969–2009): 127–130.

2. Rukopisné práce [Manuscripts]

- Jandová J. (2011): Bionomie druhu *Dicranum viride*. – Ms. [Bakalářská práce; depon. in: Jihočeská univerzita v Českých Budějovicích.]
- Manukjanová A. (2011): Vybrané ekologické charakteristiky mechu *Hamatocaulis vernicosus* (Calliergonaceae) [Some chosen ecology characteristics of the moss *Hamatocaulis vernicosus* (Calliergonaceae)]. – Ms. [Diplomová práce; depon. in: Jihočeská univerzita v Českých Budějovicích.]
- Petrželková R. (2011): Biologicky aktivní látky v mechanostech a jejich účinky. – Ms. [Bakalářská práce; depon. in: Knihovna katedry biologie a ekologie, Ostravská Univerzita.]
- Váňa J. (2010): Mechanosti obory v Uhříněvsi (stav v r. 2010). – Ms. [Depon. in: ŽS-IOŽP-MIS Praha.]
- Váňa J. (2010): Mechanosti Pitkovichého údolí (stav v r. 2010). – Ms. [Depon. in: ŽS-IOŽP-MIS Praha.]

3. Nebryologické práce s údaji o výskytu na území ČR a SR [Non-bryological papers with bryophyte records from the territory of CR and SR]

- Dudová L. (2010): Vznik a vývoj rašelinistě Rejvíz a historie okolní krajiny. – Campanula. Sborník referátů, konference ke 40. výročí CHKO Jeseníky (1969–2009): 46–53.
- Hájková P., Roleček J., Hájek M., Horská M., Fajmon K., Polák M. & Jamrichová E. (2011): Prehistoric origin of the extremely species-rich semi-dry grasslands in the Bílé Karpaty Mts (Czech Republic and Slovakia). – Preslia 83: 185–204.
- Halda J., Hauer T., Kociánová M., Mühlsteinová R., Řeháková K. & Šťastná P. (2011): Biodiverzita cévnatých rostlin, lišejníků, sinic a řas na skalách s ledopády v Labském dole [Biodiversity of vacular plants, lichenized fungi, cyanophytes and algae on rocks with icefalls in the Labský důl valley]. – Opera Corcontica 48: 45–68.
- Řezáč M. (2010): Poznámky k vegetaci a květeně západní části Klánovického lesa v Praze [Vegetation and flora of the western part of the Klánovický les forest in Prague]. – Natura Pragensis 20: 107–111.