

## NOVÁ BRYOLOGICKÁ LITERATURA XXIV.

## New bryological literature, XXII

Jan Kučera

Jihočeská Univerzita, Přírodovědecká fakulta, Branišovská 31, CZ–370 05 České Budějovice,  
e-mail: kucera@prf.jcu.cz

## Výběr ze světové bryologické literatury [Selection from the world bryological literature]

- Aboal J. R., Fernández J. A., Boquete T. & Carballeira A. (2010): Is it possible to estimate atmospheric deposition of heavy metals by analysis of terrestrial mosses? – *Science of the Total Environment* 408: 6291–6297.
- Adoli F. K. L., Lalah J. O. & Okoth A. O. (2011): Analysis of moss and topsoil to monitor metal emissions from a pulp and paper mill in Western Kenya. – *Bulletin of Environmental Contamination and Toxicology* 86: 137–143.
- Afonina O. M., Czernyadjeva I. V., Ignatova E. A. & Kučera J. (2010) [2011]: Five species of *Didymodon* (Pottiaceae, Bryophyta) new for Russia. – *Arctoa* 19: 51–62.
- Ahmed Md. G. U. & Lee C. H. (2010): Induction of protonemal gemmae and gametophyte of *Cratoneuron decipiens* (Brid.) G. Roth using IAA and kinetin. – *Plant Omics* 3: 52–56.
- Ah-Peng C., Bardat J., Staménoff P., Hedderson T. A. J. & Strasberg D. (2010): Bryophytes de l'île de La Réunion: diversité, endémicité et conservation. – *Cryptogamie Bryologie* 31: 241–270.
- Akiyama H., Chang Y., Yamaguchi T. & Tan B. C. (2011): *Yakushimabryum longissimum* (Pylaisiadelphaceae) gen. & sp. nov., from the Yakushima Island, Japan. – *Journal of Bryology* 33: 42–49.
- Alboresi A., Caffarri S., Nogue F., Bassi R. & Morosinotto T. (2008): In *silico* and biochemical analysis of *Physcomitrella patens* photosynthetic antenna: identification of subunits which evolved upon land adaptation.. – *Plos One* 3: e2033.
- Alboresi A., Gerotto C., Giacometti G. M., Bassi R. & Morosinotto T. (2010): *Physcomitrella patens* mutants affected on heat dissipation clarify the evolution of photoprotection mechanisms upon land colonization. – *Proceedings of the National Academy of Sciences of the United States of America* 107: 11128–11133.
- Alekseenok Y. V., Frontasyeva M. V. & Korokin A. Z. (2009): Pd, Cu, and Pb Atmospheric Deposition Study in Minsk Region of Belarus based on Moss Analysis and AAS. – *Nuclear Physics Methods and Accelerators In Biology and Medicine* 1204: 229–230.
- Allen B. H. (2010): Moss flora of Central America. Part 3. Anomodontaceae–Symphyodontaceae. – *Monographs in Systematic Botany from the Missouri Botanical Garden* 117: i–x, 1–731.
- Angel Fernández J., Ares Á., Rey-Asensio A., Carballeira A. & Aboal J. R. (2009): Effect of growth on active biomonitoring with terrestrial mosses. – *Journal of Atmospheric Chemistry* 63: 1–11.
- Aranda S. C., Gabriel R., Borges P. A. V. & Lobo J. M. (2010): Assessing the completeness of bryophytes inventories: an oceanic island as a case study (Terceira, Azorean archipelago). – *Biodiversity and Conservation* 19: 2469–2484.
- Ares Á., Angel Fernández J., Aboal J. R. & Carballeira A. (2011): Study of the air quality in industrial areas of Santa Cruz de Tenerife (Spain) by active biomonitoring with *Pseudoscleropodium purum*. – *Ecotoxicology and Environmental Safety* 74: 533–541.
- Armitage H. F., Britton A. J., Woodin S. J. & van der Wal R. (2011): Assessing the recovery potential of alpine moss-sedge heath: Reciprocal transplants along a nitrogen deposition gradient. – *Environmental Pollution* 159: 140–147.
- Asthana A. K. & Sahu V. (2011): Growth responses of a moss *Brachymerium capitulatum* (Mitt.) Par. in different culture media. – *National Academy Science Letters – India* 34: 23–26.
- Bainard J. D. & Newmaster S. G. (2010): Endopolyploidy in bryophytes: widespread in mosses and absent in liverworts. – *Journal of Botany* 2010 (316356):1–7.
- Bainard J. D., Fazekas A. J. & Newmaster S. G. (2010): Methodology significantly affects genome size estimates: quantitative evidence using bryophytes. – *Cytometry, A*: 725–732.
- Balabanova B., Stafilov T., Bačeva K. & Šajn R. (2010): Biomonitoring of atmospheric pollution with heavy metals in the copper mine vicinity located near Radoviš, Republic of Macedonia. – *Journal of Environmental Science and Health, Part a – Toxic/Hazardous Substances and Environmental Engineering* 45: 1504–1518.

- Balan C., Bulai P., Bilba D. & Macoveanu M. (2010): *Sphagnum* moss peat: a green and economical sorbent for removal of heavy metals (Cd and Cr) from wastewaters. – Environmental Engineering and Management Journal 9: 469–477.
- Baldwin L. K. & Bradfield G. E. (2010): Resilience of bryophyte communities in regenerating matrix forests after logging in temperate rainforests of coastal British Columbia. – Botany–Botanique 88: 297–314.
- Baniya C. B. (2010): Vascular and cryptogam richness in the world's highest alpine zone, Tibet. – Mountain Research and Development 30: 275–281.
- Banu-Fattah K. & Sarker S. K. (2010): Bryophyte flora of Greater Mymensingh district of Bangladesh – class: Hepaticopsida and Anthocerotopsida. – Bangladesh Journal of Plant Taxonomy 17: 55–67.
- Bartholomew-Began S. E. & Vitt D. (2011): Barbara J. Crandall-Stotler and Raymond E. Stotler – a tribute. – Bryologist 114: 1–8.
- Basile A., Sorbo S., Conte B., Golia B., Montanari S., Cobianchi R. C. & Esposito S. (2011): Antioxidant activity in extracts from *Leptodictyum riparium* (Bryophyta), stressed by heavy metals, heat shock, and salinity. – Plant Biosystems 145: 77–80.
- Belivermiş M. & Çotuk Y. (2010): Radioactivity measurements in moss (*Hypnum cupressiforme*) and lichen (*Cladonia rangiformis*) samples collected from Marmara region of Turkey. – Journal of Environmental Radioactivity 101: 945–951.
- Bell N. & Hyvönen J. (2010): Molecular evidence for the phylogenetic position of the New Caledonian endemic genus *Parisia*. – Journal of Bryology 32: 303–305.
- Benavides J. C. & Gutierrez C. A. (2011): Effect of deforestation in palm-epiphytic bryophyte communities in a cloud forest in the northern Andes. – Bryologist 114: 155–165.
- Benavides J. C. & Sastre-De Jesus I. (2011): Diversity and rarity of epiphyllous bryophytes in a superhumid tropical lowland forest of Chocó-Colombia. – Cryptogamie Bryologie 21: 119–133.
- Bennie J., Anderson K. & Wetherelt A. (2011): Measuring biodiversity across spatial scales in a raised bog using a novel paired-sample diversity index. – Journal of Ecology 99: 482–490.
- Bisang I., Korpelainen H. & Hedenäs L. (2010): Can the sex-specific molecular marker of *Drepanocladus trifarius* uncover gender in related species? – Journal of Bryology 32: 305–308.
- Blasi C., Marchetti M., Chiavetta U., Aleffi M., Audisio P., Azzella M. M., Brunialti G., Capotorti G., Del Vico E., Lattanzi E., Persiani A. M., Ravera S., Tilia A. & Burrascano S. (2010): Multi-taxon and forest structure sampling for identification of indicators and monitoring of old-growth forest. – Plant Biosystems 144: 160–170.
- Blievernicht A., Irrgang S., Zander M. & Ulrichs C. (2011): Sustainable *Sphagnum* production to replace peat in commercial horticulture. – Gesunde Pflanzen 62: 125–131.
- Blockeel T. L. (2010): The bryophytes of Greece: new records and observations, 2. – Nova Hedwigia Suppl. 138: 129–146.
- Bogdanović M., Sabovljević M., Sabovljević A. & Grubišić G. (2009): The influence of gypsiferous substrata on bryophyte growth: are there obligatory gypsophilous bryophytes? – Botanica Serbica 33: 75–82.
- Boiko M. F. (2009): Mokhopodibni stepovoï zony Ukraïny [Bryobionta of the steppe zone of Ukraine]. – Ailant, Kherson, Ukraine. [264 pp.]
- Bombosch A., Wieneke A., Busch A., Jonas R., Hentschel J., Kreier H.-P., Shaw B., Shaw A. J. & Heinrichs J. (2010): Narrow species concepts in the *Frullania dilatata-appalachiana-eboracensis* complex (Porellales, Jungermanniopsida): evidence from nuclear and chloroplast DNA markers. – Plant Systematics and Evolution 290: 151–158.
- Bond-Lamberty B., Gower S. T., Amiro B. & Ewers B. E. (2011): Measurement and modelling of bryophyte evaporation in a boreal forest chronosequence. – Ecohydrology 4: 26–35.
- Breeuwer A., Heijmans M. M. P. D., Robroek B. J. M. & Berendse F. (2010): Field simulation of global change: transplanting northern bog mesocosms southward. – Ecosystems 13: 712–726.
- Brown R. C., Lemmon B. E. & Shimamura M. (2010): Diversity in meiotic spindle origin and determination of cytokinetic planes in sporogenesis of complex thalloid liverworts (Marchantiopsida). – Journal of Plant Research 123: 589–605.
- Browning B. J., Jordan G. J., Dalton P. J., Grove S. J., Wardlaw T. J. & Turner P. A. M. (2010): Succession of mosses, liverworts and ferns on coarse woody debris, in relation to forest age and log decay in Tasmanian wet eucalypt forest. – Forest Ecology and Management 260: 1896–1905.
- Bruggeman-Nannenga M. A. & Arts T. (2010): a revision of the Fissidentaceae (Musci) of La Réunion (including all species known from Mauritius and Rodriguez). – Journal of Bryology 32: 170–207.
- Brugués M., Ros R. M., Sáez L. & Cano M. J. (2010): *Entosthodon commutatus* Durieu et Mont. (Funariaceae, Bryopsida), new to Europe and Morocco. – Cryptogamie Bryologie 31: 271–276.

- Brugués M., Sérgio C. (2010): On the identity of some *Entosthodon* species endemic to Macaronesia or the North of Africa. – *Cryptogamie Bryologie* 31: 277–280.
- Brunialti G., Frati L., Aleffi M., Marignani M., Rosati L., Burrascano S. & Ravera S. (2010): Lichens and bryophytes as indicators of old-growth features in Mediterranean forests. – *Plant Biosystems* 144: 221–233.
- Buchner O. & Neuner G. (2010): Freezing cytorrhysis and critical temperature thresholds for photosystem II in the peat moss *Sphagnum capillifolium*. – *Protoplasma* 243: 63–71.
- Buck W. R. & Goffinet B. (2010): Preliminary key to the mosses of Isla Navarino, Chile (Prov. Antártica Chilena). – *Nova Hedwigia Suppl.* 138, Suppl. 138: 215–229.
- Buck W. R. & Porto K. C. (2010): *Schwetschkea* (Brachytheciaceae) in Africa, and new to South America. – *Botany–Botanique* 88: 359–365.
- Buczowska K., Sawicki J., Szczecińska M., Klama H., Milewicz M. & Bączkiewicz A. (2010): Genetic variation in the liverwort *Bazzania trilobata* inferred from ISSR markers. – *Journal of Bryology* 32: 265–274.
- Budke J. M. (2010): Examining the gametophytic calyptra and its role in moss sporophyte development using the cord moss (*Funaria hygrometrica*). – *Integrative and Comparative Biology* 50, Suppl. 1: E19–E19.
- Bulgariu L., Bulgariu D. & Macoveanu M. (2010): Kinetics and equilibrium study of nickel(II) removal using peat moss. – *Environmental Engineering and Management Journal* 9: 667–674.
- Burley S. T., Harper K. A. & Lundholm J. T. (2010): Vegetation composition, structure and soil properties across coastal forest-barren ecotones. – *Plant Ecology* 211: 279–296.
- Büttner-Mainik A., Parsons J., Jérôme H., Hartmann A., Lamer S., Schaaf A., Schlosser A., Zipfel P. F., Reski R. & Decker E. L. (2011): Production of biologically active recombinant human factor H in *Physcomitrella*. – *Plant Biotechnology Journal* 9: 373–383.
- Câmara P. E. A. S. (2011): a re-circumscription of the moss genus *Taxithelium* (Pylaisiadelphaceae) with a taxonomic revision of subgenus *Vernieri*. – *Systematic Botany* 36: 7–21.
- Caners R. T., Macdonald S. E., Belland R. J. (2010): Responses of boreal epiphytic bryophytes to different levels of partial canopy harvest. – *Botany–Botanique* 88: 315–328.
- Cano M. J., Jiménez J. A. & Jiménez J. F. (2010): a systematic revision of the genus *Erythrophyllopsis* (Pottiaceae, Bryophyta). – *Systematic Botany* 35: 683–694.
- Cano M. J., Jiménez J. A., Gallego M. T. & Jiménez J. F. (2010): *Guerramontesia microdonta* (Pottiaceae, Bryophyta) a new monotypic genus from South America. – *Systematic Botany* 35: 453–460.
- Carter B. E. (2010): The taxonomic status of the Tasmanian endemic moss, *Scleropodium australe* (Brachytheciaceae). – *Bryologist* 113: 775–780.
- Carvajal B., Aboal J. R., Fernández J. A., Real C. & Carballeira A. (2010): Influence of roads and inhabited areas on metal concentrations in terrestrial mosses. – *Atmospheric Environment* 44: 3432–3441.
- Cezón K., Muñoz J. & Hespanhol H. (2010): The discovery of *Bryum minii* Podp. ex Machado-Guim. in Spain, with new synonyms and correct authorship. – *Bryologist* 113: 371–375.
- Chateigner-Boutin A.-L. & Small I. (2010): Plant RNA editing. – *RNA Biology* 7: 213–219.
- Chen Y. E., Yuan S., Su Y. Q. & Wang L. (2010): Comparison of heavy metal accumulation capacity of some indigenous mosses in Southwest China cities: a case study in Chengdu city. – *Plant Soil and Environment* 56: 60–66.
- Chimner R. A., Lemly J. M. & Cooper D. J. (2010): Mountain fen distribution, types and restoration priorities, San Juan Mountains, Colorado, USA. – *Wetlands* 30: 763–771.
- Chudzińska E. & Odrzykoski I. J. (2010): Fast identification method for an allopolyploid liverwort *Pellia borealis* Lorbeer (Hepaticae, Metzgeriales). – *Cryptogamie Bryologie* 31: 207–215.
- Cogoni A., Brundu G. & Zedda L. (2011): Diversity and ecology of terricolous bryophyte and lichen communities in coastal areas of Sardinia (Italy). – *Nova Hedwigia* 92: 159–175.
- Coleman L. A. & Wilson P. S. (2010): Moss floristics in Sequoia National Park. – *Integrative and Comparative Biology* 50, Suppl. 1: E217.
- Colville L. & Kranner I. (2010): Desiccation tolerant plants as model systems to study redox regulation of protein thiols. – *Plant Growth Regulation* 62: 241–255.
- Cooper E. D., Shaw A. J., Shaw B., Henwood M. J., Heslewood M. M. & Brown E. A. (2011): a multi-locus molecular phylogeny of the Lepidoziaceae: Laying the foundations for a stable classification. – *Molecular Phylogenetics and Evolution* 59: 489–509.
- Coşkun M., Cayir A., Coşkun M. & Kilic O. (2011): Heavy metal deposition in moss samples from East and South Marmara Region, Turkey. – *Environmental Monitoring and Assessment* 174: 219–227.
- Cox C. J., Goffinet B., Wickett N. J., Boles S. B. & Shaw A. J. (2010): Moss diversity: a molecular phylogenetic analysis of genera. – *Phytotaxa* 9: 175–195.

- Crandall-Stotler B., Stotler R. E., Zhang L. & Forrest L. L. (2010): On the morphology, systematics and phylogeny of *Noteroclada* (Noterocladaceae, Marchantiophyta). – *Nova Hedwigia* 91: 421–450.
- Cutler N. (2011): Nutrient limitation during long-term ecosystem development inferred from a mat-forming moss. – *Bryologist* 114: 204–214.
- d'Artenay T. D. & Renzaglia K. S. (2011): Architecture of the mid-stage spermatid of *Aneura pinguis* (Aneuraceae). – *Bryologist* 114: 28–37.
- Daley T. J., Barber K. E., Street-Perrott F. A., Loader N. J., Marshall J. D., Crowley S. F. & Fisher E. H. (2010): Holocene climate variability revealed by oxygen isotope analysis of *Sphagnum* cellulose from Walton Moss, northern England. – *Quaternary Science Reviews* 29: 1590–1601.
- Daniels A. E. D., Raja R. D. A. & Daniel P. (2010): *Indopotia zanderi* (Bryophyta, Pottiaceae) gen. et sp. nov. from the Western Ghats of India. – *Journal of Bryology* 32: 216–219.
- Darell P. & Cronberg N. (2011): Bryophytes in black alder swamps in south Sweden: habitat classification, environmental factors and life-strategies. – *Lindbergia* 34: 9–29.
- Désamoré A., Vanderpoorten A., Laenen B., Gradstein S. R. & Kok P. J. R. (2010): Biogeography of the Lost World (Pantepui region, northeastern South America): Insights from bryophytes. – *Phytotaxa* 9: 254–265.
- Dirkse G. M. & Losada-Lima A. (2011): Additions and amendments to the moss flora of the Canary Islands. – *Cryptogamie Bryologie* 32: 37–41.
- Djupström L. B., Perhans K., Weslien J., Schroeder L. M., Gustafsson L. & Wikberg S. (2010): Co-variation of lichens, bryophytes, saproxylic beetles and dead wood in Swedish boreal forests. – *Systematics and Biodiversity* 8: 247–256.
- Dmuchowski W., Gworek B., Gozdowski D., Baczewska A. & Maciaszek D. (2011): Evaluation of changes in air pollution with lead, zinc, and chromium in the steel mill region in Warsaw in 1993-2008. – *Przemysł Chemiczny* 90: 218–221.
- Dmuchowski W., Gworek B., Gozdowski D., Baczewska A. & Muszyńska A. (2011): Zanieczyszczenie powietrza metalami ciężkimi w rejonie huty cynku i ołowiu w Bukownie koło Olkuszu [Air pollution with heavy metals in the vicinity of zinc and lead smeltery at Bukowno near Olkusz]. – *Przemysł Chemiczny* 90: 223–228.
- dos Santos N. D. & da Costa D. P. (2010): Altitudinal zonation of liverworts in the Atlantic Forest, Southeastern Brazil. – *Bryologist* 113: 631–645.
- Dragović S. & Mandić L. J. (2010): Transfer of radionuclides to ants, mosses and lichens in semi-natural ecosystems. – *Radiation and Environmental Biophysics* 49: 625–634.
- Dragović S., Howard B. J., Caborn J. A., Barnett C. L. & Mihailović N. (2010): Transfer of natural and anthropogenic radionuclides to ants, bryophytes and lichen in a semi-natural ecosystem. – *Environmental Monitoring and Assessment* 166: 677–686.
- Draper I., González-Mancebo J. M., Werner O., Patiño J. & Ros R. M. (2011): Phylogeographic relationships between the mosses *Exsertotheca intermedia* from Macaronesian islands and *Neckera baetica* from southern glacial refugia of the Iberian Peninsula. – *Annales Botanici Fennici* 48: 133–141.
- Drugova T. P. (2010)[2011]: The genus *Aongstroemia* (Dicranaceae, Bryophyta) in Russia. – *Arctoa* 19: 247–252.
- Ederer S. L. & Graham L. K. (2011): Diverse bryophyte-cyanobacterial associations and environmental controls in a Wisconsin boreal forest. – *Journal of Phycology* 47, Sp. Iss. SI Suppl. 1: 6–6.
- Edmondson J. L., Carroll J. A., Price E. A. C. & Caporn S. J. M. (2010): Bio-indicators of nitrogen pollution in heather moorland. – *Science of the Total Environment* 408: 6202–6209.
- Ekberg A., Arneth A. & Holst T. (2011): Isoprene emission from *Sphagnum* species occupying different growth positions above the water table. – *Boreal Environment Research* 16: 47–59.
- Elibol B., Ezer T., Kara R., Çelik G. Y. & Çolak E. (2011): Antifungal and antibacterial effects of some acrocarpic mosses. – *African Journal of Biotechnology* 10: 986–989.
- Elumeeva T. G., Soudzilovskaia N. A., During H. J. & Cornelissen J. H. C. (2011): The importance of colony structure versus shoot morphology for the water balance of 22 subarctic bryophyte species. – *Journal of Vegetation Science* 22: 152–164.
- Engel J. J. (2010): Austral Hepaticae 45. a monograph of the genus *Chiloscyphus* Corda (Lophocoleaceae) for Australasia. – *Fieldiana: Botany, New Series*, 48: i–vi, 1–209.
- Espiñeira J. M., Novo Uzal E., Gómez Ros L. V., Carrión J. S., Merino F., Ros Barceló A. & Pomar F. (2011): Distribution of lignin monomers and the evolution of lignification among lower plants. – *Plant Biology* 13: 59–68.
- Fabian K., Reimann C., McEnroe S. A. & Willemoes-Wissing B. (2011): Magnetic properties of terrestrial moss (*Hylocomium splendens*) along a north-south profile crossing the city of Oslo, Norway. – *Science of the Total Environment* 409: 2252–2260.

- Fabón G., Martínez-Abaigar J., Tomás R. & Núñez-Olivera E. (2010): Effects of enhanced UV-B radiation on hydroxycinnamic acid derivatives extracted from different cell compartments in the aquatic liverwort *Jungermannia exsertifolia* subsp. *cordifolia*. – *Physiologia Plantarum* 140: 269–279.
- Fabure J., Meyer C., Denayer F., Gaudry A., Gilbert D. & Bernard N. (2010): Accumulation capacities of particulate matter in an acrocarpous and a pleurocarpous moss exposed at three differently polluted sites (industrial, urban and rural). – *Water, Air & Soil Pollution* 212: 205–217.
- Favreau M. & Brassard G. R. (2011): An analysis of *Meesia* (Meesiaceae, Musci) in arctic North America and Greenland. – *Carnets de Bryologie* 1: 2–9.
- Fedosov V. E., Ignatova E. A., Ignatov M. S. & Doroshina G. Ya. (2010)[2011]: On the genus *Entosthodon* (Funariaceae, Musci) in the Caucasus. – *Arctoa* 19: 75–86.
- Fenton N. J., Bergeron Y. & Paré D. (2010): Decomposition rates of bryophytes in managed boreal forests: influence of bryophyte species and forest harvesting. – *Plant and Soil* 336: 499–508.
- Fife A. J., Glenny D., Beever J. E., Braggins J. E., Brownsey P. J., Renner M. A. M. & Hitchmough R. (2010): New Zealand's threatened bryophytes: Conservation efforts. – *Phytotaxa* 9: 275–278.
- Finet C., Timme R. E., Delwiche C. F. & Marleta F. (2010): Multigene phylogeny of the green lineage reveals the origin and diversification of land plants. – *Current Biology* 20: 2217–2222.
- Foan L., Sablayrolles C., Elustondo D., Lasheras E., González L., Ederra A., Simon V. & Santamaría J. M. (2010): Reconstructing historical trends of polycyclic aromatic hydrocarbon deposition in a remote area of Spain using herbarium moss material. – *Atmospheric Environment* 44: 3207–3214.
- Fontinha S., Sim-Sim M. & Sérgio C. (2010): The genus *Porella* (Porellaceae, Marchantiophyta) in Madeira Archipelago ecological and morphological approaches. – *Nova Hedwigia Suppl.* 138, Suppl. 138: 297–310.
- Forrest L. L., Allen N. S., Gudiño J. A., Korpelainen H. & Long D. G. (2011): Molecular and morphological evidence for distinct species in *Dumortiera* (Dumortieraceae). – *Bryologist* 114: 102–115.
- Forrest L. L., Wickett N. J., Cox C. J. & Goffinet B. (2011): Deep sequencing of *Ptilidium* (Ptilidiaceae) suggests evolutionary stasis in liverwort plastid genome structure. – *Plant Ecology and Evolution* 144: 29–43.
- Frahm J.-P. (2010): Mosses and Liverworts of the Mascarenes and the Seychelles. – Books on Demand GmbH, Norderstedt, Germany. [144 pp.]
- Frahm J.-P. (2010): Mosses and Liverworts of the Mediterranean: An Illustrated Field Guide. – Books on Demand GmbH, Norderstedt, Germany. [138 pp.]
- Frahm J.-P. (2010): *Tortula aestiva* (Hedw.) P. Beauv., eine gute Art. – *Archive for Bryology* 66: 1–4.
- Frahm J.-P. (2011): Kommentierte Liste der Laubmoose Deutschlands. – *Archive for Bryology* 79: 1–51.
- Frahm J.-P. (2011): *Leucobryum juniperoideum* (Brid.) C.Müll. in North America. – *Archive for Bryology* 84: 1–6.
- Frahm J.-P. (2011): The taxonomic position of *Orthotrichum fastigiatum*. – *Archive for Bryology* 81: 1–7.
- Frey W. & Kürschner H. (2011): Asexual reproduction, habitat colonization and habitat maintenance in bryophytes. – *Flora* 206: 173–184.
- Frey W., Pfeiffer T. & Stech M. (2010): Geomolecular divergence patterns of Gondwanan and Palaeoaustral bryophytes – An overview Studies in austral temperate rain forest bryophytes 34. – *Nova Hedwigia* 91: 317–348.
- Fritz S., Lieske K. & Frey W. (2010): Vegetative (asexual) reproduction by propagules in six dioicous pleurocarpous mosses (*Abietinella abietina*, *Homalothecium lutescens*, *H. sericeum*, *Pleurozium schreberi*, *Pseudoscleropodium purum*, *Rhytidiadelphus squarrosus*). a morphological study. – *Nova Hedwigia Suppl.* 138, Suppl. 138: 325–333.
- Fuell C., Elliott K. A., Hanfrey C. C., Franceschetti M. & Michael A. J. (2010): Polyamine biosynthetic diversity in plants and algae. – *Plant Physiology and Biochemistry* 48: 513–520.
- Fuselier L. C., Shaw B., Engel J. J., von Konrat M., Costa D. P., Devos N. & Shaw A. J. (2011): The status and phylogeography of the liverwort genus *Apometzgeria* Kuwah. (Metzgeriaceae). – *Bryologist* 114: 92–101.
- Fuselier L., Bougary A. & Malott M. (2011): From trace evidence to bioinformatics: Putting bryophytes into molecular biology education. – *Biochemistry and Molecular Biology Education* 39: 38–46.
- Garibotti I. A., Pissolito C. I. & Villalba R. (2011): Vegetation Development on deglaciated rock outcrops from Glaciar Frías, Argentina. – *Arctic Antarctic and Alpine Research* 43: 35–45.
- Gauthier R. (2011): Le genre *Tetrodontium* (Tetraphidaceae, Musci) au Québec. – *Carnets de Bryologie* 1 : 10–14.

- Gerrienne P., Gonez P. (2011): Early evolution of life cycles in embryophytes: a focus on the fossil evidence of gametophyte/sporophyte size and morphological complexity. – *Journal of Systematics and Evolution* 49: 1–16.
- Gilabert M., Ramos A. N., Schiavone M. M., Arena M. E. & Bardon A. (2011): Bioactive sesqui- and diterpenoids from the Argentine liverwort *Porella chilensis*. – *Journal of Natural Products* 74: 574–579.
- González-Mancebo J. M., Patiño J., Rodríguez-Romero A., Werner O. & Ros R. M. (2010): *Gymnostomiella* (Pottiaceae, Bryophyta) revisited new insights revealed by a morphometric analysis. – *Nova Hedwigia Suppl.* 138, Suppl. 138: 69–83.
- González-Miqueo L., Elustondo D., Lasheras E., Bermejo R. & Santamaría J. M. (2010): Heavy metal and nitrogen monitoring using moss and topsoil samples in a Pyrenean forest catchment. – *Water, Air & Soil Pollution* 210: 335–346.
- Gradstein S. R. & Sporn S. G. (2010): Land-use change and epiphytic bryophyte diversity in the Tropics. – *Nova Hedwigia Suppl.* 138: 311–323.
- Gradstein S. R., Ilkiu-Borges A. L., & Vanderpoorten A. (2011): Habitat specialization triggers the evolution of unusual morphologies – the case of *Cololejeunea stotleriana* sp. nov. from Ecuador. – *Bryologist* 114: 9–22
- Graf U., Wildi O., Feldmeyer-Christe E. & Kuchler M. (2010): a phytosociological classification of Swiss mire vegetation. – *Botanica Helvetica* 120: 1–13.
- Graham L. E., Cook M. E., Hanson D. T., Pigg K. B. & Graham J. M. (2010): Rolled liverwort mats explain major prototaxites features: response to commentaries. – *American Journal of Botany* 97: 1079–1086.
- Graham L. E., Kim E., Arancibia-Avila P., Graham J. M. & Wilcox L. W. (2010): Evolutionary and ecophysiological significance of sugar utilization by the peat moss *Sphagnum compactum* (Sphagnaceae) and the common charophycean associates *Cylindrocystis brebissonii* and *Mougeotia* sp. (Zygnemataceae). – *American Journal of Botany* 97: 1485–1491.
- Granath G., Strengbom J. & Rydin H. (2010): Rapid ecosystem shifts in peatlands: linking plant physiology and succession. – *Ecology* 91: 3047–3056.
- Grdović S., Vitorović G., Mitrović B., Andrić V., Petrujkić B. & Obradović M. (2010): Natural and anthropogenic radioactivity of feedstuffs, mosses and soil in the Belgrade environment, Serbia. – *Archives of Biological Sciences* 62: 301–307.
- Gudelis A., Druteikienė R., Lukšienė B., Gvozdaitė R., Nielsen S. P., Hou X., Mažeika J. & Petrošius R. (2010): Assessing deposition levels of <sup>55</sup>Fe, <sup>60</sup>Co and <sup>63</sup>Ni in the Ignalina NPP environment. – *Journal of Environmental Radioactivity* 101: 464–467.
- Guerra J., Brugués M., Cano M. J. & Cros R. M. (2010): Funariales: Ephemeraceae, Gigaspermaceae, Funariaceae; Splachnales: Splachnaceae, Meesiaceae; Schistostegales: Schistostegaceae; Bryales: Bryaceae, Mielichhoferiaceae, Mniaceae, Aulacomniaceae, Catosciaceae, Bartramiaceae, Orthodontiaceae; Timmiales: Timmiaceae. – In: Guerra J. & Cros R. M., eds, *Flora Briofítica Ibérica*, Vol. IV, p. 1–317, Sociedad Española de Briología, Murcia, Spain.
- Guerra J., Jiménez-Martínez J. F. & Jiménez J. A. (2010): *Neckera baetica* sp. nov. (Neckeraceae, Bryophyta) from southern Spain, based on morphological and molecular data. – *Nova Hedwigia* 91: 255–263.
- Gupta A. (2010): European *Cryptothallus mirabilis* v. Malmb. matured in India – its morphological and LM & SEM palynological studies, with comments on utility for modern and fossil studies. – *Current Science* 99: 1116–1120.
- Haapala J. K., Mörsky S. K., Saarnio S., Suokanerva H., Kyrö E., Silvola J. & Holopainen T. (2010): Long-term effects of elevated UV-B radiation on photosynthesis and ultrastructure of *Eriophorum russeolum* and *Warnstorfia exannulata*. – *Science of the Total Environment* 409: 370–377.
- Hagemann U., Moroni M. T., Gleissner J. & Makeschin F. (2010): Accumulation and preservation of dead wood upon burial by bryophytes. – *Ecosystems* 13: 600–611.
- Hájek M., Horsák M., Tichý L., Hájková P., Dítě D. & Jamrichová E. (2011): Testing a relict distributional pattern of fen plant and terrestrial snail species at the Holocene scale: a null model approach. – *Journal of Biogeography* 38: 742–755.
- Hájek T., Ballance S., Limpens J., Zijlstra M. & Verhoeven J. T. A. (2011): Cell-wall polysaccharides play an important role in decay resistance of *Sphagnum* and actively depressed decomposition *in vitro*. – *Biogeochemistry* 103: 45–57.
- Hallingbäck T. & Tan B. C. (2010): Past and present activities and future strategy of bryophyte conservation. – *Phytotaxa* 9: 266–274.
- Hardman A. & McCune B. (2010): Bryoid layer response to soil disturbance by fuel reduction treatments in a dry conifer forest. – *Bryologist* 113: 235–245.

- Harmens H., Norris D. A., Steinnes E., Kubin E., Piispanen J., Alber R., Aleksiyenak Y., Blum O., Coşkun M., Dam M., De Temmerman L., Fernández J. A., Frolova M., Frontasyeva M., González-Miqueo L., Grodzińska K., Jeran Z., Korzekwa S., Krmar M., Kvietskus K., Leblond S., Liiv S., Magnússon S. H., Maňková B., Pesch R., Rühling Å., Santamaría J. M., Schröder W., Spiric Z., Suchara I., Thöni L., Urumov V., Yurukova L. & Zechmeister H. G. (2010): Mosses as biomonitors of atmospheric heavy metal deposition: Spatial patterns and temporal trends in Europe. – *Environmental Pollution* 158: 3144–3156.
- Harpel J. A. (2010): a history of bryology in the San Juan Islands, Washington State, USA. – *Botany–Botanique* 88: 329–335.
- Harrison C. J., Alvey E. & Henderson I. R. (2010): Meiosis in flowering plants and other green organisms. – *Journal of Experimental Botany* 61: 2863–2875.
- Hartung W. (2010): The evolution of abscisic acid (ABA) and ABA function in lower plants, fungi and lichen. – *Functional Plant Biology* 37: 806–812.
- Hautala H., Laaka-Lindberg S. & Vanha-Majamaa I. (2011): Effects of retention felling on epixylic species in boreal spruce forests in Southern Finland. – *Restoration Ecology* 19: 418–429.
- He X. & Glenn D. (2010): *Perssoniella* and the genera of Schistochilaceae: a new classification based on molecular phylogenies. – *Australian Systematic Botany* 23: 229–238.
- Hedenäs L. (2010): ITS distinguishes *Palustriella falcata* from *P. commutata* and *P. decipiens*, but not from *P. pluristratosa* (Amblystegiaceae, Bryophyta). – *Nova Hedwigia Suppl.* 138: 51–60.
- Hedenäs L. (2010): Phylogeography and origin of European *Sanionia uncinata* (Amblystegiaceae, Bryophyta). – *Systematics and Biodiversity* 8: 177–191.
- Hedenäs L. & Zare H. (2010): The Euxinian-Hyrcanian endemic species *Forsstroemia remotifolia* (Lindb. ex Broth.) Hedenäs & Zare, comb. nov. (Neckeraceae, Bryophyta). – *Nova Hedwigia Suppl.* 138: 61–67.
- Hedwall P.-O., Nordin A., Brunet J. & Bergh J. (2010): Compositional changes of forest-floor vegetation in young stands of Norway spruce as an effect of repeated fertilisation. – *Forest Ecology and Management* 259: 2418–2425.
- Heinrichs J., Hentschel J., Bombosch A., Fiebig A., Reise J., Edelmann M., Kreier H.-P., Schäfer-Verwimp A., Caspari S., Schmidt A. R., Zhu R.-L., von Konrat M., Shaw B. & Shaw A. J. (2010): One species or at least eight? Delimitation and distribution of *Frullania tamarisci* (L.) Dumort. s. l. (Jungermanniopsida, Porellales) inferred from nuclear and chloroplast DNA markers. – *Molecular Phylogenetics and Evolution* 56: 1105–1114.
- Hejzman M., Száková J., Schellberg J., Šrek P., Tlustoš P. & Balík J. (2010): The Rengen Grassland experiment: bryophytes biomass and element concentrations after 65 years of fertilizer application. – *Environmental Monitoring and Assessment* 166: 653–662.
- Hermesen C., Koprivova A., Matthewman C., Wesenberg D., Krauss G.-J. & Kopriva S. (2010): Regulation of sulfate assimilation in *Physcomitrella patens*: mosses are different! – *Planta* 232: 461–470.
- Hills S. F. K., Stevens M. I. & Gemmill C. E. C. (2010): Molecular support for Pleistocene persistence of the continental Antarctic moss *Bryum argenteum*. – *Antarctic Science* 22: 721–726.
- Hodgetts N. G. (2011): a revised Red List of bryophytes in Britain. – *Field Bryology* 103: 40–49.
- Holm K., Källman T., Gyllenstrand N., Hedman H. & Lagercrantz U. (2010): Does the core circadian clock in the moss *Physcomitrella patens* (Bryophyta) comprise a single loop? – *BMC Plant Biology* 10: –109.
- Holy M., Pesch R., Schröder W., Harmens H., Ilyin I., Alber R., Aleksiyenak Y., Blum O., Coşkun M., Dam M., De Temmerman L., Fedorets N., Figueira R., Frolova M., Frontasyeva M., Goltsova N., Gonzalez M. L., Grodzińska K., Jeran Z., Korzekwa S., Krmar M., Kubin E., Kvietskus K., Larsen M., Leblond S., Liiv S., Magnússon S., Maňková B., Mocanu R., Piispanen J., Rühling Å., Santamaría J., Steinnes E., Suchara I., Thöni L., Turcsányi G., Urumov V., Wolterbeek B., Yurukova L. & Zechmeister H. G. (2009): First thorough identification of factors associated with Cd, Hg and Pb concentrations in mosses sampled in the European Surveys 1990, 1995, 2000 and 2005. – *Journal of Atmospheric Chemistry* 63: 109–124.
- Holyoak D. T. & Köckinger H. (2010): a taxonomic revision of some European and Asian bulbiferous species of *Anomobryum* (Bryophyta: Bryaceae). – *Journal of Bryology* 32: 153–169.
- Hooper E. J., Duckett J. G., Cumming A. C., Kunin W. E. & Pressel S. (2010): *Ephemerella readeri* Mull. Hal. (*Physcomitrella readeri* (Müll. Hal.) I.G. Stone & G.A.M. Scott, Funariidae, Bryophyta): a genus and species new to Europe. – *Journal of Bryology* 32: 256–264.
- Horsley K., Stark L. R. & McLetchie D. N. (2011): Does the silver moss *Bryum argenteum* exhibit sex-specific patterns in vegetative growth rate, asexual fitness or prezygotic reproductive investment? – *Annals of Botany* 107: 897–907.
- Hrivnák R., O'áhel'ová H., Valachovič M., Pa'ove-Balang P. & Kubinská A. (2010): Effect of environmental variables on the aquatic macrophyte composition pattern in streams: a case study from Slovakia. – *Fundamental and Applied Limnology* 177: 115–124.

- Hugonnot V. (2010): Mousses et hépatiques de Païolive (Ardèche et Gard, France). – Bulletin de la Société Botanique de Centre-Ouest, nouvelle série, Numéro Spécial 34: 1–293.
- Hugonnot V. (2010): Towards an improved understanding of the taxonomy of *Riccia ciliata* Hoffm. (Marchantiopsida: Ricciaceae). – Journal of Bryology 32: 300–303.
- Hugonnot V. (2011): Bryophytes of the peatlands of Aubrac (Massif Central – France). – Cryptogamie Bryologie 32: 43–63.
- Humphreys C. P., Franks P. J., Rees M., Bidartondo M. I., Leake J. R. & Beerling D. J. (2010): Mutualistic mycorrhiza-like symbiosis in the most ancient group of land plants. – Nature Communications 1: 103.
- Huttunen S. & Ignatov M. S. (2010): Evolution and taxonomy of aquatic species in the genus *Rhynchostegium* (Brachytheciaceae, Bryophyta). – Taxon 59: 791–808.
- Hylander K. & Johnson S. (2010): *In situ* survival of forest bryophytes in small-scale refugia after an intense forest fire. – Journal of Vegetation Science 21: 1099–1109.
- Ignatov M. S. & Milyutina I. A. (2010)[2011]: The genus *Brachythecium* (Brachytheciaceae, Musci) in Russia: comments on species and key for identification. – Arctoa 19: 1–30.
- Ignatov M. S., Fedosov V. E., Ignatova E. A., Doroshina G. Ya. & Zolotov V. I. (2010)[2011]: Moss flora of Gunib area in Dagestan, the Eastern Caucasus. – Arctoa 19: 87–96.
- Ignatova E. A., Blom H. H., Goryunov D. V. & Milyutina I. A. (2010)[2011]: On the genus *Schistidium* (Grimmiaceae, Musci) in Russia. . – Arctoa 19: 195–233.
- Ignatova E. A., Ignatov M. S. & Milyutina I. A. (2010)[2011]: a revision of the genus *Lindbergia* (Leskeaceae, Bryophyta) in Russia. – Arctoa 19: 97–116.
- Ilomets M., Truus L., Pajula R., & Sepp K. (2010): Species composition and structure of vascular plants and bryophytes on the water level gradient within a calcareous fen in North Estonia. – Estonian Journal of Ecology 59:19–38.
- Infante M. (2010): Notes on the genus *Dendroceros* in West Africa and South Atlantic Islands. – Journal of Bryology 32: 283–287.
- Iwatsuki Z. (2010): Type specimens of mosses located in the herbarium of the Hattori Botanical Laboratory (NICH): Part 1, species from Japan. – Hattoria 1: 31–70.
- Jackson B. G., Martin P., Nilsson M.-C. & Wardle D. A. (2011): Response of feather moss associated N<sub>2</sub> fixation and litter decomposition to variations in simulated rainfall intensity and frequency. – Oikos 120: 570–581.
- Janauer G. A., Schmidt-Mumm U. & Schmidt B. (2010): Aquatic macrophytes and water current velocity in the Danube River. – Ecological Engineering 36: 1138–1145.
- Jassey V. E. J., Gilbert D., Binet P., Toussaint M.-L. & Chiapusio G. (2011): Effect of a temperature gradient on *Sphagnum fallax* and its associated living microbial communities: a study under controlled conditions. – Canadian Journal of Microbiology 57: 226–235.
- Jensen S., Omarsdottir S., Ogmundsdottir H. & Olafsdottir E. (2010): Bioprospecting Icelandic liverworts for anticancer activity. – Planta Medica 76: 1208.
- Jesson L. K., Cavanagh A. P. & Perley D. S. (2011): Polyploidy influences sexual system and mating patterns in the moss *Atrichum undulatum* sensu lato. – Annals of Botany 107: 135–143.
- Jiroušek M., Hájek M. & Bragazza L. (2011): Nutrient stoichiometry in *Sphagnum* along a nitrogen deposition gradient in highly polluted region of Central-East Europe. – Environmental Pollution 159: 585–590.
- Juutinen R. & Kotiaho J. S. (2011): Finnish Forest Act as a conservation tool in protecting boreal springs and associated bryophyte flora. – Boreal Environment Research 16: 136–148.
- Juutinen R. (2011): The decrease of rich fen bryophytes in springs as a consequence of large-scale environmental loss. a 50-year re-sampling study. – Lindbergia 34: 2–8.
- Kalachova G. S., Gladyshev M. I., Sushchik N. N. & Makhutova O. N. (2011): Water moss as a food item of the zoobenthos in the Yenisei River. – Central European Journal of Biology 6: 236–245.
- Karlin E. F., Andrus R. E., Boles S. B. & Shaw A. J. (2011): One haploid parent contributes 100% of the gene pool for a widespread species in northwest North America. – Molecular Ecology 20: 753–767.
- Karlin E. F., Boles S. B., Seppelt R. D., Terracciano S. & Shaw A. J. (2011): The peat moss *Sphagnum cuspidatum* in Australia: microsatellites provide a global perspective. – Systematic Botany 36: 22–32.
- Kempter H., Krachler M. & Shotyk W. (2010): Atmospheric Pb and Ti accumulation rates from *Sphagnum* moss: dependence upon plant productivity. – Environmental Science & Technology 44: 5509–5515.
- Keuper F., Dorrepaal E., Van Bodegom P. M., Aerts R., Van Logtestijn R. S. P., Callaghan T. V. & Cornelissen J. H. C. (2011): a race for space? How *Sphagnum fuscum* stabilizes vegetation composition during long-term climate manipulations. – Global Change Biology 17: 2162–2171.



- Kicsi A., Bilba D. & Macoveanu M. (2010): Equilibrium and kinetic modeling of Zn (II) sorption from aqueous solutions by *Sphagnum* moss peat. – Environmental Engineering and Management Journal 9: 341–349.
- Kicsi A., Cojocaru C., Macoveanu M. & Bilba D. (2010): Response surface methodology applied for zinc removal from aqueous solutions using *Sphagnum* peat moss as sorbent. – Journal of Environmental Protection and Ecology 11: 614–622.
- Király I. & Ódor P. (2010): The effect of stand structure and tree species composition on epiphytic bryophytes in mixed deciduous-coniferous forests of Western Hungary. – Biological Conservation 143: 2063–2069.
- Köckinger H., Werner O. & Ros R. M. (2010): a new taxonomic approach to the genus *Oxystegus* (Pottiaceae, Bryophyta) in Europe based on molecular data. – Nova Hedwigia Suppl. 138: 31–49.
- Konno H., Nakashima S. & Katoh K. (2010): Metal-tolerant moss *Scopelophila cataractae* accumulates copper in the cell wall pectin of the protonema. – Journal of Plant Physiology 167: 358–364.
- Konstantinova N. A. & Mamontov Yu. S. (2010)[2011]: a revision of the genus *Calycularia* Mitt. (Calyculariaceae, Marchantiophyta). – Arctoa 19: 117–130.
- Körner C. (2011): Coldest places on earth with angiosperm plant life. – Alpine Botany 121: 11–22.
- Korpelainen H., von Craeutlein M., Laaka-Lindberg S. & Huttunen S. (2011): Fine-scale spatial genetic structure of a liverwort (*Barbilophozia attenuata*) within a network of ant trails. – Evolutionary Ecology 25: 45–57.
- Kosior G., Samecka-Cymerman A., Kolon K. & Kempers A. J. (2010): Bioindication capacity of metal pollution of native and transplanted *Pleurozium schreberi* under various levels of pollution. – Chemosphere 81: 321–326.
- Košnar J. & Kučera J. (2010): The taxonomic identity and typification of *Barbula montenegrina* Breidl. & Szyszyl. (Bryopsida, Pottiaceae). – Journal of Bryology 32: 275–278.
- Koster K. L., Balsamo R. A., Espinoza C. & Oliver M. J. (2010): Desiccation sensitivity and tolerance in the moss *Physcomitrella patens*: assessing limits and damage. – Plant Growth Regulation 62: 293–302.
- Kreier H.-P., Feldberg K., Mahr F., Bombosch A., Schmidt A. R., Zhu R.-L., von Konrat M., Shaw B., Shaw A. J. & Heinrichs J. (2010): Phylogeny of the leafy liverwort *Ptilidium*: Cryptic speciation and shared haplotypes between the Northern and Southern Hemispheres. – Molecular Phylogenetics and Evolution 57: 1260–1267.
- Kruijer J. D., Raes N. & Stech M. (2010): Modelling the distribution of the moss species *Hypopterygium tamarisci* (Hypopterygiaceae, Bryophyta) in Central and South America. – Nova Hedwigia 91: 399–420.
- Kučera J. (2010): 5. *Grimmia dissimulata* (Czech Republic). – In: Ellis L.T. et al., New national and regional bryophyte records, 24, Journal of Bryology 32: 232–237, p. 233.
- Kürschner H. & Frey W. (2011): Liverworts, mosses and hornworts of Southwest Asia (Marchantiophyta, Bryophyta, Anthocerotophyta). – Nova Hedwigia Suppl. 139. – J. Cramer, Stuttgart. [240 pp.]
- Kürschner H. (2010): Amendments to the bryophyte flora of Lebanon, based on collections of C. B. Arzeni 1962–1964. – Bryologist 113: 717–720.
- Laenen B., Desamore A., Devos N., Shaw A. J., González-Mancebo M. J., Carine M. A. & Vanderpoorten A. (2011): Macaronesia: a source of hidden genetic diversity for post-glacial recolonization of western Europe in the leafy liverwort *Radula lindenbergiana*. – Journal of Biogeography 38: 631–639.
- Lafleur B., Paré D., Fenton N. J. & Bergeron Y. (2010): Do harvest methods and soil type impact the regeneration and growth of black spruce stands in northwestern Quebec? – Canadian Journal of Forest Research 40: 1843–1851.
- Lang A. & Naciri Y. (2010): New chloroplast primers for intraspecific variation in *Dicranum scoparium* Hedw. (Dicranaceae) and amplification success in other bryophyte species. – Molecular Ecology Resources 10: 735–737.
- Lara F., Mazimpaka V., Medina R., Caparros R. & Garilleti R. (2010): Northeastern Turkey, an unnoticed but very important area for the Orthotrichaceae (Bryophyta). – Nova Hedwigia Suppl. 138: 165–180.
- Larmola T., Tuittila E.-S., Tiirola M., Nykänen H., Martikainen P. J., Yrjälä K., Tuomivirta T. & Fritze H. (2010): The role of *Sphagnum* mosses in the methane cycling of a boreal mire. – Ecology 91: 2356–2365.
- Leal I. R., Bieber A. G. D., Tabarelli M. & Andersen A. N. (2010): Biodiversity surrogacy: indicator taxa as predictors of total species richness in Brazilian Atlantic forest and Caatinga. – Biodiversity and Conservation 19: 3347–3360.
- Lee L., Rosenstiel T. N. & Eppley S. M. (2010): Variation of photoperiod response in moss gametangial formation. – Bryologist 113: 673–678.
- Lehtonen S., Myllys L. & Huttunen S. (2009): Phylogenetic analysis of non-coding plastid DNA in the presence of short inversions. – Phytotaxa 1: 3–20.

- Lenne T., Bryant G., Hocart C. H., Huang C. X. & Ball M. C. (2010): Freeze avoidance: a dehydrating moss gathers no ice. – *Plant Cell and Environment* 33: 1731–1741.
- Li Y., Wang Z., Xu T., Tu W., Liu C., Zhang Y. & Yang C. (2010): Reorganization of photosystem II is involved in the rapid photosynthetic recovery of desert moss *Syntrichia caninervis* upon rehydration. – *Journal of Plant Physiology* 167: 1390–1397.
- Liang S.-F., Sun Y. & Zhu R.-L. (2010): *In vitro* micropropagation of *Bryum argenteum* Hedw. – *Cryptogamie Bryologie* 31: 233–239.
- Lindo Z. & Gonzalez A. (2010): The bryosphere: An integral and influential component of the Earth's biosphere. – *Ecosystems* 13: 612–627.
- Lindo Z. & Whiteley J. A. (2011): Old trees contribute bio-available nitrogen through canopy bryophytes. – *Plant and Soil* 342: 141–148.
- Linis V. C. (2010): The mossflora of Camiguin Island, Philippines and their floristic relations to some adjacent islands in the archipelago. – *Telopea* 12: 525–542.
- Liu X.-Y., Xiao H.-Y., Liu C.-Q., Li Y.-Y., Xiao H.-W. & Wang Y.-L. (2010): Response of stable carbon isotope in epilithic mosses to atmospheric nitrogen deposition. – *Environmental Pollution* 158: 2273–2281.
- Liu Y., Cao T. & Ge X.-J. (2011): a case study of DNA barcoding in Chinese Grimmiaceae and a moss recorded in China for the first time. – *Taxon* 60: 185–193.
- Liu Y., Ge X.-J., Sun Q.-B. & Cao T. (2010): Development of microsatellite markers for the moss *Ptychomitrium gardneri* (Ptychomitriaceae). – *American Journal of Botany* 97: E14–E16.
- Lobe C., Sim-Sim M., Luís L. & Stech M. (2011): Chorological data and threat status evaluation of *Fissidens* species (Fissidentaceae, Bryophyta) on Madeira Archipelago. – *Nova Hedwigia* 92: 121–131.
- Löbel S., Rydin H. (2010): Trade-offs and habitat constraints in the establishment of epiphytic bryophytes. – *Functional Ecology* 24: 887–897.
- Locky D. A. & Bayley S. E. (2010): Plant diversity in wooded moderate-rich fens across boreal western Canada: an ecoregional perspective. – *Biodiversity and Conservation* 19: 3525–3543.
- Lucaciu A., Moțoc C., Jelea M. & Jelea S. G. (2010): Survey of heavy metal deposition in Romania: Transylvanian Plateau and Western Carpathians Mountains. – *University Politehnica of Bucharest Scientific Bulletin, Series A*, 72: 171–178.
- Lücking R., Mata-Lorenzen J. & Dauphin L. G. (2010): Epizoic liverworts, lichens and fungi growing on Costa Rican Shield Mantis (Mantodea: Choeradodis). – *Studies on Neotropical Fauna and Environment* 45: 175–186.
- Ludwiczuk A., Gradstein S. R., Nagashima F. & Asakawa Y. (2011): Chemosystematics of *Porella* (Marchantiophyta, Porellaceae). – *Natural Product Communications* 6: 315–321.
- Ludwiczuk A., Raharivelomanana P., Komala I., Pham A., Bianchini J. & Asakawa Y. (2010): Chemodiversity of Polynesian liverworts. – *Planta Medica* 76: 1198.
- Ludwig-Müller J. (2011): Auxin conjugates: their role for plant development and in the evolution of land plants. – *Journal of Experimental Botany* 62: 1757–1773.
- Lüth M. (2010): Bildatlas der Moose Deutschlands, Vol. 7: Brachytheciaceae – Sematophyllaceae. – Bryologische Arbeitsgemeinschaft Deutschlands, Bonn. [121 color plates]
- Lüth M. (2011): Bildatlas der Moose Deutschlands, Vol. 1b: Sphagnaceae. – Bryologische Arbeitsgemeinschaft Deutschlands, Bonn. [36 color plates]
- Maciel da Silva A. S., Pôrto K. C. & Simabukuro E. A. (2010): Effects of light and nutrients on different germination phases of the cosmopolitan moss *Bryum argenteum* Hedw. (Bryaceae). – *Brazilian Archives of Biology and Technology* 53: 763–769.
- Magill R. E. (2010): Moss diversity: New look at old numbers. – *Phytotaxa* 9: 167–174.
- Mälson K., Sundberg S. & Rydin H. (2010): Peat disturbance, mowing, and ditch blocking as tools in rich fen restoration. – *Restoration Ecology* 18: 469–478.
- Mandl N., Lehnert M., Kessler M. & Gradstein S. R. (2010): a comparison of alpha and beta diversity patterns of ferns, bryophytes and macrolichens in tropical montane forests of southern Ecuador. – *Biodiversity and Conservation* 19: 2359–2369.
- Manolaki P., Tsakiri E. & Papastergiadou E. (2011): Inventory of aquatic and riparian flora of Acheron and Louros Rivers, and Zirou Lake in Western Greece. – *Fresenius Environmental Bulletin* 20: 861–874.
- Marešová J., Pipiška M., Rozložník M., Horník M., Remenárová L. & Augustín J. (2011): Cobalt and strontium sorption by moss biosorbent: Modeling of single and binary metal systems. – *Desalination* 266: 134–141.
- Mariet C., Gaudry A., Ayrault S., Moskura M., Denayer F. & Bernard N. (2011): Heavy metal bioaccumulation by the bryophyte *Scleropodium purum* at three French sites under various influences: rural conditions, traffic, and industry. – *Environmental Monitoring and Assessment* 174: 107–118.

- Marinova S., Frontasyeva M., Yurukova L., Strelkova L., Steinnes E. & Marinov A. (2009): The moss techniques for air pollution study in Bulgaria. – 7th International Conference of the Balkan Physical Union Vols 1 and 2 1203: 777–782.
- Marinova S., Yurukova L., Frontasyeva M. V., Steinnes E., Strelkova L. P., Marinov A. & Karadzhinova A. G. (2010): Air pollution studies in Bulgaria using the moss biomonitoring technique. – *Ecological Chemistry and Engineering s* 17: 37–52.
- Marka J. & Sabovljević M. (2011): New bryophyte records from Albania. – *Journal of Bryology* 33: 74–76.
- Marstaller R. (2010): The bryophyte communities of the order *Hylocomietalia splendentis* Gillet ex Vadam 1990. 141<sup>th</sup> contribution to the bryophyte vegetation of Thuringia. – *Herzogia* 23: 279–311.
- Martin J. L., Cardoso P., Arechavaleta M., Borges P. A. V., Faria B. F., Abreu C., Aguiar A. F., Carvalho J. A., Costa A. C., Cunha R. T., Fernandes F. M., Gabriel R., Jardim R., Lobo C., Martins A. M. F., Oliveira P., Rodrigues P., Silva L., Teixeira D., Amorim I. R., Homem N., Martins B., Martins M. & Mendonça E. (2010): Using taxonomically unbiased criteria to prioritize resource allocation for oceanic island species conservation. – *Biodiversity and Conservation* 19: 1659–1682.
- Martins R. J. E. & Boaventura R. A. R. (2011): Modelling of lead removal by an aquatic moss. – *Water Science and Technology* 63: 136–142.
- Martins R. J. E., Vilar V. J. P. & Boaventura R. A. R. (2010): Removal of Pb(II) from wastewaters by *Fontinalis antipyretica* biomass: Experimental study and modelling. – *Desalination and Water Treatment* 20: 179–188.
- Mazimpaka V., Medina N. G., Lo Giudice R., Garilleti R. & Lara F. (2010): Tree age-dependent changes among epiphytic bryophyte communities in Mediterranean environments. a case study from Sicily (Italy). – *Plant Biosystems* 144: 241–249.
- McMullan-Fisher S. J. M., Kirkpatrick J. B., May T. W. & Pharo E. J. (2010): Surrogates for macrofungi and mosses in reservation planning. – *Conservation Biology* 24: 730–736.
- Meyer C., Bernard N., Moskura M., Toussaint M. L., Denayer F. & Gilbert D. (2010): Effects of urban particulate deposition on microbial communities living in bryophytes: An experimental study. – *Ecotoxicology and Environmental Safety* 73: 1776–1784.
- Michel P., Overton J. McC, Mason N. W. H., Hurst J. M. & Lee W. G. (2011): Species-environment relationships of mosses in New Zealand indigenous forest and shrubland ecosystems. – *Plant Ecology* 212: 353–367.
- Mierzeńska M. & Vončina G. (2010): New localities of the rare liverwort *Aneura maxima* (Schiffn.) Steph. (Metzgeriales, Marchantiophyta) in Poland (Pieninski National Park, Western Carpathians). – *Cryptogamie Bryologie* 31: 305–312.
- Migaszewski Z. M., Dołęgowska S., Halas S. & Trembaczowski A. (2010): Stable sulphur isotope ratios in the moss species *Hylocomium splendens* (Hedw.) B. S. G. and *Pleurozium schreberi* (Brid.) Mitt. from the Kielce area (south-central Poland). – *Isotopes In Environmental and Health Studies* 46: 219–224.
- Migaszewski Z. M., Galuszka A., Dołęgowska S., Crock J. G. & Lamothe P. J. (2010): Mercury in mosses *Hylocomium splendens* (Hedw.) BSG and *Pleurozium schreberi* (Brid.) Mitt. from Poland and Alaska: Understanding the origin of pollution sources. – *Ecotoxicology and Environmental Safety* 73: 1345–1351.
- Mikulášková, E. 2009. Biology, ecology and expansion of *Campylopus introflexus* in the Czech Republic. – *Nowellia Bryologica* numéro spécial 156–157.
- Miller G. R., Geddes C. & Mardon D. K. (2010): Effects of excluding sheep from an alpine dwarf-herb community. – *Plant Ecology & Diversity* 3: 87–93.
- Miller N. G. & Robinson S. C. (2010): Introduction and recent range expansion in the moss *Ptychomitrium serratum* (Ptychomitriaceae) in the Southern and Eastern United States. – *Botany–Botanique* 88: 336–344.
- Milyutina I. A., Goryunov D. V., Ignatov M. S., Ignatova E. A. & Troitsky A. V. (2010): The phylogeny of *Schistidium* (Bryophyta, Grimmiaceae) based on the primary and secondary structure of nuclear rDNA internal transcribed spacers. – *Molecular Biology* 44: 883–897.
- Montero A. & Dieguez C. (2010): Devonian floral assemblages and plant megafossils from the Iberian Peninsula: a review. – *Review of Palaeobotany and Palynology* 162: 231–238.
- Mulec J. & Kubešová S. (2010): Diversity of bryophytes in show caves in Slovenia and relation to light intensities. – *Acta Carsologica* 39: 587–596.
- Nascimbene J., Spitale D., Thues H. & Cantonati M. (2011): Congruencies between photoautotrophic groups in springs of the Italian Alps: implications for conservation strategies. – *Journal of Limnology* 70, Suppl. 1: 3–8.
- Nasrulhaq-B. A., Mohamed M. A. H., Lim A. L., Barakbah S. S., Yong K. T. & Nor D. M. (2011): Comparative morphological and photosynthetic studies on three Malaysian species of *Pogonatum* from habitats of varying light irradiances. – *Journal of Bryology* 33: 35–41.

- Neta T., Cheng Q., Bello R. L. & Hu B. (2010): Lichens and mosses moisture content assessment through high-spectral resolution remote sensing technology: a case study of the Hudson Bay Lowlands, Canada. – *Hydrological Processes* 24: 2617–2628.
- Neta T., Cheng Q., Bello R. L. & Hu B. (2011): Development of new spectral reflectance indices for the detection of lichens and mosses moisture content in the Hudson Bay Lowlands, Canada. – *Hydrological Processes* 25: 933–944.
- Newsham K. K. & Bridge P. D. (2010): Sebacinales are associates of the leafy liverwort *Lophozia excisa* in the southern maritime Antarctic. – *Mycorrhiza* 20: 307–313.
- Nguyen Viet H., Frontasyeva Ma. V., Trinh Thi T. M., Gilbert D. & Bernard N. (2010): Atmospheric heavy metal deposition in Northern Vietnam: Hanoi and Thainguyen case study using the moss biomonitoring technique, INAA and AAS. – *Environmental Science and Pollution Research* 17: 1045–1052.
- Nishihira N., Shindo A., Saigusa M., Ono F., Matsushima Y., Mori Y., Takarabe K., Saini N. L. & Yamashita M. (2010): Preserving life of moss *Ptychomitrium* under very high pressure. – *Journal of Physics and Chemistry of Solids* 71: 1123–1126.
- Novotný I. (2009): The *Ceratodon* moss genus in the Czech Republic and Slovakia. – *Nowellia Bryologica* numéro spécial: 68–77.
- Núñez-Olivera E., Otero S., Tomás R., Fabón G. & Martínez-Abaigar J. (2010): Cyclic environmental factors only partially explain the seasonal variability of photoprotection and physiology in two mosses from an unforested headwater stream. – *Bryologist* 113: 277–291.
- O'Shea B. (2010): Mosses of Venezuela. – *Archive for Bryology* 75: 1–23.
- Ochyra R. & Bednarek-Ochyra H. (2010): *Kuerschneria*, a new pleurocarpous moss genus from Brazil. – *Nova Hedwigia Suppl.* 138, Suppl. 138: 25–29.
- Olsson S., Enroth J., Buchbender V., Hedenäs L., Huttunen S. & Quandt D. (2011): *Neckera* and *Thamnobryum* (Neckeraceae, Bryopsida): Paraphyletic assemblages. – *Taxon* 60: 36–50.
- Özdemir T., Apaydın G., Mendil D., Bulut V. N., Cengiz E., Gündoğdu A. & Aylıkçı V. (2010): Determination of some elements in moss samples from North Eastern Anatolia, Turkey. – *Asian Journal of Chemistry* 22: 346–352.
- Oztopcu-Vatan P., Savaroglu F., Filik-Isçen C., Kabadere S., İlhan S. & Uyar R. (2011): Antimicrobial and antiproliferative activities of *Homalothecium sericeum* (Hedw.) Schimp. extracts. – *Fresenius Environmental Bulletin* 20: 461–466.
- Papp B. & Erzberger P. (2010): Contributions to the bryophyte flora of Durmitor National Park, Montenegro. – *Nova Hedwigia Suppl.* 138: 147–163.
- Parolly G. & Zippel E. (2010): Phytogeographical composition of neotropical trunk-epiphytic bryophyte communities. Ecosociological studies in Ecuadorian bryophyte communities, VII. – *Nova Hedwigia Suppl.* 138: 243–258.
- Pata I. M. C., Pata S.-M. & Macoveanu M. (2010): Biomonitoring of atmospheric pollution of Fe and Zn using native epigeic mosses. – *Environmental Engineering and Management Journal* 9: 1217–1225.
- Patiño J. & González-Mancebo J. M. (2011): Exploring the effect of host tree identity on epiphyte bryophyte communities in different Canarian subtropical cloud forests. – *Plant Ecology* 212: 433–449.
- Patiño J., Hylander K. & González-Mancebo J. M. (2010): Effect of forest clear-cutting on subtropical bryophyte communities in waterfalls, on dripping walls, and along streams. – *Ecological Applications* 20: 1648–1663.
- Patiño J., Werner O. & González-Mancebo J. M. (2010): The impact of forest disturbance on the genetic diversity and population structure of a late-successional moss. – *Journal of Bryology* 32: 220–231.
- Pauliuk F., Müller J. & Heinken T. (2011): Bryophyte dispersal by sheep on dry grassland. – *Nova Hedwigia* 92: 327–341.
- Peck J. E. & Moldenke A. R. (2010): Identifying pathways of biological invasion: can commercial moss harbor potential stowaways? – *Biological Invasions* 12: 2395–2398.
- Pejin B., Vujisić L., Sabovljević M., Sabovljević A., Tesević V. & Vajs V. (2010): Preliminary analysis of fatty acid chemistry of *Kindbergia praelonga* and *Kindbergia stokesii* (Brachytheciaceae). – *Journal of the Serbian Chemical Society* 75: 1637–1640.
- Peng-cheng W., Crosby M. & He S. (2011): Moss Flora of China, English Version. Volume 5, Erpodiaceae to Climaciaceae. – Missouri Botanical Garden & Science Press, Beijing [432 pp.]
- Peralta D. F. & Yano O. (2010): Taxonomic treatment of the Polytrichaceae from Brazil. – *Bryologist* 113: 646–672.
- Pérez-Llamazares A., Aboal J. R., Carballeira A. & Fernández J. A. (2011): Cellular location of K, Na, Cd and Zn in the moss *Pseudoscleropodium purum* in an extensive survey. – *Science of the Total Environment* 409: 1198–1204.

- Pérez-Llamazares A., Fernández J. A., Aboal J. R., Giordano S. & Carballeira A. (2011): Evaluation of the efficacy of the sequential elution technique, by use of electron microscopy methods. – *Journal of Bryology* 33: 54–61.
- Petrone R. M., Solondz D. S., Macrae M. L., Gignac D. & Devito K. J. (2011): Microtopographical and canopy cover controls on moss carbon dioxide exchange in a western boreal plain peatland. – *Ecohydrology* 4: 115–129.
- Philippi G. (2010): Epiphytic bryophyte vegetation on stems of *Acer campestre* in SW Germany and the eastern part of France.. – *Herzogia* 23: 269–278.
- Plášek V. & Sawicki J. (2009): New endemic moss species from the Czech Republic. – *Nowellia Bryologica* numéro spécial 43–52.
- Plášek V. & Sawicki J. (2009): The moss *Orthotrichum moravicum*, an endemic species of Moravskoslezské Beskydy Mts. (Western Carpathians Mountain Range). – *Nowellia Bryologica* 38: 10–12.
- Popper Z. A. & Tuohy M. G. (2010): Beyond the green: Understanding the evolutionary puzzle of plant and algal cell walls. – *Plant Physiology* 153: 373–383.
- Pouliot R., Marchand-R. M., Rochefort L. & Gauthier G. (2010): Estimating moss growth in Arctic conditions: a comparison of three methods. – *Bryologist* 113: 322–332.
- Pressel S. & Duckett J. G. (2010): Cytological insights into the desiccation biology of a model system: moss protonemata. – *New Phytologist* 185: 944–963.
- Pressel S., Bidartondo M. I., Ligrone R. & Duckett J. G. (2010): Fungal symbioses in bryophytes: New insights in the Twenty First Century. – *Phytotaxa* 9: 238–253.
- Pressel S., P'ng K. M. Y. & Duckett J. G. (2010): a cryo-scanning electron microscope study of the water relations of the remarkable cell wall in the moss *Rhacocarpus purpurascens* (Rhacocarpaceae, Bryophyta). – *Nova Hedwigia* 91: 289–299.
- Pressel S., P'ng K. M. Y. & Duckett J. G. (2011): An ultrastructural study of the liverwort *Mizutania riccardioides* Furuki et Iwatsuki: new insights into its systematic affinities and unique surface ornamentation. – *Bryologist* 114: 28–37.
- Preston C. D., Harrower C. A. & Hill M. O. (2011): Distribution patterns in British and Irish liverworts and hornworts. – *Journal of Bryology* 33: 3–16.
- Preussing M., Olsson S., Schäfer-Verwimp A., Wickett N. J., Wicke S., Quandt D. & Nebel M. (2010): New insights in the evolution of the liverwort family Aneuraceae (Metzgeriales, Marchantiophyta), with emphasis on the genus *Lobatiriccardia*. – *Taxon* 59: 1424–1440.
- Price M. J. (2010): Lectotypification of the name *Tetraphis pellucida* Hedw. (Bryophyta). – *Candollea* 65: 15–19.
- Prigge M. J. & Bezanilla M. (2010): Evolutionary crossroads in developmental biology: *Physcomitrella patens*. – *Development* 137: 3535–3543.
- Proctor M. C. F. (2010): Recovery rates of chlorophyll-fluorescence parameters in desiccation-tolerant plants: fitted logistic curves as a versatile and robust source of comparative data. – *Plant Growth Regulation* 62: 233–240.
- Proctor M. C. F. & Smirnov N. (2011): Ecophysiology of photosynthesis in bryophytes: major roles for oxygen photoreduction and non-photochemical quenching? – *Physiologia Plantarum* 141: 130–140.
- Proust H., Hoffmann B., Xie X., Yoneyama K., Schaefer D. G., Yoneyama K., Nogue F. & Rameau C. (2011): Strigolactones regulate protonema branching and act as a quorum sensing-like signal in the moss *Physcomitrella patens*. – *Development* 138: 1531–1539.
- Puglisi M. (2010): Contribution to the bryophyte vegetation of the Pelagian Archipelago the Island of Linosa (Sicily Channel, Italy). – *Nova Hedwigia Suppl.* 138, Suppl. 138: 259–284.
- Puglisi M., Privitera M. & Di Pietro R. (2011): New records of arctic montane and alpine mosses from Central Italy. – *Cryptogamie Bryologie* 32: 75–81.
- Purvis O. W., Tittley I., Chimonides P. D. J., Bamber R., Hayes P. A., James P. W., Rumsey F. J. & Read H. (2010): Long-term biomonitoring of lichen and bryophyte biodiversity at Burnham Beeches SAC and global environmental change. – *Systematics and Biodiversity* 8: 193–208.
- Pyšek P., Jarošík V., Hulme P. E., Kühn I., Wild J., Arianoutsou M., Bacher S., Chiron F., Didžiulis V., Essl F., Genovesi P., Gherardi F., Hejda M., Kark S., Lambdon P. W., Desprez-Loustau M.-L., Nentwig W., Pergl J., Poboljšaj K., Rabitsch W., Roques A., Roy D. B., Shirley S., Solarz W., Vilà M. & Winter M. (2010): Disentangling the role of environmental and human pressures on biological invasions across Europe. – *Proceedings of the National Academy of Sciences of the United States of America* 107: 12157–12162.

- Raabe S., Müller J., Manthey M., Dürhammer O., Teuber U., Göttlein A., Förster B., Brandl R. & Bässler C. (2010): Drivers of bryophyte diversity allow implications for forest management with a focus on climate change. – *Forest Ecology and Management* 260: 1956–1964.
- Ramaiya M., Johnson M. G., Shaw B., Heinrichs J., Hentschel J., von Konrat M., Davison P. G. & Shaw A. J. (2010): Morphologically cryptic biological species within the liverwort *Frullania asagrayana*. – *American Journal of Botany* 97: 1707–1718.
- Ramsay H. P. (2010): Scanning electron microscope studies on four species of *Ulota* from Australia. – *Botany–Botanique* 88: 435–438.
- Rancura S., Ribeiro M. M. & Nordi N. (2010): Considerations on harvesting *Sphagnum* in Cananéia municipality, São Paulo state, Brazil. – *Acta Botanica Brasilica* 24: 328–334.
- Raymond B. A., Bassingthwaite T. & Shaw D. P. (2010): Measuring nitrogen and sulphur deposition in the Georgia Basin, British Columbia, using lichens and moss. – *Journal of Limnology* 69, Suppl. 1: 22–32.
- Reiner-Drehwald M. E. (2010): On *Lejeunea* subgenus *Nanolejeunea* (Lejeuneaceae, Jungermanniopsida). – *Nova Hedwigia Suppl.* 138: 117–128.
- Reiner-Drehwald M. E., Rubies M. & Schiavone M. M. (2010): a tribute to Gabriela Hässel de Menéndez (1927–2009). – *Nova Hedwigia* 91: 279–288.
- Ren H., Ma G., Zhang Q., Guo Q., Wang J. & Wang Z. (2010): Moss is a key nurse plant for reintroduction of the endangered herb, *Primulina tabacum* Hance. – *Plant Ecology* 209: 313–320.
- Rivera M., Zechmeister H., Medina-Ramón M., Basagaña X., Foraster M., Bouso L., Moreno T., Solanas P., Ramos R., Koellensperger G., Deltell A., Vizcaya D. & Künzli N. (2011): Monitoring of heavy metal concentrations in home outdoor air using moss bags. – *Environmental Pollution* 159: 954–962.
- Root H. T. & McCune B. (2010): Forest floor lichen and bryophyte communities in thinned *Pseudotsuga menziesii* – *Tsuga heterophylla* forests. – *Bryologist* 113: 619–630.
- Rooy J. van & van Wyk A. E. (2011): The bryofloristic elements of southern Africa. – *Journal of Bryology* 33: 17–29.
- Ross J. J. & Reid J. B. (2010): Evolution of growth-promoting plant hormones. – *Functional Plant Biology* 37: 795–805.
- Rowntree J. K., Cowan R. S., Leggett M., Ramsay M. M. & Fay M. F. (2010): Which moss is which? Identification of the threatened moss *Orthodontium gracile* using molecular and morphological techniques. – *Conservation Genetics* 11: 1033–1042.
- Rozhina V. I., Ignatov M. S., Ignatova E. A. & Napreenko M. G. (2010)[2011]: *Dicranella staphylina* (Dicranaceae, Bryophyta) in Russia. – *Arctoa* 19: 135–138.
- Rubasinghe S. C. K., Milne R., Forrest L. L. & Long D. G. (2011): Realignment of the genera of Cleveaceae (Marchantiopsida, Marchantiidae). – *Bryologist* 114: 116–127.
- Rudolphi J. & Gustafsson L. (2011): Forests regenerating after clear-cutting function as habitat for bryophyte and lichen species of conservation concern. – *PLoS ONE* 6: e18639.
- Rydberg J., Karlsson J., Nyman R., Wanhatalo I., Näthe K. & Bindler R. (2010): Importance of vegetation type for mercury sequestration in the northern Swedish mire, Rödmosamyran. – *Geochimica et Cosmochimica Acta* 74: 7116–7126.
- Rydgren K., Halvorsen R. & Cronberg N. (2010): Infrequent sporophyte production maintains a female-biased sex ratio in the unisexual clonal moss *Hylocomium splendens*. – *Journal of Ecology* 98: 1224–1231.
- Sabovljević A., Sabovljević M. & Grubisić D. (2010): Gibberellin influence on the morphogenesis of the moss *Bryum argenteum* Hedw. in *in vitro* conditions. – *Archives of Biological Sciences* 62: 373–380.
- Sabovljević A., Sabovljević M. & Vukojević V. (2010): Effects of different cytokinins on chlorophyll retention in the moss *Bryum argenteum* (Bryaceae). – *Periodicum Biologorum* 112: 301–305.
- Sabovljević A., Soković M., Glamoclija J., Cirić A., Vujčić M., Pejin B. & Sabovljević M. (2010): Comparison of extract bio-activities of in-situ and in vitro grown selected bryophyte species. – *African Journal of Microbiology Research* 4: 808–812.
- Sabovljević M. & Frahm J.-P. (2011): Genetic diversity of the relict moss *Rhytidium rugosum* (Hypnales) in Europe inferred from the ITS region (nrDNA). – *Biologia* 66: 42–49.
- Sabovljević M. & Marka J. (2009): The biological evidence of climate changes: a case study of liverwort *Lunularia cruciata* (L.) Dum. ex Lindb. in Serbia. – *Botanica Serbica* 33: 185–187.
- Sabovljević M., Papp B. & Szurdoki E. (2010): New bryophyte records to some countries of the South-Eastern Europe. – *Cryptogamie Bryologie* 31: 289–292.
- Sabovljević M., Vujčić M. & Sabovljević A. (2010): Diversity of saproxylic bryophytes in old-growth and managed beech forests in the central Balkans. – *Plant Biosystems* 144: 234–240.
- Sadler K. D. & Bradfield G. E. (2010): Ecological facets of plant species rarity in rock outcrop ecosystems of the Gulf Islands, British Columbia. – *Botany–Botanique* 88: 429–434.

- Samecka-Cymerman A., Kolon K. & Kempers A. J. (2010): Influence of *Quercus robur* throughfall on elemental composition of *Pleurozium schreberi* (Brid.) Mitt. and *Hypnum cupressiforme* Hedw. – Polish Journal of Environmental Studies 19: 763–769.
- Samecka-Cymerman A., Wojtuń B., Kolon K. & Kempers A. J. (2011): *Sanionia uncinata* (Hedw.) Loeske as bioindicator of metal pollution in polar regions. – Polar Biology 34: 381–388.
- Sassmann S., Wernitznig S., Lichtscheidl I. K. & Lang I. (2010): Comparing copper resistance in two bryophytes: *Mielichhoferia elongata* Hornsch. versus *Physcomitrella patens* Hedw.. – Protoplasma 246: 119–123.
- Sawicki J., Plášek V. & Szczecińska M. (2010): Molecular studies resolve *Nyholmiella* (Orthotrichaceae) as a separate genus. – Journal of Systematics and Evolution 48: 183–194.
- Schäfer-Verwimp A. (2010): a checklist of the liverworts and hornworts of Dominica, West Indies. – Cryptogamie Bryologie 31: 313–415.
- Schirmel J. (2011): Response of the grasshopper *Myrmeleotettix maculatus* (Orthoptera: Acrididae) to invasion by the exotic moss *Campylopus introflexus* in acidic coastal dunes. – Journal of Coastal Conservation 15: 159–162.
- Schirmel J., Timler L. & Buchholz S. (2011): Impact of the invasive moss *Campylopus introflexus* on carabid beetles (Coleoptera: Carabidae) and spiders (Araneae) in acidic coastal dunes at the southern Baltic Sea. – Biological Invasions 13: 605–620.
- Schmidt A. R., Hentschel J. & Heinrichs J. (2010): The fossil hornwort described from Dominican amber is an angiosperm flower. – Review of Palaeobotany and Palynology 160: 209–211.
- Schramm J. & Wollert H. (2010): The bryophyte flora of an unspoiled percolation mire of northern Germany in the valley of the Blinde Trebel (Mecklenburg-Western Pomerania). – Herzogia 23: 257–267.
- Schröder W. & Pesch R. (2010): Long-term monitoring of the metal accumulation in forests measured by use of the moss technique. – European Journal of Forest Research 129: 475–488.
- Schröder W., Holy M., Pesch R., Harmens H., Fagerli H., Alber R., Coşkun M., De Temmerman L., Frolova M., González-Miqueo L., Jeran Z., Kubin E., Leblond S., Liiv S., Maňková B., Piispanen J., Santamaría J. M., Simonè P., Suchara I., Yurukova L., Thöni L. & Zechmeister H. G. (2010): First Europe-wide correlation analysis identifying factors best explaining the total nitrogen concentration in mosses. – Atmospheric Environment 44: 3485–3491.
- Schröder W., Holy M., Pesch R., Harmens H., Ilyin Il., Steinnes E., Alber R., Aleksiyenak Y., Blum O., Coşkun M., Dam M., De Temmerman L., Frolova M., Frontasyeva M., Gonzalez M. L., Grodzińska K., Jeran Z., Korzekwa S., Krmar M., Kubin E., Kvietkus K., Leblond S., Liiv S., Magnússon S., Maňková B., Piispanen J., Rühling Å., Santamaría J., Spiric Z., Suchara I., Thöni L., Urumov V., Yurukova L. & Zechmeister H. G. (2010): Are cadmium, lead and mercury concentrations in mosses across Europe primarily determined by atmospheric deposition of these metals? – Journal of Soils and Sediments 10: 1572–1584.
- Schuette S. & Renzaglia K. S. (2010): Development of multicellular spores in the hornwort genus *Dendroceros* (Dendrocerotaceae, Anthocerotophyta) and the occurrence of endospory in Bryophytes. – Nova Hedwigia 91: 301–316.
- Segreto R., Hassel K., Bardal R. & Stenøien H. K. (2010): Desiccation tolerance and natural cold acclimation allow cryopreservation of bryophytes without pretreatment or use of cryoprotectants. – Bryologist 113: 760–769.
- Seppelt R. D., Türk R., Green T. G. A., Moser G., Pannowitz S., Sancho L. G. & Schröter B. (2010): Lichen and moss communities of Botany Bay, Granite Harbour, Ross Sea, Antarctica. – Antarctic Science 22: 691–702.
- Sérgio C. & Garcia C. (2011): Bryophyte flora of São Tomé e Príncipe Archipelago (West Africa): annotated catalogue. – Cryptogamie Bryologie 21: 145–196.
- Sérgio C., Sim-Sim M. & Luís L. (2010): *Atrichum androgynum* (Mull. Hal.) A. Jaeger in Portugal, Azores and Madeira, new to the European bryoflora. – Cryptogamie Bryologie 31: 281–288.
- Sérgio C., Vieira C., Claro D. & Garcia C. (2011): *Conocephalum salebrosum* Szwedkowski, Buczkowska & Odrzykoski (Marchantiopsida): modelling the occurrence of a hygrophytic species new to the bryoflora of Portugal, Madeira and Azores. – Journal of Bryology 33: 30–34.
- Shaw A. J., Cox C. J., Buck W. R., Devos N., Buchanan A. M., Cave L., Seppelt R., Shaw B., Larrain J., Andrus R., Greilhuber J. & Tensch E. M. (2010): Newly resolved relationships in an early land plant lineage: bryophyta class Sphagnopsida (peat mosses). – American Journal of Botany 97: 1511–1531.
- Shaw A. J., Devos N., Cox C. J., Boles S. B., Shaw B., Buchanan A. M., Cave L. & Seppelt R. (2010): Peatmoss (*Sphagnum*) diversification associated with Miocene Northern Hemisphere climatic cooling? – Molecular Phylogenetics and Evolution 55: 1139–1145.

- Shaw A. J., Szövényi P. & Shaw B. (2011): Bryophyte diversity and evolution: windows into the early evolution of land plants. – *American Journal of Botany* 98: 352–369.
- Shmakova N. Yu. & Markovskaya E. F. (2010): Photosynthetic pigments of plants and lichens inhabiting Arctic tundra of West Spitsbergen. – *Russian Journal of Plant Physiology* 57: 764–769.
- Sim-Sim M., Bergamini A., Luís L., Fontinha S., Martins S., Lobo C. & Stech M. (2011): Epiphytic bryophyte diversity on Madeira Island: Effects of tree species on bryophyte species richness and composition. – *Bryologist* 114: 142–154.
- Sim-Sim M., Fontinha S., Luís L., Lobo C. & Stech M. (2010): The Selvagens Islands bryoflora and its relation with islands of the Madeira and Canary Islands Archipelagos. – *Nova Hedwigia Suppl.* 138: 187–199.
- Singh M., Singh S., Nath V., Sahu V. & Rawat A. K. S. (2011): Antibacterial activity of some bryophytes used traditionally for the treatment of burn infections. – *Pharmaceutical Biology* 49: 526–530.
- Singh S. M., Pereira N. & Ravindra R. (2010): Adaptive mechanisms for stress tolerance in Antarctic plants. – *Current Science* 99: 334–340.
- Sklenář P., Kovář P., Palice Z., Stančík D. & Soldán Z. (2010): Primary succession of high-altitude Andean vegetation on lahars of Volcán Cotopaxi, Ecuador. – *Phytocoenologia* 40: 15–28.
- Skripnikov A. Yu., Anikanov N. A., Kazakov V. S., Dolgov S. V., Ziganshin R. Kh., Govorun V. M., Ivanov V. T. (2011): The search for and identification of peptides from the moss *Physcomitrella patens*. – *Russian Journal of Bioorganic Chemistry* 37: 95–104.
- Słowińska S., Słowiński M. & Lamentowicz M. (2010): Relationships between local climate and hydrology in *Sphagnum* mire: implications for palaeohydrological studies and ecosystem management. – *Polish Journal of Environmental Studies* 19: 779–787.
- Smart S. M., Henrys P. A., Scott W. A., Hall J. R., Evans C. D., Crowe A., Rowe E. C., Dragosits U., Page T., Whyatt J. D., Sowerby A. & Clark J. M. (2010): Impacts of pollution and climate change on ombrotrophic *Sphagnum* species in the UK: analysis of uncertainties in two empirical niche models. – *Climate Research* 45: 163–507.
- Smart S. M., Scott W. A., Whitaker J., Hill M. O., Roy D. B., Critchley C. N., Marini L., Evans C., Emmett B. A., Rowe E. C., Crowe A., Le Duc M. & Marrs R. H. (2010): Empirical realised niche models for British higher and lower plants – development and preliminary testing. – *Journal of Vegetation Science* 21: 643–656.
- Söderström L., De Roo R. & Hedderson T. (2010): Taxonomic novelties resulting from recent reclassification of the Lophoziaceae/Scapaniaceae clade. – *Phytotaxa* 3: 47–53.
- Söderström L., Gradstein S. R. & Hagborg A. (2010): Checklist of the hornworts and liverworts of Java. – *Phytotaxa* 9: 53–149.
- Söderström L., Hagborg A., Váňa J. & von Konrat M. (2011): Land of wood and water: a checklist of liverworts and hornworts of Jamaica. – *Bryologist* 114: 67–91.
- Šoltés R., Školek J., Homolová Z. & Kyselová Z. (2010): Early successional pathways in the Tatra Mountains (Slovakia) forest ecosystems following natural disturbances. – *Biologia* 65: 958–964.
- Song L., Liu W.-Y., Ma W.-Z. & Tan Z.-H. (2011): Bole epiphytic bryophytes on *Lithocarpus xylocarpus* (Kurz) Markgr. in the Ailao Mountains, SW China. – *Ecological Research* 26: 351–363.
- Sørensen I., Domozych D. & Willats W. G. T. (2010): How have plant cell walls evolved? – *Plant Physiology* 153: 366–372.
- Soudzilovskaia N. A., Cornelissen J. H. C., During H. J., van Logtestijn R. S. P., Lang S. I. & Aerts R. (2010): Similar cation exchange capacities among bryophyte species refute a presumed mechanism of peatland acidification. – *Ecology* 91: 2716–2726.
- Sparrius L. B. & Kooijman A. M. (2011): Invasiveness of *Campylopus introflexus* in drift sands depends on nitrogen deposition and soil organic matter. – *Applied Vegetation Science* 14: 221–229.
- Spence J. R. (2009): Nomenclatural changes in the Bryaceae (Bryopsida) for North America III. – *Phytologia* 91: 494–500.
- Spirina U. N. & Ignatov M. S. (2010)[2011]: On the branch development in *Leucodon* (Leucodontaceae, Bryophyta). – *Arctoa* 19: 139–150.
- Stark L. R., Brinda J. C. & McLetchie D. N. (2011): Effects of increased summer precipitation and N deposition on Mojave Desert populations of the biological crust moss *Syntrichia caninervis*. – *Journal of Arid Environments* 75: 457–463.
- Stark L. R., McLetchie D. N. & Eppley S. M. (2010): Sex ratios and the shy male hypothesis in the moss *Bryum argenteum* (Bryaceae). – *Bryologist* 113: 788–797.
- Stebel A., Ochyra R. & Vončina G. (2010): Mosses of the Pieniny Range (Polish Western Carpathians). – *Sorus*, Poznań. [114 pp.]



- Stech M. & Quandt D. (2010): 20,000 species and five key markers: The status of molecular bryophyte phylogenetics. – *Phytotaxa* 9: 196–228.
- Stech M., Kolvoort E., Loonen M. J. J. E., Vrieling K. & Kruijer J. D. (2011): Bryophyte DNA sequences from faeces of an arctic herbivore, barnacle goose (*Branta leucopsis*). – *Molecular Ecology Resources* 11: 404–408.
- Stech M., Sim-Sim M., Esquivel M. G., Luis L., Fontinha S., Lobo C., Garcia C., Martins S., Vieira C., Barroso J., Pedro L. G. & Figueiredo A. C. S. (2010): Molecular, phytochemical and morphological characterization of the liverwort genus *Radula* in Portugal (mainland, Madeira, Azores). – *Systematics and Biodiversity* 8: 257–268.
- Stech M., Werner O., González-Mancebo M. J., Patiño J., Sim-Sim M., Fontinha S., Hildebrandt I. & Ros R. M. (2011): Phylogenetic inference in *Leucodon* Schwägr. subg. *Leucodon* (Leucodontaceae, Bryophyta) in the North Atlantic region. – *Taxon* 60: 79–88.
- Stehn S. E., Webster C. R., Glime J. M. & Jenkins M. A. (2010): Elevational gradients of bryophyte diversity, life forms, and community assemblage in the southern Appalachian Mountains. – *Canadian Journal of Forest Research* 40: 2164–2174.
- Stenøien H. K., Shaw A. J., Shaw B., Hassel K. & Gunnarsson U. (2011): North American origin and recent European establishments of the amphi-atlantic peat moss *Sphagnum angermanicum*. – *Evolution* 65: 1181–1194.
- Stenøien H. K., Shaw A. J., Stengrundet K. & Flatberg K. I. (2011): The narrow endemic Norwegian peat moss *Sphagnum troendelagicum* originated before the last glacial maximum. – *Heredity* 106: 370–382.
- Stevens C., Dupré C., Gaudnik C., Dorland E., Dise N., Gowing D., Bleeker A., Alard D., Bobbink R., Fowler D., Vandvik V., Corcket E., Mountford J. O., Aarrestad P. A., Muller S. & Diekmann M. (2011): Changes in species composition of European acid grasslands observed along a gradient of nitrogen deposition. – *Journal of Vegetation Science* 22: 207–215.
- Suárez G. M. & Schiavone M. M. (2010): New combinations and range extension for Neotropical mosses. – *Bryologist* 113: 679–681.
- Suárez G. M. & Schiavone M. M. (2010): *Pohlia* section *Apalodictyon* (Bryaceae, Bryophyta) in Central and South America. – *Nova Hedwigia* 91: 377–388.
- Suárez G. M. & Schiavone M. M. (2011): *Pohlia* section *Pohlia* (Bryaceae, Bryophyta) in Central and South America. – *Nova Hedwigia* 92: 453–477.
- Suchara I., Sucharová J., Holá M., Reimann C., Boyd R., Filzmoser P. & Englmaier P. (2011): The performance of moss, grass, and 1- and 2-year old spruce needles as bioindicators of contamination: a comparative study at the scale of the Czech Republic. – *Science of the Total Environment* 409: 2281–2297.
- Sugier P. & Czarnecka B. (2010): Vascular plants versus mosses in lakeland and riverine mires in two regions of Eastern Poland. – *Polish Journal of Ecology* 58: 635–644.
- Swatland H. J. (2011): Microphotometry of underwater shadowing by a moss from a Niagara escarpment waterfall. – *Microscopy and Microanalysis* 17: 125–131.
- Szmeja J. (2010): Changes in the aquatic moss *Sphagnum denticulatum* Brid. population abundance in a softwater lake over a period of three years. – *Acta Societatis Botanicorum Poloniae* 79: 167–173.
- Szmeja J., Bociąg K. & Merdalski M. (2010): Effect of light competition with filamentous algae on the population dynamics and development of the moss species *Warnstorfia exannulata* in a soft water lake. – *Polish Journal of Ecology* 58: 221–230.
- Szövényi P., Rensing S. A., Lang D., Wray G. A. & Shaw A. J. (2011): Generation-biased gene expression in a bryophyte model system. – *Molecular Biology and Evolution* 28: 803–812.
- Taira T., Mahoe Y., Kawamoto N., Onaga S., Iwasaki H., Ohnuma T. & Fukamizo T. (2011): Cloning and characterization of a small family 19 chitinase from moss (*Bryum coronatum*). – *Glycobiology* 21: 644–654.
- Taylor T. N., Taylor E. L., Decombeix A.-L., Schwendemann A., Serbet R., Escapa I. & Krings M. (2010): The enigmatic Devonian fossil *Prototaxites* is not a rolled-up liverwort mat: comment on the paper by Graham et al. (*AJB* 97: 268-275). – *American Journal of Botany* 97: 1074–1078.
- Thébaud G., Goubet P., Skrzypczak R. & Sourp E. (2009): Plant communities of ombrotrophic mires in the eastern Massif central (France). – *Acta Botanica Gallica* 156: 341–377.
- Thöni L., Yurukova L., Bergamini A., Ilyin I. & Matthaei D. (2011): Temporal trends and spatial patterns of heavy metal concentrations in mosses in Bulgaria and Switzerland: 1990-2005. – *Atmospheric Environment* 45: 1899–1912.
- Tobias M. & Niinemets Ü. (2010): Acclimation of photosynthetic characteristics of the moss *Pleurozium schreberi* to among-habitat and within-canopy light gradients. – *Plant Biology* 12: 743–754.
- Tomaselli M., Spitale D. & Petraglia A. (2011): Phytosociological and ecological study of springs in Trentino (south-eastern Alps, Italy). – *Journal of Limnology* 70, Suppl. 1: 23–53.

- Tretiach M., Pittao E., Crisafulli P. & Adamo P. (2011): Influence of exposure sites on trace element enrichment in moss-bags and characterization of particles deposited on the biomonitor surface. – *Science of the Total Environment* 409: 822–830.
- Tuba Z., Saxena D. K., Srivastava K., Singh S., Czobel S. & Kalaji H. M. (2010): Chlorophyll a fluorescence measurements for validating the tolerant bryophytes for heavy metal (Pb) biomapping. – *Current Science* 98: 1505–1508.
- Tubanova D. Ya., Goryunov D. V., Ignatova E. A. & Ignatov M. S. (2010)[2011]: On the taxonomy of *Dicranum acutifolium* and *D. fuscescens* complexes (Dicranaceae, Bryophyta) in Russia. – *Arctoa* 19: 151–164.
- Turetsky M. R., Mack M. C., Hollingsworth T. N. & Harden J. W. (2010): The role of mosses in ecosystem succession and function in Alaska's boreal forest. – *Canadian Journal of Forest Research* 40: 1237–1264.
- Uzawa M. & Higuchi M. (2010): Comparative development of the sporophyte-gametophyte junction in six moss species. – *Journal of Plant Research* 123: 777–787.
- van Leeuwen J. L. (2010): Launched at 36,000g. – *Science* 329: 395–396.
- van Rooy J. & van Wyk A. E. (2011): The bryofloristic elements of southern Africa. – *Journal of Bryology* 33: 17–29.
- Váňa J., Lebouvier M., Ochyra R., Bednarek-Ochyra H. & Cykowska B. (2010): Two noteworthy records of hepatics from Ile Amsterdam in the South Indian Ocean. – *Nova Hedwigia Suppl.* 138: 231–241.
- Váňa J., Söderström L., Hagborg A. & von Konrat M. (2010): Nomenclatural novelties and lectotypifications in Gymnomitriaceae. – *Novon* 20: 225–227.
- Váňa J., Söderström L., Hagborg A., von Konrat M. & Engel J. J. (2010): Early Land Plants Today: Taxonomy, systematics and nomenclature of Gymnomitriaceae. – *Phytotaxa* 11: 1–80.
- Vanderpoorten A. & Shaw A. J. (2010): The application of molecular data to the phylogenetic delimitation of species in bryophytes: a note of caution. – *Phytotaxa* 9: 229–237.
- Vanderpoorten A., Gradstein S. R., Carine M. A. & Devos N. (2010): The ghosts of Gondwana and Laurasia in modern liverwort distributions. – *Biological Reviews* 85: 471–487.
- Varela Z., Fernández J. A., Aboal J. R., Real C. & Carballeira A. (2010): Determination of the optimal size of area to be sampled by use of the moss biomonitoring technique. – *Journal of Atmospheric Chemistry* 65: 37–48.
- Veljić M., Cirić A., Soković M., Janacković P. & Marin P. D. (2010): Antibacterial and antifungal activity of the liverwort (*Ptilidium pulcherrimum*) methanol extract. – *Archives of Biological Sciences* 62: 381–385.
- Verhoeven J. T. A., Beltman B., Dorland E., Robat S. A. & Bobbink R. (2011): Differential effects of ammonium and nitrate deposition on fen phanerogams and bryophytes. – *Applied Vegetation Science* 14: 149–157.
- Villarreal J. C., Cargill D. C., Hagborg A., Söderström L. & Renzaglia K. S. (2010): a synthesis of hornwort diversity: Patterns, causes and future work. – *Phytotaxa* 9: 150–166.
- Vilnet A. A., Konstantinova N. A. & Troitsky A. V. (2010)[2011]: Molecular insight on phylogeny and systematics of the Lophoziales, Scapaniaceae, Gymnomitriaceae and Jungermanniaceae. – *Arctoa* 19: 31–50.
- Virtanen R. & Crawley M. J. (2010): Contrasting patterns in bryophyte and vascular plant species richness in relation to elevation, biomass and Soay sheep on St Kilda, Scotland. – *Plant Ecology & Diversity* 3: 77–85.
- Vittoz P., Bayfield N., Brooker R., Elston D. A., Duff E. I., Theurillat J.-P. & Guisan A. (2010): Reproducibility of species lists, visual cover estimates and frequency methods for recording high-mountain vegetation. – *Journal of Vegetation Science* 21: 1035–1047.
- Vittoz P., Camenisch M., Mayor R., Miserere L., Vust M. & Theurillat J.-P. (2010): Subalpine-nival gradient of species richness for vascular plants, bryophytes and lichens in the Swiss Inner Alps. – *Botanica Helvetica* 120: 139–149.
- Volkmar U. & Knoop V. (2010): Introducing Intron Locus *cox1i624* for phylogenetic analyses in bryophytes: on the issue of takakia as sister genus to all other extant mosses. – *Journal of Molecular Evolution* 70: 506–518.
- von Cräutlein M., Korpelainen H., Pietiläinen M. & Rikkinen J. (2011): DNA barcoding: a tool for improved taxon identification and detection of species diversity. – *Biodiversity and Conservation* 20: 373–389.
- von Konrat M., Shaw A. J. & Renzaglia K. S. (2010): a special issue of *Phytotaxa* dedicated to Bryophytes: The closest living relatives of early land plants. – *Phytotaxa* 9: 5–10.
- von Konrat M., Söderström L. & Hagborg A. (2010): The Early Land Plants Today project (ELPT): a community-driven effort and a new partnership with *Phytotaxa*. – *Phytotaxa* 9: 11–21.

- von Konrat M., Söderström L., Renner M. A. M., Hagborg A., Briscoe L. & Engel J. J. (2010): Early Land Plants Today (ELPT): How many liverwort species are there? – *Phytotaxa* 9: 22–40.
- Vuori K.-M. & Helisten H. (2010): The use of aquatic mosses in assessment of metal pollution: appraisal of type-specific background concentrations and inter-specific differences in metal accumulation. – *Hydrobiologia* 656: 99–106.
- Waite M. & Sack L. (2011): Does global stoichiometric theory apply to bryophytes? Tests across an elevation × soil age ecosystem matrix on Mauna Loa, Hawaii. – *Journal of Ecology* 99: 122–134.
- Waite M. & Sack L. (2011): Shifts in bryophyte carbon isotope ratio across an elevation × soil age matrix on Mauna Loa, Hawaii: do bryophytes behave like vascular plants? – *Oecologia* 166: 11–22.
- Whitaker D. L. & Edwards J. (2010): *Sphagnum* moss disperses spores with vortex rings. – *Science* 329: 406–406.
- Whitton B. A. & Neal C. (2011): Organic phosphate in UK rivers and its relevance to algal and bryophyte surveys. – *Annales de Limnologie – International Journal of Limnology* 47: 3–10.
- Wodniok S., Brinkmann H., Glöckner G., Heidel A. J., Philippe H., Melkonian M. & Becker B. (2011): Origin of land plants: Do conjugating green algae hold the key? – *BMC Evolutionary Biology* 2011, 11: 104.
- Wolf L., Rizzini L., Stracke R., Ulm R. & Rensing S. A. (2010): The Molecular and Physiological Responses of *Physcomitrella patens* to Ultraviolet-B Radiation. – *Plant Physiology* 153: 1123–1134.
- Xiao H.-Y., Tang C.-G., Xiao H.-W., Liu X.-Y. & Liu C.-Q. (2010): Mosses indicating atmospheric nitrogen deposition and sources in the Yangtze river drainage basin, China. – *Journal of Geophysical Research – Atmospheres* 115: –D14301.
- Xiao H.-Y., Tang C.-G., Xiao H.-W., Liu X.-Y. & Liu C.-Q. (2010): Stable sulphur and nitrogen isotopes of the moss *Haplocladium microphyllum* at urban, rural and forested sites. – *Atmospheric Environment* 44: 4312–4317.
- Xiao H.-Y., Zhu R.-G., Lin B.-N. & Liu C.-Q. (2011): Sulfur isotopic signatures in rainwater and moss *Haplocladium microphyllum* indicating atmospheric sulfur sources in Nanchang City (SE China). – *Science of the Total Environment* 409: 2127–2132.
- Xu S.-J., Jiang P.-A., Wang Z.-W. & Wang Y. (2009): Crystal structures and chemical composition of leaf surface wax depositions on the desert moss *Syntrichia caninervis*. – *Biochemical Systematics and Ecology* 37: 723–730.
- Yao H., Song J., Liu C., Luo K., Han J., Li Y., Pang X., Xu H., Zhu Y., Xiao P. & Chen S. (2010): Use of ITS2 region as the universal DNA barcode for plants and animals. – *PLoS ONE* 5: e13102.
- Zander R. H. (2010): Taxon mapping exemplifies punctuated equilibrium and atavistic saltation. – *Plant Systematics & Evolution* 286: 69–90.
- Zare H., Akbarinia M., Hedenäs L. & Maassumi A. A. (2011): Eighteen mosses from the Hyrcanian forest region new to Iran. – *Journal of Bryology* 33: 62–65.
- Zechmeister H. G., Dullinger S., Koellensperger G., Ertl S., Lettner C. & Reiter K. (2011): Do metal concentrations in moss from the Zackenberg area, Northeast Greenland, provide a baseline for monitoring? – *Environmental Science and Pollution Research* 18: 91–98.
- Zedda L., Cogoni A., Flore F. & Brundu G. (2010): Impacts of alien plants and man-made disturbance on soil-growing bryophyte and lichen diversity in coastal areas of Sardinia (Italy). – *Plant Biosystems* 144: 547–562.
- Zhang J., Zhang Y.-M., Downing A., Wu N. & Zhang B.-C. (2011): Photosynthetic and cytological recovery on remoistening *Syntrichia caninervis* Mitt., a desiccation-tolerant moss from Northwestern China. – *Photosynthetica* 49: 13–20.
- Zheng Y., Xu M., Zhao J., Zhang B., Bei S. & Hao L. (2011): Morphological adaptations to drought and reproductive strategy of the moss *Syntrichia caninervis* in the Gurbantunggut Desert, China. – *Arid Land Research and Management* 25: 116–127.
- Zhu Y., Buck W. R., & Wang Y. (2010): a revision of *Entodon* (Entodontaceae) in East Asia. – *Bryologist* 113: 516–589.