

Absence of anthraquinone pigments is paraphyletic and a phylogenetically unreliable character in the *Teloschistaceae*

Jan VONDRÁK, Jaroslav ŠOUN, Olga VONDRÁKOVÁ, Alan M. FRYDAY,
Alexander KHODOSOVTSSEV and Evgeny A. DAVYDOV

Abstract: It has been suggested that the absence of anthraquinones is not a synapomorphic character, but appears independently in unrelated lineages of *Teloschistaceae*. We analyzed ITS nrDNA regions in species of the genus *Caloplaca* and present evidence for five such examples: the *Caloplaca cerina* group, *C. obscurella*, the *C. servitiana* group, the *C. xerica* group and the *C. variabilis* group (*Pyrenodesmia*). In some cases, loss of anthraquinones is observed only in individuals within ordinarily pigmented populations, but sometimes the loss covers whole lineages containing one or more species. Both situations are observed in the *C. servitiana* group. Loss of anthraquinones is always followed by the synthesis of 'alternative' pigments (often *Sedifolia*-grey). In the specimens with anthraquinone-containing apothecia studied, these pigments are not visible in apothecial sections after dissolving anthraquinones in K. Fully unpigmented apothecia have not been observed.

The *Caloplaca xerica* group is a newly established, infraspecific grouping of species related to, and similar to, *C. xerica*. The *Caloplaca servitiana* group is also newly established and represents an isolated lineage covering two rather different, but related species. *Caloplaca neotaurica* is described here as a new species with apothecia of two colour variants; orange-red (with anthraquinones) and grey (with *Sedifolia*-grey).

The genus *Huea* represents another taxon lacking anthraquinones within *Teloschistaceae*. The genera *Apatoplaca* and *Cephalophysis*, which lack anthraquinones, are tentatively placed in *Teloschistaceae*, but their phylogenetic identity has not been recognized. *Hueidea* is reported to have no anthraquinones, but its secondary metabolites should be studied further and its possible placement in *Teloschistaceae* assessed.

We suggest that *Caloplaca abbreviata* var. *lecideoides* and *C. celata* represent variants of *C. stillicidiorum* lacking anthraquinones.

Key words: *Apatoplaca*, *Caloplaca neotaurica*, *Cephalophysis*, *Huea*, *Hueidea*, lichens, phylogeny, taxonomy, UV-radiation

Accepted for publication 11 October 2011

J. Vondrák: Institute of Botany, Academy of Sciences, Zámek 1, Průhonice, CZ-25243, Czech Republic, and Faculty of Science, University of South Bohemia, Branišovská 31, CZ-370 05, České Budejovice, Czech Republic. Email: j.vondrak@seznam.cz

J. Šoun: Department of Botany, Faculty of Science, University of South Bohemia, Branišovská 31, České Budejovice, CZ-370 05, Czech Republic.

O. Vondráková: Institute of Steppe (Urals Branch of Russian Academy of Sciences), Pionerskaya st. 11, Orenburg, RF-460000, Russia.

A. M. Fryday: Herbarium, Department of Plant Biology, Michigan State University, East Lansing, MI 48824-1312, USA.

A. Khodosovtsev: Kherson State University, 40 Rokiv Zhovtnya str. 27, 73000 Kherson, Ukraine.

E. A. Davydov: Altai State University, Lenin Ave. 61, Barnaul, RF-656049, Russia; Tigirek State Reserve, Nikitina, 111, Barnaul, RF-656043, Barnaul, Russia.

Introduction

In one of the largest lichen families, *Teloschistaceae*, the majority of lichens produce yellow-orange-red anthraquinone pigments in the superficial tissues of their fruiting bodies and/or thallus (e.g., Søchting 1997). They are produced in various *Teloschistaceae* in 1) both thallus and apothecia, 2) in apothecial disc and margin only, 3) in apothecial disc only or 4) are entirely absent. Anthraquinones are known to protect lichens from absorption of UV radiation (Solhaug *et al.* 2003); their content in *Xanthoria parietina* (L.) Beltr. (*Teloschistaceae*) significantly increases with exposition to solar radiation (Gauslaa & McEvoy 2005).

Some *Teloschistaceae* also have the ability to synthesize green or grey pigments of unknown structure, for example, ‘*Lecidea-green*’ (Wetmore 1996; Arup *et al.* 2007) or ‘*Sedifolia-grey*’ (e.g., Wetmore 1996; Meyer & Printzen 2000), which may have the same function, that is, protection against UV radiation (e.g., Hauck *et al.* 2007). These pigments usually occur in parts of the thalli or apothecia where anthraquinones are absent, although both pigments may co-occur in the superficial tissues of the apothecia of some species. These species, for example, *Caloplaca conversa* (Kremp.) Jatta, *C. exsecuta* (Nyl.) Dalla Torre & Sarnth., *C. oleicola* (J. Steiner) van den Boom & Breuss and *C. pollinii* (A. Massal.) Jatta, may resemble those with an entire absence of anthraquinones, because their old apothecia turn black as they contain large amounts of melanin pigments. In these cases, the anthraquinones may be identified by the reactions of the apothecial sections (K+ strongly purple-violet, K → HCl+ yellow, N+ orange, N → K+ purple-violet-blue, N → K → HCl+ yellow; Vondrák *et al.* 2010a).

The absence of anthraquinones is known at various levels; their absence in the thallus has arisen in many lineages throughout *Teloschistaceae*, for example, the *C. cerina* group (Šoun *et al.* 2011). Occasional loss of anthraquinones in the thallus is also documented in individuals or populations of species that usually have a pigmented thallus (Søchting 1973; Vondrák *et al.* 2010b). Aberrant populations lacking anthraquinones completely are suggested in *C. tirolensis* Zahlbr. (Hansen *et al.* 1987; Søchting 1989) or *C. xerica* Poelt & Vězda (Poelt 1975).

Species (or individuals) of *Teloschistaceae* with a complete loss of yellow-orange-red pigments, sometimes grouped under *Pyrenodesmia* (*Caloplaca variabilis* group), have been studied since the 19th century but attempts to treat them systematically began with Magnusson (1950), followed by Wunder (1974) in the Old World and Wetmore (1994) in North America. More recently, several works have dealt with this group using single locus (ITS) molecular data (Arup & Grube 1999; Tretiach *et al.* 2003; Tretiach & Muggia

2006; Muggia *et al.* 2008; Vondrák *et al.* 2008; Xahidin *et al.* 2010).

Although the absence of anthraquinones in *Teloschistaceae* has often been treated as an important grouping character, no studies to date have tested whether or not this is a reliable character for classification, or whether species with such characters represent a monophyletic group. The purpose of the present study is to test this hypothesis using a broad sampling of species lacking anthraquinones, which would traditionally be treated as closely related (e.g., Muggia *et al.* 2008). Specifically, we will test whether or not these species can be recovered as a well-supported monophyletic group.

Materials and Methods

Lichen samples were collected by the authors from various localities in the Northern Hemisphere. Detailed herbarium data for specimens used in our phylogenetic studies are given in Table 1. Data for paratypes of *C. neotaurica* are abbreviated; full information for samples from CBFS is available at <http://botanika.bf.jcu.cz/lichenology/data.php>.

Morphological examinations

A detailed morphological description is given only for the new species. Measurements are accurate to 0.25 µm for cells (e.g., conidia and ascospores), 1 µm for width of asci, or 10 µm for larger structures (e.g., hymenium, width of exciple). All measurements of cells (ascospores, conidia, asci, paraphyses) include their walls. Paraphyses tips were observed after treatment with c. 10% KOH. Only those ascospores with well-developed septa were measured; in these ascospores, loculi were connected with a thin cytoplasmic channel, never disconnected. Measurements with more than ten repeats ($n \geq 10$) are given as (min.–) $\bar{x} \pm SD$ (–max.), where \bar{x} = mean value and SD = standard deviation. Morphological terminology follows Smith *et al.* (2009).

Chemistry

The acetone extracts from apothecia were subjected to high-performance liquid chromatography (HPLC) analysis. Reverse phase column (C18, 5 µm, Lichrocart 250-4) was eluted with MeOH / 30%MeOH+1% H_3PO_4 for 77 min and the absorbances at 270 nm were recorded (for details see Søchting 1997). The compounds were determined on the basis of their retention times and absorption spectra. In the newly described species, five specimens were examined (CBFS JV2048, 5925, 6023, 7094, 7105).

TABLE 1. Sample data, engaging to groups and GenBank accession numbers of the ITS sequences newly generated for phylogenetic analyses.

Species	Specimen	Group	GenBank accession number
<i>Caloplaca albolutescens</i>	Greece: Crete: Sternes, on calcareous stone, 2005, <i>ř. Vondrák</i> (CBFS JV4098)	<i>C. xerica</i> group	JN641762
<i>C. aractina</i>	Sweden: Bohuslän: Askum par., Barlindarna, 2006, <i>U. Søchting</i> 10579 (C)	<i>C. haematites</i> group	JN641763
<i>C. atroflava</i>	Czech Republic: North Moravia: Šumperk, Raškov, on serpentinite, 2007, <i>Palice</i> 11741 (PRA)	<i>C. xerica</i> group	JN641764
<i>C. cerina</i> , grey	Greece: Astakos, on bark of <i>Olea</i> , 2010, <i>ř. Vondrák</i> (CBFS JV8329)	<i>C. cerina</i> group	JN641765
<i>C. cerina</i> , orange	Greece: Astakos, on bark of <i>Olea</i> , 2010, <i>ř. Vondrák</i> (CBFS JV8329)	<i>C. cerina</i> group	JN641766
<i>C. erythrocarpa</i>	Bulgaria: Strandzha Mts: Malko Tarnovo, on limestone, 2005, <i>ř. Vondrák</i> (CBFS JV3208)	<i>C. xerica</i> group	JN641767
<i>C. fuscoatroides</i>	Bulgaria: Black Sea coast: Burgas, Sozopol, 2007, <i>ř. Vondrák</i> (CBFS JV6468)	<i>C. xerica</i> group	JN641768
<i>C. haematites</i>	Russia: Caucasus: “Dagestanskij zapovednik” National park, 2009, <i>A. Gabibova</i> (CBFS)	<i>C. haematites</i> group	JN641769
<i>C. neotaurica</i>	Great Britain: Wales: seashore rocks, <i>U. Arup</i> (LD, material lost)	<i>C. xerica</i> group	JN641770
<i>C. neotaurica</i>	Greece: Peloponnese: 2010, on serpentinite, <i>ř. Vondrák</i> & <i>O. Vondráková</i> (CBFS JV8322)	<i>C. xerica</i> group	JN641771
<i>C. neotaurica</i>	Ukraine: Crimean Peninsula: Alushta, 2007, on siliceous stone, <i>J. Vondrák</i> (CBFS JV6023)	<i>C. xerica</i> group	JN641772
<i>C. neotaurica</i>	Ukraine: Crimean Peninsula: Karadag, 2007, on siliceous stone, <i>ř. Vondrák</i> (CBFS JV6229, isotype)	<i>C. xerica</i> group	JN641773
<i>C. neotaurica</i>	Ukraine: Crimean Peninsula: Yalta, on siliceous stone, 2009, <i>ř. Vondrák</i> & <i>A. Khodosovtsev</i> (CBFS JV7121)	<i>C. xerica</i> group	JN641774
<i>C. neotaurica</i> , grey	Ukraine: Crimean Peninsula: Alushta, on schist, 2007, <i>ř. Vondrák</i> (CBFS JV5475)	<i>C. xerica</i> group	JN641775
<i>C. neotaurica</i> , grey (1)	Greece: Peloponnese: Argolis Peninsula, on serpentinite, 2010, <i>ř. Vondrák</i> & <i>O. Vondráková</i> (CBFS JV8322)	<i>C. xerica</i> group	JN641776
<i>C. neotaurica</i> , grey (2)	Greece: Peloponnese: Argolis Peninsula, on serpentinite, 2010, <i>ř. Vondrák</i> & <i>O. Vondráková</i> (CBFS JV8322)	<i>C. xerica</i> group	JN641777
<i>C. servitiana</i>	Greece: Pindos Mts: Profitis Ilias, on bark of <i>Quercus coccifera</i> , 2005, <i>T. Spribille</i> 16225 (CBFS JV6974)	<i>C. servitii</i> group	JN641778
<i>C. servitiana</i>	Greece: Crete: Psiloritis Mts, upper end of Rouvas Gorge, on <i>Quercus coccifera</i> , 2004, <i>T. Spribille</i> 13700 (hb. T. Spribille)	<i>C. servitii</i> group	JN641779
<i>C. soralifera</i>	Ukraine: Khmel'nitsk' region: Kamenets-Podolsky, on limestone, 2006, <i>ř. Vondrák</i> (CBFS JV4594)	<i>C. xerica</i> group	JN641780
<i>C. aff. soralifera</i>	Czech Republic: Prachatice, on gneiss, 2009, <i>ř. Vondrák</i> & <i>O. Vondráková</i> (CBFS JV8325)	<i>C. xerica</i> group	JN641781

TABLE 1. *Continued*

Species	Specimen	Group	GenBank accession number
<i>Caloplaca</i> sp.	Russia: <i>Southern Ural Mts:</i> Orenburg, on limestone, 2009, <i>ř. Vondřák</i> (CBFS JV8182)	<i>Pyrenodesmia</i> 2	JN641784
<i>Caloplaca</i> sp.	Russia: <i>Southern Ural Mts:</i> Orenburg, on conglomerate, 2009, <i>ř. Vondřák</i> (CBFS JV8179)	<i>Pyrenodesmia</i> 2	JN641783
<i>Caloplaca</i> sp.	Russia: <i>Southern Ural Mts:</i> Orenburg, on conglomerate, 2009, <i>ř. Vondřák</i> (CBFS JV8326)	<i>Pyrenodesmia</i> 2	JN641782
<i>Caloplaca</i> sp.	Kazakhstan: <i>Mangistau region:</i> Beyney, on limestone, 2009, <i>ř. Vondřák</i> & <i>A. Khodosovtsev</i> (CBFS JV7635)	<i>Pyrenodesmia</i> 2	JN641785
<i>Caloplaca</i> sp.	Ukraine: <i>Doneck upland:</i> Luhansk, Rozkishne, on marl, 2007, <i>O. Nadyeina</i> (hb. <i>O. Nadyeina</i>)	<i>Pyrenodesmia</i> 2	JN641786
<i>Caloplaca</i> sp., grey	China: <i>Xinjiang:</i> Qinghe county, Mongolian Altai, west slope of Mt. Kara-Balchigtau, alt. 2400 m, on plant debris, 2007, <i>E. A. Davydov</i> 6876b (CBFS JV8413)	<i>C. servitii</i> group	JN641787
<i>Caloplaca</i> sp., yellow	China: <i>Xinjiang:</i> Qinghe county, Mongolian Altai, west slope of mt. Kara-Balchigtau, alt. 2400 m, on plant debris, 2007, <i>E. A. Davydov</i> 6876a (CBFS JV8414)	<i>C. servitii</i> group	JN641788
<i>C. stillicidiorum</i> , grey	USA: <i>Alaska:</i> Atkasuk, on bone, 2001, <i>A. Fryday</i> 8158 (MSC 57393)	<i>C. cerina</i> group	JN641789
<i>C. teicholyta</i>	Romania: <i>Dobrogea:</i> Tulcea, on limestone, 2007, <i>ř. Vondřák</i> (CBFS JV5381)	<i>C. xerica</i> group	JN641790
<i>C. teicholyta</i>	Ukraine: <i>Crimean Peninsula:</i> Bakhchisaray, on limestone, 2006, <i>ř. Vondřák</i> (CBFS JV5675)	<i>C. xerica</i> group	JN641791
<i>C. xerica</i>	Greece: <i>Crete:</i> Ano Viannos, on siliceous rock, 2005, <i>ř. Vondřák</i> (CBFS JV4119)	<i>C. xerica</i> group	JN641792
<i>C. xerica</i>	Czech Republic: <i>Křivoklátsko:</i> Hořovice, on siliceous rock, 2003, <i>ř. Vondřák</i> (CBFS JV1124)	<i>C. xerica</i> group	JN641793

Apothecial pigments were also examined by reactions in sections with Cl (C), KOH (K), HCl and HNO₃ (N) according to Meyer & Printzen (2000).

DNA extraction and amplification

Direct PCR was used for PCR-amplification of the ITS regions including the 5.8S gene of the nuclear rDNA following Arup (2006). Primers for amplification were ITS1F (Gardes & Bruns 1993) and ITS4 (White *et al.* 1990). PCR cycling parameters follow Ekman (2001).

Phylogenetic analyses

Sequences were aligned using MAFFT 6 (on-line version in the Q-INS-i mode; see Katoh *et al.* 2002) and

manually trimmed to eliminate the unaligned ends and ambiguously aligned regions of ITS1 and ITS2. Bayesian phylogenetic analyses were carried out using the program MrBayes 3.1.1 (Ronquist & Huelsenbeck 2003). The optimal nucleotide substitution models were found using the program MrModeltest v2.3 (Nylander 2004). The MCMC analyses were performed in two runs, each with four chains starting from a random tree and using the default temperature of 0.2. Every 100th tree was sampled and standard deviation of splits between runs less than 0.01 as a convergence criterion was used to assess burn-in. Models of nucleotide substitution, numbers of generations and discarded generations, numbers of sequences and lengths of each alignment are given in Table 2. Posterior probabilities are shown in all depicted phylogenetic trees (Figs 1–6). Tree terminals of samples

TABLE 2. Models of nucleotide substitution, numbers of generations and discarded generations, numbers of sequences and lengths of alignments for the molecular analyses.

Tree	Number of sequences	Number of retained positions	Substitution model	Number of generations	Number of discarded generations
<i>Caloplaca cerina</i> group	25	433	GTR+G	1×10^6	297 000
<i>Caloplaca obscurella</i>	21	528	GTR+I+G	1×10^6	259 000
<i>Caloplaca servitii</i> group	15	536	SYM+G	1×10^6	135 000
<i>Caloplaca xerica</i> group	21	477	GTR+G	10×10^6	726 000
<i>Pyrenodesmia</i> group	28	480	GTR+G	10×10^6	615 000
Teloschistaceae	33	398	GTR+I+G	10×10^6	1 582 000

without anthraquinones are in green (absence on population level) or yellow (absence on species level).

Results and Discussion

We generated 29 new sequences and used them in six different alignments. Five alignments were for specifically studied groups and one for a broad context. We found that the character of anthraquinone absence is polyphyletic, occurring in five different clades; the *Caloplaca cerina* group, *C. obscurella*, the *C. servitiana* group, the *C. xerica* group and the *C. variabilis* group (*Pyrenodesmia*). In some cases, whole lineages of anthraquinone-containing species contain one or more species that lack these pigments ('absence on species level'), whereas in others loss of yellow-orange pigments is restricted to individuals or populations of species that normally contain anthraquinones ('absence on population level'). In the latter case, specimens lacking anthraquinones always have some alternative pigment in their apothecia (usually *Sedifolia*-grey), which is not identified in apothecia with anthraquinones; that is, it is not apparent in the apothecial sections after dissolving the anthraquinones in K. Each case will be discussed specifically.

Caloplaca cerina group (absence on population level)

This homogeneous group (Šoun *et al.* 2011) clusters crustose *Teloschistaceae* with a white/

grey/black thallus without anthraquinones, and lecanorine apothecia with distinct thalline exciple lacking anthraquinones but with anthraquinone pigmented (yellow/orange/red) discs.

We have observed individuals entirely lacking anthraquinones in the Mediterranean lineage of *Caloplaca cerina* (Hedw.) Th. Fr. s. lat. (group A in Šoun *et al.* 2011) and in the arctic lineage of *Caloplaca stillicidiorum* (Vahl) Lynge (group 3 in Šoun *et al.* 2011). Both examples are morphologically similar to their closest relatives; important similarities are lecanorine exciples, distinct cortex in lower exciple and presence of *Sedifolia*-grey pigment in the thalline exciple. The arctic sample has smaller ascospores (*c.* $10\text{--}15 \times 6\text{--}8 \mu\text{m}$) than the phylogenetically closest observed specimens of *C. stillicidiorum* with orange discs (*c.* $13\text{--}16 \times 7\text{--}9 \mu\text{m}$). Both samples differ from other specimens in the *C. cerina* group by the presence of *Sedifolia*-grey in the apothecial disc (in section: K+ violet, C+ orange-carmine, N+ orange-red-purple); we have not observed this pigment in orange, anthraquinone-containing discs. Both specimens investigated are placed into the named groups (PP = 1.00 in both; Fig. 1A). In the available material, non-anthraquinone phenotypes occur side by side with their counterparts with orange discs (Fig. 1B & C).

The arctic specimen of *C. stillicidiorum* lacking anthraquinones is probably conspecific with *C. celata* Th. Fr., which appears to




FIG. 1. *Caloplaca cerina* group. A, Bayesian phylogeny (ITS nrDNA regions) of a part of the group containing lineages of *C. cerina* and *C. stillicidiorum* with individuals lacking anthraquinones (in green); B, two phenotypes of *C. cerina* (CBFS JV8329), grey apothecia without anthraquinones (left) and normally coloured apothecia (right); C, *C. stillicidiorum* (Fryday 8158; MSC) with grey, white pruinose apothecia (right) and with orange, normally coloured apothecia (left) and with *C. cf. tirolensis* in the upper part. Scales: B & C = 1 mm.

have the same ecology and identical morphology (Magnusson 1950; Poelt 1955; Hansen et al. 1987; Søchting et al. 2008). *Caloplaca abbreviata* var. *lecideoides* H. Magn., a lignicolous lichen from the Russian arctic, seems to be another name referring to *C. stillicidiorum* lacking anthraquinones; the type specimen in S (<http://andor.nrm.se/kryptos/kbo/kryptobase/200803/max/32948.jpg>) apparently contains a mixed population with orange and grey apothecia.

The description of *Pyrenodesmia monacensis* Leder., currently known as *Caloplaca monacensis* (Leder.) Lettau, is also based on non-anthraquinone containing specimens (Lederer 1896). Nevertheless, our data have shown that *C. monacensis* is a well-characterized, widely distributed and usually anthraquinone-containing species from the *C. cerina* group (Šoun et al. 2011).

***Caloplaca xerica* group (absence on population level)**

The '*Caloplaca xerica* group' represents a monophyletic group (PP = 1.00 in Fig. 2A) of mainly saxicolous species with the following characters: *thallus* without orange-red pigmentation, devoid of anthraquinones but with *Sedifolia*-grey in cortical tissues (K+ violet reaction in sections); *apothecia* orange to red, containing anthraquinones; *ascospores* never narrowly ellipsoid, length/breadth ratio 1.4–2.6; *pycnidia* grey-black, devoid of anthraquinones. The *C. xerica* group is related to *Pyrenodesmia* and the *C. haematites* complex (Fig. 3).

This species-rich group contains many unnamed taxa, one of which is *Caloplaca neotaurica* (Fig. 2B) that is formally described at the end of this paper. *Caloplaca neotaurica* is unusual within the group because it also has phenotypes with complete loss of anthraquinones, which usually occur along with phenotypes with red apothecia (Fig. 2C). Non-anthraquinone phenotypes are morphologically \pm identical to their counterparts with red apothecia, but all samples observed without anthraquinones had apothecia containing *Sedifolia*-grey (in section K+ violet, C+ orange-carmine, N+ orange-red-purple-black);

this pigment is present only in the thallus of the variant with red apothecia. The non-anthraquinone specimen from the Crimean Peninsula (CBFS JV6942) nests safely inside the *C. neotaurica* monophylum (PP = 1.00), but specimens examined from Greece have aberrant ITS sequences and the Bayesian phylogeny placed them as a sister group to other *C. neotaurica* samples (Fig. 2A).

Although he did not do so explicitly, Poelt (1975) already recognized the absence of anthraquinones at the population level within this group in *Caloplaca xerica*; its anthraquinone-lacking populations are named *C. xerica* var. *venostana*.

***Pyrenodesmia* group (absence on species level)**

The *C. variabilis* group (*Pyrenodesmia*) is a well-known group containing most of the European saxicolous species without anthraquinones but with *Sedifolia*-grey in the apothecia (in section K+ violet, C+ orange-carmine, N+ orange-red-purple-black). Our data show that *Pyrenodesmia* is unresolved within the *Caloplaca xerica* group and the *C. haematites* group (PP = 0.97 for the whole group; Fig. 3). The internal taxonomy and phylogeny of this group have been reported (Tretiach et al. 2003; Tretiach & Muggia 2006; Muggia et al. 2008; Vondrák et al. 2008), but the group is still only partially known in Europe and North America and it is little known in Asia. For instance, the group is species-rich in Central Asia (J. Vondrák, unpublished data). Based on our set of sequences, two main internal lineages are to be found within *Pyrenodesmia*; ***Pyrenodesmia* 1** with most of the European taxa (unsupported) and ***Pyrenodesmia* 2** containing mainly unknown taxa from the southern Ural Mountains, the Near East and the steppe-desert zone of the most SE region of Europe (PP = 0.97).

***Caloplaca obscurella* (absence on species level)**

Currently, we know little about the closest phylogenetic relatives of *Caloplaca obscurella*




FIG. 2. *Caloplaca neotaurica*. A, Bayesian phylogeny (ITS nrDNA regions) of the *C. xerica* group (delimited by the red line) with the monophylum of *C. neotaurica* (delimited by the blue line); species without anthraquinones in yellow; specimens of *C. neotaurica* without anthraquinones in green; B, *C. neotaurica*, isotype (CBFS JV6229); C, *C. neotaurica*, phenotype with anthraquinones (left) and phenotype with grey apothecia (right) in Peloponnese (CBFS JV8322). Scales: B & C = 1 mm.




FIG. 3. Bayesian phylogeny of ITS nrDNA regions showing the two groups of *Pyrenodesmia* in polytomy with the *C. haematites* group and the *C. xerica* group; species without anthraquinones highlighted in yellow.

(J. Lahm) Th. Fr. (Fig. 4A). It forms a strongly supported clade (PP = 1.00), sister to the large monophylum containing the *C. haematites* group, the *C. xerica* group and *Pyrenodesmia* (PP = 1.00), but we failed to find closer relatives. It is also unclear whether

the *C. obscurella* clade contains only a single species. We suggest that the North American species *C. brunneola* Wetmore, *C. camptidia* (Tuck.) Zahlbr. *C. dakotensis* Wetmore (Wetmore 1994) and *C. lecanoroides* Lendemer (Lendemer *et al.* 2010) might belong here




FIG. 4. *Caloplaca obscurella*. A, Bayesian phylogeny of ITS nrDNA regions; sister relationship of *C. obscurella* is shown to the clade containing *C. haematites* group, *C. xerica* group and *Pyrenodesmia* (delimited by the red square); species without anthraquinones highlighted in yellow; B, richly fertile specimen from Russia (CBFS JV7641). Scale = 1 mm.




FIG. 5. *Caloplaca servitiana* group. A, Bayesian phylogeny of ITS nrDNA regions containing *C. servitiana* with absence on species level (in yellow) and a species from China with absence on population level (in green); B, *C. servitiana* (Spribille 13705); C, species from China (Davydov 6876; CBFS) with yellow and grey apothecia growing side by side; D, species from China, the phenotype with grey apothecia (Davydov 6876b; CBFS) together with *C. stillicidiorum* (orange apothecia with white margins). Scales: B–D = 1 mm.

because all these species contain brown pigments in the epihyemium that resemble those in *C. obscurella* (K–, C–, N–, acetone insoluble, not detectable by HPLC).

***Caloplaca servitiana* group (absence on both species and population level)**

We revealed two species in the *Caloplaca servitiana* group (Fig. 5A); 1) *C. servitiana* Szatala s. str. (absence on species level; Fig. 5B) and 2) an unnamed species from China

(absence on population level; Fig. 5C & D). This group is the only one with the lack of anthraquinones found in the 'Lineage 1' of *Teloschistaceae* (sensu Gaya et al. 2008; Fig. 6). It is a rather isolated group with no known closely related lineages.

1) A brief description of *C. servitiana* is available in Vondrák et al. (2010a). It is known only from several sites in the eastern Mediterranean, where it grows on trees with various other *Caloplaca* species. Our comparison of this phenotype with co-occurring




FIG. 6. Bayesian phylogeny (ITS nrDNA regions) of *Teloschistaceae* with two sequences of *Physciaceae* as an outgroup. Representatives of the main lineages within the family have been selected for the tree terminals. Lineages lacking anthraquinones at the population level are in green and at species level in yellow.

Caloplaca species showed that it had a strong similarity with *C. aegatica* Giralt *et al.*, which challenged us to test if *C. aegatica* and *C. servitiana* are a single species. Our results show that *C. aegatica* is closely related to *C. cerinella* (Nyl.) Flagey, whereas *C. servitiana*

is not closely related to any known European species.

2) The unnamed specimen from the Altai Mountains (north-west China) is characterized by an inconspicuous, simplified thallus and biatorine to zeorine apothecia with a

distinct, *c.* 70–100 µm wide, proper exciple. The thalline exciple is usually reduced and restricted to the lower part of the apothecial margin. The ascospores, *c.* 14–18 × 7–9 µm, have a characteristic ‘obtuse rhomboid’ shape. In the specimen observed, the apothecia have two colour variants (Fig. 5C & D). The first is yellow, contains anthraquinones and lacks *Sedifolia*-grey, whereas the second is grey, devoid of anthraquinones, and contains *Sedifolia*-grey (in section K+ violet, C+ red, N+ red) in the epihymenium and superficial cells of the excipulum.

Other *Teloschistaceae* without anthraquinones

Among the genera of *Teloschistaceae*, the entire loss of anthraquinones is known only in the paraphyletic genus *Caloplaca* (Fig. 6). Species or populations with entire loss of anthraquinones are not known within foliose and fruticose genera of *Teloschistaceae*, although in some specimens of, for example, *Seirophora lacunosa* (Rupr.) Frödén and some *Teloschistes* species, whole thalli are grey with anthraquinones restricted to the apothecial discs and the walls of pycnidia (e.g., Søchting & Frödén 2002).

Our study has been carried out only on specimens from arctic-temperate-(subtropical) zones of the Northern Hemisphere, but species without anthraquinones are known from the tropics and the Southern Hemisphere: for example, *Caloplaca filsonii* Hafellner *et al.* (Australia; Kondratyuk *et al.* 2007), *C. gambiensis* Aptroot (Gambia; Aptroot 2001), *C. jatolensis* Y. Joshi & Upreti (India; Joshi *et al.* 2008), *C. lewis-smithii* (Antarctica; Søchting & Øvstedal 1998), *Caloplaca yammeraensis* S. Y. Kondr. *et al.* (Australia; Kondratyuk *et al.* 2009) and some tropical species with tri- to tetralocular ascospores (Hafellner & Poelt 1979). It is probable that these species belong to diverse groups that are unrelated to the lineages shown in this study.

Additional lineages with species or individuals that do not produce anthraquinones will also certainly be found in the Northern Hemisphere. We have examined the pig-

ments contained in herbarium specimens of some little-known arctic species, *Caloplaca atrocyanesces* (Th. Fr.) H. Olivier, *C. concinerasces* (Nyl.) H. Olivier and *C. diphyes* (Nyl.) H. Olivier. These species have blackish apothecia without anthraquinones, but contain different pigments (blue-green-olive) with various reactions with K, N and C in sections, but the identity of these pigments is unknown. The phylogenetic positions of these taxa are also not yet known. Other rare and little-known European species with black apothecia that contain unknown pigments are *C. concilians* (Nyl.) H. Olivier and *C. conciliascens* (Nyl.) Zahlbr. In addition, *Caloplaca sorocarpa* (Vain.) Zahlbr. has brown apothecia but is very rarely fertile and it is not known which pigments are present in its apothecia.

Caloplaca atosanguinea (G. Merr.) I. M. Lamb is a North American corticolous species which has brown (young) or black (old) apothecia. Its apothecial pigment is violet-blue in section and reacts: K+ violet-blue intensifying, K → HCl+ sordid green, N– unchanged, N → K+ weekly purple → discoloured, N → K → HCl– discoloured, C– discoloured. These reactions do not correspond to anthraquinones, which are typically K → HCl+ strongly yellow. The HPLC chromatogram shows one major unknown substance with a short retention time (Rt = 13.6 min, which is less than any known anthraquinone in *Teloschistaceae*) but also a minor substance (Rt = 33.3 min) with the absorption spectrum similar to emodin that we identified as an unknown anthraquinone. *Caloplaca atosanguinea* is related to, but not within, the *C. cerina* group (J. Vondrák, unpublished data).

Caloplaca demissa (Körb.) Arup & Grube also lacks anthraquinones, but it has probably never been found with apothecia. If it is able to produce apothecia, they may contain anthraquinones because its closest related species, *C. carphinea* (Fr.) Jatta (Gaya *et al.* 2008), has anthraquinone-containing apothecia.

Another lineage in *Teloschistaceae* that lacks anthraquinones is represented by the genus *Huea* C. W. Dodge & G. E. Baker (Dodge & Baker 1938). This genus was erected for a group of crustose Antarctic species of *Telo-*

schistaceae characterized by lacking anthraquinones, and having black apothecia with a carbonaceous exciple and a bright blue epihymenium. The status of *Huea* is controversial, with some authors accepting it (Øvstedal & Lewis Smith 2001) whereas others include the species in *Caloplaca* (e.g., Poelt & Hafellner 1980). Søchting *et al.* (2004) cited unpublished preliminary molecular data showing that the two species transferred to *Huea* by Dodge & Baker (1938) formed a distinct clade in *Caloplaca*. Unfortunately, there are problems with the typification of *Huea* because Dodge & Baker (1938) chose as the type species their newly described species *Huea flava* C. W. Dodge & G. E. Baker. This is usually considered to be a sterile crust of uncertain identity but was shown by Fryday (2011) to be conspecific with *Lecidea capsulata* C.W. Dodge & G. E. Baker. As this species is now included in the synonymy of *Carbonea vorticosa* (Flörke) Hertel, (<http://www.indexfungorum.org/Names/NamesRecord.asp?RecordID=107792>) this makes *Huea* an earlier name for *Carbonea*. Søchting *et al.* (2004) proposed a conserved type for *Huea* [*Huea coralligera* (Hue) C. W. Dodge & G. E. Baker] but, because they did not do so as a formal proposal, this has no relevance. The typification of *Huea* is dealt with in detail by Fryday (2011).

Apatoplaca Poelt & Hafellner and *Cephalophys* H. Kilius are monotypic genera with an absence of anthraquinones affiliated to *Teloschistaceae* mainly because of their *Teloschistaceae*-type asci (Poelt & Hafellner 1980; Kilius 1985); however, their phylogenetic identities are uncertain.

Also of interest is the genus *Hueidea* Kantvilas & P. M. McCarthy (Kantvilas & McCarthy 2003), which was erected for a single Australian species, *H. australiensis* Kantvilas & P. M. McCarthy, with characters intermediate between *Teloschistaceae* and *Fuscideaceae* V. Wirth & Vězda. The authors characterized their new genus as lacking thalline chemistry and showing all spot-test reactions negative, and placed it in *Fuscideaceae*. However, inspection of a specimen of *H. australiensis* revealed a weak K+ crimson reaction in the epihymenium,

suggesting the presence of anthraquinones and that the genus may be better placed in *Teloschistaceae*.

Conclusions

1. Non-anthraquinone lineages or populations represent a minor part of *Teloschistaceae* but they originated repeatedly in various sites in the phylogeny of the family (Fig. 6).
2. We have observed individuals without anthraquinones (absence on population level) in the *Caloplaca cerina* group, the *C. servitiana* group and the *C. xerica* group. Lineages containing non-anthraquinone species (absence on species level) were recognized in *Caloplaca obscurella*, the *C. servitiana* group and *Pyrenodesmia*.
3. Loss of anthraquinones is always followed by synthesis of alternative pigments (often *Sedifolia*-grey); fully unpigmented apothecia have not been observed by us. Anthraquinones and the alternative pigments may perform the same function, perhaps protection of ascospores against absorption of the UV-B radiation.

Taxonomy

Caloplaca neotaurica Vondrák, *Khodosovtsev, Arup & Søchting* sp. nov.

MycoBank No: MB563136

Caloplacae fuscoatroidis similis sed thallo ambitu squamulis nullis, tenui, atrogiseo vel nigrescenti, epruinoso vel raro pruinoso.

Typus: Ukraine, Crimean Peninsula, Sudak, Karadag Mts, Mt Svyataya, alt. 320 m, 44°56'03.27"N, 35°13'06.17"E, on volcanic rock, 24 May 2007, *Ĵ. Vondrák* (CBFS JV5925—holotypus; CBFS JV5960, 6022, 6229 & C—*isotypi*). ITS sequence of the *isotypus*: JN641773.

(Fig. 2)

Thallus thin, less than 150 (rarely 250) μm high, grey to brown-black, rarely white pruinose, indistinctly areolate; *areoles* (150–) $322 \pm 112(-600)$ μm wide ($n = 20$). *Prothallus* \pm present, dark grey. Marginal squamules and vegetative diaspores absent. *Cortex* absent; *alveolate cortex* (*sensu* Vondrák *et al.* 2009) present in spots (< 20 μm thick).

Apothecia <0.7 mm diam., biatorine, orange to red (grey in variants without anthraquinones), K+ purple, C+ purple, I- (disc I+ blue due to reaction with asci), P- (but red pigment dissolves in acetone solution of P), UV-; no visible spot reactions in grey apothecia. *Exciple* usually paler than the disc. *Proper exciple* c. 100–130 µm thick prosoplectenchymatous. *Thalline exciple* c. 60–120 µm thick. *Hymenium* c. 90–110 µm high. *Paraphyses* ± branched and anastomosed, with tips widened to (3.3–)4.4 ± 1.5(–6.0) µm (*n* = 10). *Asci* clavate, c. 55–75 × 18–26 µm. *Ascospores* (13.0–)14.7 ± 1.4(–17.3) × (7.0–) 8.9 ± 1.0(–11.3) µm (*n* = 25); ratio of ascospore length/breadth (1.4–)1.7 ± 0.2(–2.0); (*n* = 25); *ascospore septa* (3.5–) 5.7 ± 0.9(–8.0) µm wide (*n* = 25); ascospore septa/spore length (0.27–)0.39 ± 0.07(–0.54); (*n* = 25).

Pycnidia common, usually observable as darker spots in thallus (higher concentration of *Sedifolia*-grey around ostiole). *Conidia* ellipsoid c. 2.5–3.5 × 1.0–1.5 µm.

Chemistry. In apothecia, the main anthraquinone is 7–Cl-emodin; additional anthraquinones: 7–Cl-emodinal, 7–Cl-emodic acid, 7–Cl-citreorsein, ± citreorsein, ± emodinal, ± emodin (chemosyndrome C2, *sensu* Søchting 2001). *Sedifolia*-grey in thallus and in grey variants of apothecia (K+ violet, C+ orange-red, N+ purple in section).

Etymology. The species is very common in the Crimean Peninsula, which was known as 'Taurica' by the Greeks and Romans. The name *Caloplaca taurica* Mereschk. nomen nudum [= *C. inconnexa* (Nyl.) Zahlbr.] already exists and, although this is not validly published and so does not prevent the use of that name for our species, we have decided not to use it in order to avoid confusion.

Diagnostic characters. Thallus dark grey to brown-black (rarely white pruinose), thin, usually up to 150 µm thick; without marginal squamules and vegetative diaspores. Apothecia small (up to 0.7 mm), biatorine, orange-red, C+ purple (except in grey apothecial variants). Ascospores c. 14–17 × 7.5–10.5 µm, with septa c. 4.0–7.5 µm wide. Pycnidia

grey, conidia ellipsoid. Chemosyndrome C2 (details above).

Variability. Apart from two variants of apothecia (with/without anthraquinones), the species varies strongly in the content of *Sedifolia*-grey in the thallus. While northern specimens from Great Britain have a pale grey thallus with an indistinct amount of the grey pigment, specimens from the Mediterranean regions have a darker thallus, with an observable *Sedifolia*-grey content in sections.

Similar species. *Caloplaca atroflava* (Turner) Mong. differs in its paler and C- apothecia (absence of chlorinated anthraquinones). *Caloplaca crenularia* (With.) J. R. Laundon and related species have red pycnidial walls (with anthraquinones) and chemosyndrome C3 or C4 (with constant content of parietin and fragilin). *Caloplaca fuscoatroides* J. Steiner differs in its thick thallus (100–350 µm high), usual presence of marginal squamules (up to 2 mm diam.) and chemosyndrome C5 (with stable content of fragilin and absence of citreorsein and emodinal), while *Caloplaca xerica* Poelt & Vězda has a blastidiolate-lobulate thallus, ± zeorine apothecia and chemosyndrome C5 (as in *C. fuscoatroides*).

Chemosyndrome C2 (absence of parietin and fragilin and traces of citreorsein and emodinal) has been observed only in *Caloplaca ligustica* B. de Lesd. (Søchting 2001), which is probably unrelated to the new species having narrow ascospores with thin septa (Bouly de Lesdain 1936) similar to *C. subpallida* s. lat.

Phylogeny. Most sequences of the new species examined form a well-supported monophylum (PP = 1.00), but two sequences (not identical) of one Greek specimen with grey apothecia are somewhat different, and their relationship to the core clade is unsupported. Based on phenotype similarities, we decided to place the Greek specimen (containing also individuals with red apothecia) into *Caloplaca neotaurica*. The nearest related species may be a morphologically different *C. xerica*, but the relationship is unsupported.

Ecology and distribution. The species occurs on siliceous cliffs, outcrops or stones not far from the sea coast and is only rarely observed further inland (only in the Peloponnese and in the Rhodopes, Bulgaria). It is distributed in the Mediterranean and the Black Sea coast (especially in the Crimean Peninsula) and along the Atlantic coast of Europe (confirmed from Great Britain). Phenotypes with grey apothecia are known only from the Crimean Peninsula, Cyprus and Greece.

Specimens examined with red apothecia. **Bulgaria:** Black Sea coast: Burgas, Sinemorets, 2007, *Ĵ. Vondrák* (CBFS JV6497); Burgas, Sozopol, 2007, *Ĵ. Vondrák* (CBFS JV6452). Rhodopes: Momchilgrad, Bregovo, 2004, *Ĵ. Vondrák* (CBFS JV2043).—**Cyprus:** *Distr. Paphos:* Tripylos, 1987, *M. & H. Mayrhofer* 7750 (GZU).—**Great Britain:** *Wales:* V.C. 45 Permbrookshire: 1992, *U. Arup* (LD 1118360, 1118540, 1424694).—**Greece:** *Peloponnese:* Argolis Peninsula (Peiraias administrative district), Driopi, 2010, *Ĵ. Vondrák & O. Vondráková* (CBFS JV8322); Methana, Kypseli, 2010, *Ĵ. Vondrák & O. Vondráková* (CBFS JV8435); Thermisia, Vlachaiika, 2010, *Ĵ. Vondrák & O. Vondráková* (CBFS JV8436); Tripoli, Kastri, 2010, *Ĵ. Vondrák & O. Vondráková* (CBFS JV8438).—**Italy:** *Sardinia:* Núoro, Barbágia Seúlo, alt. 950–1080 m, 1986, *H. Mayrhofer* 6156, 5941 (GZU, sub *C. atroflava*).—**Turkey:** *Sea of Marmara coast:* Kapıdađı Peninsula, 2007, *Ĵ. Vondrák* (CBFS JV6459).—**Ukraine:** *Crimean Peninsula:* Alushta, Demerji Yayla, 2007, *Ĵ. Vondrák* (CBFS JV5996); Alushta, Kanakskaya balka, 2007, *Ĵ. Vondrák* (CBFS JV6023, 6041); Feodosia, Karadag, Mt Levinsona-Lessinga, 2000, *A. Khodosovtsev* (KHER 200, 201 sub *Caloplaca atroflava*); Feodosia, Mt Choba-Tepe, 2001, *A. Khodosovtsev* (KHER 2874, 2873 sub *Caloplaca scotoplaca*); Sudak, Dachnoe, 2008, *Ĵ. Vondrák* (CBFS JV7231); Sudak, Mt Echkedag, 2008, *Ĵ. Vondrák* (CBFS JV7213, 7232); Yalta, Foros, 2001, *A. V. Yena* (KHER 2889); Yalta, Oliva, 2009, *Ĵ. Vondrák* (CBFS JV7094, dupl. in C; 7103; 7105, dupl. in C; 7121).

Specimens examined with grey apothecia. **Cyprus:** *Distr. Laranca:* Stavrovouni, 1987, *M. & H. Mayrhofer* 7198 (GZU).—**Greece:** *Peloponnese:* Argolis Peninsula (Peiraias administrative district), Driopi, 2010, *Ĵ. Vondrák & O. Vondráková* (CBFS JV8322).—**Ukraine:** *Crimean Peninsula:* Alushta, Kanakskaya balka, 2007, *A. Khodosovtsev & Ĵ. Vondrák* (CBFS JV5475, KHER); *ibid.*, 2008, *A. Khodosovtsev & Ĵ. Vondrák* (CBFS JV6942, 7188, KHER); Feodosia, Karadag, 2001, *A. Khodosovtsev* (KHER 1452).

We are grateful to Toby Spribille, Ulrik Søchting and Ulf Arup for their comments on this paper. Our research received support from the Visegrad Fund (grants 51000067 and 51100753), the whole-Institute grant (AV0Z60050516) and the SYNTHESYS Project <http://www.synthesys.info/>, which is financed by European Community Research Infrastructure Action under the FP7 “Capacities” Program.

REFERENCES

- Aptroot, A. (2001) Lichens from Gambia, with a new black-fruited isidiate *Caloplaca* on savannah trees. *Cryptogamie, Mycologie* **22**: 265–270.
- Arup, U. (2006) A new taxonomy of the *Caloplaca citrina* group in the Nordic countries, except Iceland. *Lichenologist* **38**: 1–20.
- Arup, U. & Grube, M. (1999) Where does *Lecanora demissa* (Ascomycota, Lecanorales) belong? *Lichenologist* **31**: 419–430.
- Arup, U., Arneng, E. & Søchting, U. (2007) *Caloplaca fuscocorufa*—a misunderstood species in northern Europe. *Lichenologist* **39**: 409–414.
- Bouly de Lesdain, M. (1936) Notes lichénologiques. 29. *Bulletin de la Société Botanique de France* **83**: 5–9.
- Dodge, C. W. & Baker, G. E. (1938) The Second Byrd Antarctic Expedition. Botany. II. Lichens and lichen parasites. *Annals of the Missouri Botanical Garden* **25**: 515–718.
- Ekman, S. (2001) Molecular phylogeny of the *Bacidia-ceae* (Lecanorales, lichenized Ascomycota). *Mycological Research* **105**: 783–797.
- Fryday, A. M. (2011) How should we deal with the Antarctic and Subantarctic taxa published by Carroll William Dodge? *Opuscula Philolichenum* **9**: 89–98.
- Gardes, M. & Bruns, T. D. (1993) ITS primers with enhanced specificity for basidiomycetes. Application for the identification of mycorrhizae and rust. *Molecular Ecology* **2**: 113–118.
- Gauslaa, Y. & McEvoy, M. (2005) Seasonal changes in solar radiation drive acclimation of the sun-screening compound parietin in the lichen *Xanthoria parietina*. *Basic and Applied Ecology* **6**: 75–82.
- Gaya, E., Navarro-Rosinés, P., Llimona, X., Hladun, N. & Lutzoni, F. (2008) Phylogenetic reassessment of the *Teloschistaceae* (lichen-forming Ascomycota, Lecanoromycetes). *Mycological Research* **112**: 528–546.
- Hafellner, J. & Poelt, J. (1979) Die Arten der Gattung *Caloplaca* mit plurilocularen Sporen (*Meroplacis*, *Triophthalmidium*, *Xanthocarpia*). *Journal of the Hattori Botanical Laboratory* **46**: 1–41.
- Hansen, E. S., Poelt, J. & Søchting, U. (1987) Die Flechtengattung *Caloplaca* in Grönland. *Meddelelser om Grönland, Bioscience* **25**: 1–52.
- Hauck, M., Dulamsuren, C. & Mühlberg, M. (2007) Lichen diversity on steppe slopes in the northern Mongolian mountain taiga and its dependence on microclimate. *Flora* **202**: 530–546.
- Joshi, Y., Upreti, D. K. & Sati, S. C. (2008) Three new species of *Caloplaca* from India. *Lichenologist* **40**: 535–541.
- Kantvilas, G. & McCarthy, P. M. (2003) *Hueidea* (*Fuscideaceae*), a new lichen genus from alpine Australia. *Lichenologist* **35**: 397–407.
- Katoh, K., Kuma, K., Toh, H. & Miyata, T. (2002) MAFFT: a novel method for rapid multiple sequence

- alignment based on fast Fourier transform. *Nucleic Acids Research* **30**: 3059–3066.
- Kilius, H. (1985) *Cephalophysis* (Hertel) Kilius gen. nov., eine weitere Gattung der *Teloschistaceae* mit einzelligen Sporen. *Herzogia* **7**: 181–190.
- Kondratyuk, S. Y., Kärnefelt, I., Elix, J. A. & Thell, A. (2007) New species of the genus *Caloplaca* in Australia. *Bibliotheca Lichenologica* **95**: 341–386.
- Kondratyuk, S. Y., Kärnefelt, I., Elix, J. A. & Thell, A. (2009) Contributions to the *Teloschistaceae*, with particular reference to the Southern Hemisphere. *Bibliotheca Lichenologica* **100**: 207–282.
- Lederer, M. (1896) Einige für Bayern neue Flechten. *Berichte der Bayerischen Botanischen Gesellschaft* **4**: 26–30.
- Lendemer, J. C., Knudsen, K. & Fryday, A. M. (2010) New and interesting lichens, lichenicolous and allied fungi from Yosemite National Park, California, U.S.A. *Opuscula Philolichenum* **8**: 107–120.
- Magnusson, A. H. (1950) On some species of *Blastenia* and *Caloplaca* with black apothecia. *Botaniska Notiser* **1950**(3): 369–386.
- Meyer, B. & Printzen, C. (2000) Proposal for a standardized nomenclature and characterization of insoluble lichen pigments. *Lichenologist* **32**: 571–583.
- Muggia, L., Grube, M. & Tretiach, M. (2008) A combined molecular and morphological approach to species delimitation in black-fruited, endolithic *Caloplaca*: high genetic and low morphological diversity. *Mycological Research* **112**: 36–49.
- Nylander, J. A. A. (2004) *MrModeltest v2*. Program distributed by the author. Evolutionary Biology Centre, Uppsala University (<http://www.abc.se/wnylander/mrmodeltest2/mrmodeltest2.html>).
- Øvstedal, D. O. & Lewis Smith, R. I. (2001) *Lichens of Antarctica and South Georgia: A Guide to Their Identification and Ecology*. Cambridge: Cambridge University Press.
- Poelt, J. (1955) Mitteleuropäische Flechten III. *Mitteilungen der Botanischen Staatssammlung München* **2**(12): 46–56.
- Poelt, J. (1975) Mitteleuropäische Flechten X. *Mitteilungen der Botanischen Staatssammlung München* **12**: 1–32.
- Poelt, J. & Hafellner, J. (1980) *Apatopla* – genus novum Teloschistacearum (Lichenes). *Mitteilungen der Botanischen Staatssammlung München* **16**: 503–528.
- Ronquist, F. & Huelsenbeck, J. P. (2003) MrBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**: 1572–1574.
- Smith, C. W., Aptroot, A., Coppins, B. J., Fletcher, A., Gilbert, O. L., James, P. W. & Wolesey, P. A. (eds) (2009) *The Lichens of Great Britain and Ireland*. London: British Lichen Society.
- Søchting, U. (1973) Anatomical and cytological characteristics of unpigmented *Caloplaca verruculifera* from Denmark. *Botanisk Tidskrift* **68**: 152–156.
- Søchting, U. (1989) Lignicolous species of the lichen genus *Caloplaca* from Svalbard. *Opera Botanica* **100**: 241–257.
- Søchting, U. (1997) Two major anthraquinone chemosyndromes in *Teloschistaceae*. *Bibliotheca Lichenologica* **68**: 135–144.
- Søchting, U. (2001) Chemosyndromes with chlorinated anthraquinones in the lichen genus *Caloplaca*. *Bibliotheca Lichenologica* **78**: 395–404.
- Søchting, U. & Frödén, P. (2002) Chemosyndromes in the lichen genus *Teloschistes* (*Teloschistaceae*, Lecanorales). *Mycological Progress* **1**: 257–266.
- Søchting, U. & Øvstedal, D. O. (1998) *Caloplaca lewis-smithii*, a new lichen species from continental Antarctica. *Mycotaxon* **69**: 447–451.
- Søchting, U., Øvstedal, D. O. & Sancho, L. G. (2004) The lichens of Hurd Peninsula, Livingston Island, South Shetlands, Antarctica. *Bibliotheca Lichenologica* **88**: 607–658.
- Søchting, U., Lorentsen, L. B. & Arup, U. (2008) The lichen genus *Caloplaca* (Ascomycota, Lecanoromycetes) on Svalbard. Notes and additions. *Nova Hedwigia* **87**: 69–96.
- Solhaug, K. A., Gauslaa, Y., Nybakken, L. & Bilger, W. (2003) UV-induction of sun-screening pigments in lichens. *New Phytologist* **158**: 91–100.
- Šoun J., Vondrák J., Søchting U., Hrouzek P., Khodosovtsev A. & Arup, U. (2011) Taxonomy and phylogeny of the *Caloplaca cerina* group in Europe. *Lichenologist* **43**: 113–135.
- Tretiach, M. & Muggia, L. (2006) *Caloplaca badioreagens*, a new calcicolous, endolithic lichen from Italy. *Lichenologist* **38**: 223–229.
- Tretiach, M., Pinna, D. & Grube, M. (2003) *Caloplaca erodens* [sect. *Pyrenodesmia*], a new lichen species from Italy with an unusual thallus type. *Mycological Progress* **2**: 127–136.
- Vondrák, J., Khodosovtsev, A. & Říha, P. (2008) *Caloplaca concreticola* (*Teloschistaceae*), a new species from anthropogenic substrata in Eastern Europe. *Lichenologist* **40**: 97–104.
- Vondrák, J., Říha, P., Arup, U. & Søchting, U. (2009) The taxonomy of the *Caloplaca citrina* group (*Teloschistaceae*) in the Black Sea region; with contributions to the cryptic species concept in lichenology. *Lichenologist* **41**: 571–604.
- Vondrák, J., Khodosovtsev, A., Lőkös, L. & Merkulova, O. (2010a) The identity of type specimens in BP of some names in *Caloplaca*. *Mycotaxon* **111**: 241–250.
- Vondrák, J., Šoun, J., Sogaard, M., Søchting, U. & Arup, U. (2010b) *Caloplaca phlogina*, a lichen with two facies; an example of infraspecific variability resulting in the description of a redundant species. *Lichenologist* **42**: 685–692.
- Wetmore, C. M. (1994) The lichen genus *Caloplaca* in North and Central America with brown or black apothecia. *Mycologia* **86**: 813–838.
- Wetmore, C. M. (1996) The *Caloplaca sideritis* group in North and Central America. *Bryologist* **99**: 292–314.
- White, T. J., Bruns, T. D., Lee, S. & Taylor, J. (1990) Amplification and direct sequencing of fungal ribosomal DNA genes for phylogenies. In *PCR Protocols: a Guide to Methods and Applications* (M. A.

- Innis, D. H. Gelfand, J. J. Sninsky & T. J. White, eds): 315–322. San Diego: Academic Press.
- Wunder, H. (1974) Schwartzfrüchtige, saxicole Sippen der Gattung *Caloplaca* (Lichenes, *Teloschistaceae*) in Mitteleuropa, dem Mittelmeergebiet und Vorderasien. *Bibliotheca Lichenologica* **3**: 1–186.
- Xahidin, H., Abbas, A. & Wei, J.-C. (2010) *Caloplaca tianshanensis* (lichen-forming Ascomycota), a new species of subgenus *Pyrenodesmia* from China. *Mycotaxon* **114**: 1–6.